REVIEW BOARD ON THE USE OF HUMAN SUBJECTS, ICDUR, B.

ر در دانه

ation No.

obacted in

pal Investigator LOSEN GCASS Trainee Invostigator (if any)

80-010(P) Supporting Agency (if Non-ICDDR, B)

of Study Surveicepace of Project status: ( New Study HATLAS

) Continuation with change No change (do not fill out rest of form)

the appropriate answer to each of the following (If Not Applicable write NA).

rce of Population: Will signed consent form be required: Ill subjects

(a) From subjects Yes (No) Non-ill subjects **(b)** From parent or guardian

Minors or persons (if subjects are minors) Yes (No)

under guardianship No Will precautions be taken to protect is the study involve: anonymity of subjects

Physical risks to the Check documents being submitted herewith to subjects Yes Board:

Social Risks Umbreila proposal - Initially submit an Yes. Psychological risks

overview (all other requirements will to subjects be submitted with individual studies). Yes Discomfort to subjects (Yes) No Protocol (Required)

Invasion of privacy 'Abstract Summary (Required) Yes (No Disclosure of informa-Statement gaven or read to subjects on

rion damaging to subnature of study, risks, types of quest ions to be asked, and right to refuse Yes (No. ject be others is the study involve: to participate or withdraw (Required)

Use of records, (hosp-Informed consent form for subjects ital, medical, death, Informed consent form for parent or birth or other) (Yes) No guardian

Procedure for maintaining confidential-Use of fetal tissue or abortus Yes Use of organs or body Questionnaire or interview schedule A

Yes (No) If the final instrument is not completed subjects clearly informed about: prior to review, the following information

Nature and purposes of should be included in the abstract summary: stady A description of the areas to be Procedures to be covered in the questionnaire or

followed including interview which could be considered NO NA alternatives used Yes either sensitive or which would Physical risks Yes NO NA constitute an invasion of privacy.

NO NA Examples of the type of specific Sensitive questions Yes NO NA Benefits to be derived questions to be asked in the sensitive Yes Right to refuse to areas. participate or to with-An indication as to when the question-

draw from study NO NA naire will be presented to the Board Yes Confidencial handling. for review. of data Yes No NA

ment where there are risks or privacy is involved in . ony particular procedure NO NA Yes

se to obtain approval of the Review Board on the Use of Human Subjects for my changes ing the rights and welfare of subjects before making such change.

Trainee

avestigator

Compensation 6/or treat-

80-0/0 (P) Reed 30/1/80 Attachment la

#### PILOT PROJECT

#### SURVEILLANCE OF CAMPYLOBACTER IN MATLAB

#### ABSTRACT SUMMARY

Campylobacter has recently been identified as an important and previously unrecognized enteric pathogen in man. It is difficult to isolate in the laboratory without special, selective media and techniques and proper methods for field collection of specimens has Before a full protocol on Campylobacter not been fully determined. can be initiated in Matlab, isolation, transport, and preservation techniques must be worked out for a field hospital setting. pilot project deals with the determination of the optimal methods for transport of cultures from villages to the laboratory comparing four It further different transport techniques and media (Proposal A). would use the monthly surveillance of the children in the Diarrhea and Growth Study and a single survey of children in the Nutrition Growth Survey to determine the prevalence of Campylobacter among rural children and whether stool cultures or rectal swabs would have the highest yield (Proposal B).

The principal aim of this pilot project is, therefore, the development of laboratory capability to collect, transport, process, and identify <u>Campylobacter</u> in a field station so that a full-scale epidemiologic study could be performed later on.

## 1. Population

Proposal A would involve patients already registered in the routine hospital care network. Proposal B involves processing the routine rectal swabs taken from the Diarrhea and Growth Study as well as a routine rectal swab taken from patients in the Nutrition Growth Study.

### 2. Potentiál Risks

There are no significant risks to this study. Only rectal swabs and stool specimens will be collected.

# 3. Procedures for Minimizing Potential Risks

There are no risks, so this section is non-applicable.

# 4: Safeguarding Confidentiality

Confidentiality of the data collected will be insured. Patients names will not be used in analysis or publication of the data. In all laboratory studies, only the hospital number will be used to identify patients.

### 5. Informed Consent

A signed consent form will not be used since the study risks are minimal. Only routine diagnostic procedures will be done - rectal swab, stool collection, and interviewing. Information on diagnostic procedures will be provided verbally and verbal consent obtained. For minors, consent will be obtained from the authorized, legal guardian or parent of the child. The verbal statement made will include the nature and purpose of the procedures, the benefits derived, the right to refuse to participate, and the confidentiality of the data.

# 6. Interview Information

Patients found to have <u>Campylobacter</u> will be interviewed for their signs and symptoms, and history of animal and food exposure. The interview will take 10-15 minutes.

# 7. Potential Benefits to the Subject

The treatment of Campylobacter requires antibiotics different from the treatment of other diarrheal diseases. Patients identified to have this disease will receive proper treatment.

#### 8. Medical Records

This study requires use of Matlab hospital records.

# Pilot Protocol for the Study of <u>Campylobacter</u> Infection in Children

#### TRODUCTION

ates were 38 and 40%.

Campylobacter fetus ssp. jejuni is now recognized as a human enteric pathogen.

Sudies from all 6 inhabited continents have shown it to cause diarrheal illness

3-12% of the individuals studied.

The epidemiology of campylobacter enteritis is largely unknown. Most studies we been done in developed countries, however recent work in several less developed teas have also indicated its importance. A preliminary study of patients in Dacca th bloody diarrhea has shown <u>Campylobacter</u> to be the causative agent in 8% of uses; it is thus an important cause of diarrheal disease in Bangladesh. Similarly, tudies in South Africa, Brazil and Zaire have shown it to be a common cause for larrheal illness.

In North America and Europe carriage by asymptomatic adults is very infrequent.

In North America and Europe carriage by asymptomatic adults is very infrequent.

In North America and Europe carriage by asymptomatic adults is very infrequent.

In North America and Europe carriage by asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asymptomatic children carried C. fetus

Provides in Belgium however showed that 1.3% of asy

Taken together, these studies suggest that children, especially in less eveloped areas, may have a high prevelence of infection with <u>C. fetus ssp. jejuni</u>; afection may be either clinical or asymptomatic. Further clarification of this uestion is indicated.

However before such a study could be carried out examination of the best athods for transport of specimens from villages to the laboratory need to be established. Thus this proposal has two parts.

#### ROPOSAL A

In order to determine whether <u>Campylobacter fetus</u> ssp. <u>jejumi</u> is a significant cause of diarrheal disease in rural Bangladesh it is necessary to establish a apability for culturing stools for that organism. A pilot study at the outpatient reatment center has indicated that use of selective methods will detect <u>Campylobacter</u> in fecal specimens. A similar system employing antibiotic-containing media, incubation at 42°C and a microaerophilic atmosphere could be instituted at the laboratory in Matlab. Patients with diarrhea attending the clinic would have stool cultures obtained for routine pathogens (i.e. <u>Shigella</u>) and for <u>Campylobacter</u>. Approximately 100 cases should be cultured and if <u>Campylobacter</u> is isolated, then a total of 25 positive cases should be collected.

Each of these cases would be visited in their village and repeat cultures obtained. Repeat cultures will be transported to the laboratory in 1) Carey-Blair media 2) Campy-thio media 3) fresh stools 4) and after being inoculated to selective plates at the village and then transported back to the laboratory in candle jars. Using these positive patients we could determine which are the best methods for transport of specimens. Whenever possible, family contacts and household animals would also be cultured, especially if they had a recent history of a diarrheal episod

#### PROPOSAL B

The population to be studied would be a random sample of children under the age of 5. The total number of children to be examined would be 500. All children enrolled in this study would have a stool specimen or a rectal swab taken. Swabs would be transported to the laboratory at the central hospital in Matlab using the methods shown to be most effective in Part A, and cultured for Campylobacter fetus ssp. jejumi. The parent of each child would be asked about age and sex of the child and whether the child had had recent diarrheal illness.

A field study would then be undertaken in the villages surrounding Matlab bazaar

Those children who were positive would then be cultured three times within the next week, comparing the efficacy of fresh stools versus rectal swabs.

This pilot study would determine prevelence of <u>Campylobacter</u> infection among and children, and whether stool cultures or rectal swabs would have the highest yield. If <u>Campylobacter</u> is found in these children, further studies could be designed to determine the duration of excretion, and epidemiological and clinical features of <u>Campylobacter</u> infection.

#### CONSENT FORM

# COLLECTION OF REPEAT STOOL SPECIMENS FROM CAMPYLOBACTER PATIENTS IDENTIFIED IN HOSPITAL

#### Proposal A

You recently were hospitalized for diarrhea and had <u>Campylobacter</u>, a newly identified bacteria, isolated from your stool. If you are not already better, we can now prescribe further medication for your illness. To determine if whether you are still infected with this organism, we would like to reculture your stool. You may decide whether or not you wish to allow us to collect this specimen.

For purposes of this study, you will not be identified by name.

#### Proposal B

We would like to perform a rectal swab and a stool culture on your child to search for <u>Campylobacter</u>, a newly identified bacteria which causes diarrhea as well as other known diarrhea-causing organisms. We will also ask about your child's health in the past week. If your child is ill, a physician will see the child and prescribe appropriate treatment.

Your participation will help the Cholera Hospital develop better methods to diagnose diarrheal illness in the future. You may choose not to allow your child to be cultured. For purposes of this study, your child will not be identified by name.