

Importance of Age and Sociodemographic Factors in Contraceptive Acceptance Among Rural Women In Bangladesh: Lessons Learned from Matlab MCH-FP Project

J Chakraborty, Andres de Francisco, Shamim A Khan, and K Zaman

Objective: Compare selected sociodemographic variables of first-time acceptors of modern contraceptive methods with those of never-users during the same period.

Methods: Data for this study came from the ICDDR,B's Maternal and Child Health-Family Planning (MCH-FP) Project in Matlab, Bangladesh. In the project area, improved health and family planning services are provided and relevant data are collected. Sociodemographic information on women's age, parity, education, and occupation, the size of dwelling house and its construction material was recorded between 1984 and 1993 for contraceptive acceptors on the date of acceptance. This information was compared with data on non-users after the first pregnancy outcome date.

Results: Contraceptive prevalence rates in the Matlab MCH-FP Project area increased from 40% in 1985 to 63% in 1993. Modern methods accounted for almost all contraceptive use (97%), with injectables having the highest prevalence (52%) followed by oral contraceptives (30%) and female sterilization (10%). Eight thousand six hundred and fifty-one women aged less than 29 years constituted the largest group of new acceptors (80% of new users) and the number of those over 34 years of age was 852 who constituted the lowest group (8%). Among non-users, 3,950 (74%) were below 29 years and 940 (18%) above 35 years. Injectable contraceptives were the most popular in all ages. Highest oral contraceptive use was observed among women aged 28 years and below (22%). The data showed no significant difference of contraceptive acceptance among educated and non-educated group. No association was observed between contraceptive prevalence rate and household area or type.

Conclusions: This study suggests that the most appropriate group of women to be targeted for family planning in rural Bangladesh consists of the youngest group of women who are easy to reach and demographically important.



Service Delivery at Family Welfare Centres: the Clients' Perspective

Parveen A Khanam, Helene Wirzba, Tanjina Mirza, and Therese Juncker

The Family Welfare Centre (FWC) are the primary fixed centres for the provision of maternal and child health and family planning services at grass root level. Two paramedics are posted of which one is female. The objectives of this study was to evaluate the types and quality of services and determine client satisfaction of these services. Six hundred and fifty exit interviews were conducted with FWC clients using a structured questionnaire between July and September 1993 in Abhoynagar and Sirajgonj Thanas, the field sites of the MCH-FP Extension Project (Rural) of ICDDR,B. Client were asked about the services they had received that day.

Most clients were married women, aged 20 to 40 years, living less than one kilometre away from FWC. They had an average of 3.75 previous pregnancies and 55 percent of them were using a contraceptive method. At least one paramedics was present on 94 percent of the working days while interviews were conducted at the FWCs. Illness of the women (68%) and child care (51%) were the most common reasons for seeking services. Only 8 percent clients received antenatal care while 6 percent got family planning services. Over ninety percent clients received at-least two drugs that day. Half of the family planning complications were found among the IUD users, who make up only 6% of all users. Discrepancies were found between the drugs recorded in the register and those actually dispensed. Many clients did not remember or were not told the dosage of drugs. A quarter of the clients received antibiotics of which 70% percent did not know the proper dosage of drugs. Thirty percent of the clients were not satisfied about privacy, clinical examination, availability of drugs, and counselling. Their knowledge about services offered at FWC indicates