


Complications during labour and delivery in health facilities in rural Bangladesh

Globally, each year, an estimated 358,000 women die because of complications during pregnancy and childbirth^{1,2}. The absolute figure of maternal mortality is only the tip of the iceberg; behind each death lies a huge burden of life-threatening and chronic morbidities³⁻⁵. Globally, 7 to 8% of pregnant women suffer acute maternal complications resulting in about 9 million morbid events each year^{2,6}. In Bangladesh an estimated 6,000 women die from pregnancy-related complications every year⁷ while another 194,000 women reportedly suffer injuries or disabilities caused by complications during pregnancy and childbirth⁸.

Despite several national estimates on maternal mortality, there is a dearth of reliable data on levels and types of maternal morbidities. Available studies document self-

deliver in a health facility¹¹. Facility records of all women admitted during labour or up to 42 days postpartum to any public or private hospital in Matlab Upazilla and the district town Chandpur were reviewed by a physician to identify acute maternal complications. Maternal complications were categorized into four groups based on the severity of the complication type (very severe maternal complication, less severe maternal complication, caesarean section (C/S) with no maternal complication, and vaginal delivery with no maternal complication). The definition of the complication categories are shown in Figure 1. Of the total 4,817 pregnant women during the 24-month study period, 44% delivered in one of the 30 local hospitals and their hospital records were reviewed for classification into one of the 4 categories.


Figure 1: Categorization of different maternal complications



reported pregnancy-related complications which is not medically valid^{9,10}. To address this knowledge gap, between January 2007 and December 2008, a prospective study was conducted to determine the levels and types of acute maternal morbidities in the Matlab icddr,b service area, a rural area of Bangladesh where 81% of women


Research Findings:

- Among the women who delivered in local hospitals, 7% were identified with very severe and 15% with less severe maternal complications. 7% had a caesarean section with no maternal complication and 63% had a vaginal delivery without any reported maternal



complication. 8% of the hospital records were not found.

- Among women with very severe or less severe maternal complications, the majority (69%) had C/S followed by hypertensive disorders of pregnancy (13%), haemorrhage (8%), anaemia (6%) and infection (4%).
- The majority (75%) of the women with very severe maternal complications and C/S with no maternal



indications were treated in private hospitals in Chandpur and women who had a vaginal delivery without any maternal complication were mostly managed in the Matlab icddr,b hospital (Figure 2).

- A total of 401 caesarean sections (19%) were recorded. The private hospitals performed nine times as many caesarean sections compared to public hospitals of Matlab and Chandpur. Indications for caesarean sections are shown in Figure 3.

Conclusions:

- If the incidence of severe obstetric complications of 7% found in this study holds true for all women of Bangladesh, then an estimated 124,264 severe obstetric complications occur in facilities annually nationwide.
- Most women with pregnancy complications seek care from private facilities and the public facilities remain underutilized.
- Only one-quarter of all C/S deliveries currently conducted are actually needed for saving lives of mothers and another one-quarter is needed for saving babies.
- A large proportion of unnecessary C/S deliveries are conducted in private facilities. For every one in eight C/S deliveries, no indication is recorded by the provider.

Recommendations:

- The reasons behind the failure of public sector to serve women with pregnancy complications needs to be investigated
- A system to monitor the indications for caesarean sections in private facilities to avoid unnecessary surgical interference is required to improve recording indications.

Addressing these recommendations will help to improve not only the safe motherhood program in Matlab but also overall maternal health services in hospitals in Bangladesh.

- Trends in maternal Mortality: 1990 to 2008. Estimates developed by WHO, UNICEF, UNFPA and The World Bank. World Health Organization 2010.
- Hogan MC, Foreman KJ, Naghavi M, Ahn SY, Wang M, Makela SM, Lopez AD, Lozano R, Murray CJL. Maternal mortality for 181 countries, 1980–2008: a systematic analysis of progress towards Millennium Development Goal 5. Lancet 2010; 375: 1609-23.
- BETTER OFF DEAD? A report on maternal morbidity from the UK All Party Parliamentary Group on Population, Development and Reproductive Health. May 2009.
- The World Bank Group. Millennium development goals. 2000. <http://www.developmentgoals.org> (access on January 10, 2010).
- UNICEF, WHO, UNFPA. 1997. Guidelines for monitoring the availability and use of obstetric services. New York: United Nations Population Fund.
- Filippi V, Ronsmans C, Campbell OMR, Graham WJ, Mills A, Borghi J, Koblinsky M, Osrin D. Maternal health in poor countries: the broader context and a call for action. www.thelancet.com. September 28, 2006.
- Demographics of Bangladesh, 2009 estimates. http://en.wikipedia.org/wiki/Demographics_of_Bangladesh (Access on 14.06.2011).
- Bangladesh Maternal Mortality and Health Care Survey 2010: Summary of Key Findings and Implications. National Institute of Population Research and Training (NIPORT), MEASURE Evaluation, UNC-CH, USA, ICDDR,B.
- Filippi V, C. Ronsmans, et al. (2000). "Women's reports of severe (near-miss) obstetric complications in Benin." Stud Fam Plann 31 (4): 309-24.
- MotherCare Matters 1997,6:15-16; Washington, DC: JSI
- Rahman. A. et al. Effectiveness of an integrated approach to reduce perinatal mortality: Recent experiences from Matlab, Bangladesh. Unpublished data, 2011.

This Knowledge Translation Brief is based on the manuscript by Huda FA, Ahmed A, DasGupta SK, Jahan M, Ferdous J, Koblinsky M, Ronsmans C, Chowdhury ME. Profile of maternal and foetal complications during labour and delivery among women giving birth in hospitals in Matlab and Chandpur, Bangladesh. JHPN-1005:1583-MM (Accepted for publication on 24.1.11). The study was funded by USAID. For more information, please contact Dr. Fauzia Akhter Huda, Senior Research Investigator, Centre for Reproductive Health, icddr,b, Mohakhali, Dhaka-1212, Bangladesh. Email:fauzia@icddr.org Tel: 8860523-32 Ext 2257, Cell: 01716-962841.

This is the product of a grant to icddr,b from the Maternal Health Task Force (MHTF) at EngenderHealth. The views expressed in this brief are not necessarily those of the MHTF and full responsibility for all contents remains with the author(s) of the Knowledge Translation brief.