

Community Support Group: a sociable approach for improving maternal health in rural Bangladesh

Improving maternal health remains a significant challenge in Bangladesh. Lack of awareness and socio-cultural and religious taboos, among other factors, contribute to the current maternal mortality rate of 194 per 100,000 live births in Bangladesh [1]. One evidence-based intervention [2, 3] to increase awareness of the causes of maternal mortality and to encourage advocacy that will improve maternal health outcomes is social mobilization through community support groups (CSGs). As part of the demand side of a larger maternal and neonatal health

local health facilities and health service providers (family welfare assistants, family welfare visitors, health assistants, family planning inspectors, and most importantly the community-based skilled birth attendants (CSBAs)), to ensure the availability of maternal health care services when required. The effectiveness of CSGs was assessed using a baseline and a follow-up structured questionnaire survey during November'08 to March'09 and October'10 to November'10 among mothers who delivered six months prior to the interview date.



Advocacy meeting for the formation of CSG



CSG volunteers in a courtyard-session

(MNH) intervention, icddr,b formed CSGs by involving community members in Shahjadpur sub-district. The objectives of the CSGs are: to identify pregnant women and sensitize pregnant women and members of her family about the importance of using skilled maternal care; to provide support to the poor and disadvantaged through fund raising and arranging transport to a facility for mothers with complications; and to establish links between the community and the facility.

Approach:

By October 2010, 64 CSGs had been formed. Each CSG was comprised of 30 committee members serving in one of three different capacities - an advisory committee, executive committee and volunteer committee. The aim of the committee members was to persuade pregnant mothers and their family members to communicate with

Results:

In the baseline and follow-up surveys, 3,158 and 2,725 mothers were recruited respectively for interview. Following the baseline survey, at least 640 courtyard-sessions of CSGs were conducted by female volunteers covering 6,500 mothers. During the baseline survey nearly three-fourths of the mothers (70%) reported receiving any antenatal care (ANC); which increased to 84% after formation of CSGs and covering almost all the unions of Shahjadpur sub-district. Skilled attendance at delivery was 26% at baseline and increased to 42% at the follow-up survey. Mothers who attended the courtyard-sessions conducted by CSGs during the time period received substantially more ANC and skilled delivery care than the mothers who did not attend (**Figures 1 and 2**).




Figure 1: Number of ANC visits by attendance in courtyard CSG session




Figure 2: Percentage of mothers receiving skilled delivery care by attendance in courtyard CSG session

Conclusion:

The intervention findings suggest that the courtyard-sessions using CSGs made noticeable contribution to improving maternal health-seeking behaviour in terms of ANC visits and skilled attendance at delivery.

Implications:

Participatory approaches through courtyard-sessions and family involvement in care stimulated by the CSGs appear effective in improving maternal health-seeking behaviour in Bangladesh. There was some operational cost (US\$ 225 in 2010) to form each CSG and provide hand-outs for training to the volunteer committee. However, involvement of elite groups of villagers in the CSGs may increase the likelihood that the program will be sustainable in the long run.

References:

1. National Institute of Population Research and Training (NIPORT), ORC Macro, Johns Hopkins University and ICDDR,B. Bangladesh Maternal Health Services and Maternal Mortality Survey 2010. Dhaka, Bangladesh 2011.
2. UNICEF. Maternal and Neonatal Health in Bangladesh. January 2009.
3. 3ie Enduring Questions Brief. Access to health: How to reduce child and maternal mortality? International Initiative for Impact Evaluation. Number 14, June 2010. 3ie, Global Development Network, Second Floor, East Wing, ISID Complex, Plot No.4, Vasant Kunj Institutional Area, New Delhi 110 070. www.3ieimpact.org.

This Knowledge Translation Brief was prepared by Anisuddin Ahmed, Nafis Al Haque, Nafisa Lira Huq, Sadika Akther, Sheikh A Shahed Hossain, Laura Reichenbach, Moyazzam Hossain, Md. Jamal Uddin, M A Quaiyum. It is based on the baseline and follow-up evaluation of the Shahjapur Integrated Maternal and Neonatal Intervention Project (SIMNHIP), being implemented by Centre for Reproductive Health of icddr,b at Shahjapur sub-district of Sirajganj district, Bangladesh. The study was funded by AusAid.

For more information, please contact Mr. Anisuddin Ahmed, Senior Statistical Officer, Centre for Reproductive Health, icddr,b, Mohakhali, Dhaka-1212, Bangladesh. Email: anisuddin@icddr.org.

Tel: 8860523-32 Ext 2232, Cell: 01917-618005.

This is the product of a grant to icddr,b from the Maternal Health Task Force (MHTF) at EngenderHealth. The views expressed in this brief are not necessarily those of the MHTF and full responsibility for all contents remains with the author(s) of the Knowledge Translation brief.

Recommendations:

- Formation of CSGs over the entire sub-district to increase community awareness regarding appropriate maternal health actions;
- Encourage CSGs to be self-sustainable and to take independent initiatives to address a range of maternal and neonatal health issues;
- Birth preparedness initiatives should first empower household members through CSGs; and
- To further address the gap between knowledge and action, CSGs should also focus on arrangement of transport and blood donors, to help limit potential delays in referral to facility.