DEMOGRAPHIC SURVEILLANCE SYSTEM— MATLAB

VOLUME TWENTY FOUR

REGISTRATION OF DEMOGRAPHIC EVENTS-1993

> SCIENTIFIC REPORT NO. 76 May 1996



INTERNATIONAL CENTRE FOR DIARRHOEAL DISEASE RESEARCH, BANGLADESH

DEMOGRAPHIC SURVEILLANCE SYSTEM - MATLAB Volume Twenty Four Registration of Demographic Events - 1993

With a Special Supplement

Fertility Trends in Matlab by Birth Order, Age at Birth and Birth Interval, 1980-1993



International Centre for Diarrhoeal Disease Research, Bangladesh DSS reports are not copyrighted and may be freely quoted as long as the source is properly indicated. The following citation is suggested for this report; this form of citation is also appropriate for previous DSS annual reports.

ICDDR,B. <u>Demographic Surveillance System - Matlab: Registration of Demographic Events - 1993</u>. Dhaka, Bangladesh: 1996.

This report was prepared by the staff of the Demographic Surveillance System, assisted by Dr. John Blacker, of the London School of Hygiene and Tropical Medicine.

ISBN 984-551-055-8

Cover design: Asem Ansari

May 1996

Scientific Report No. 76

Published by:

International Centre for Diarrhoeal Disease Research, Bangladesh GPO Box 128, Dhaka 1000, Bangladesh

Telephone: 871751-60 (10 lines); Cable: CHOLERA DHAKA; Telex: 675612 ICDD BJ; Fax: 880-2-883116 and 880-2-886050

PREFACE

The International Centre for Diarrhoeal Disease Research, Bangladesh (ICDDR,B) is an autonomous, international, philanthropic, non-profit centre for research, education, training, and clinical service. The Centre is derived from the Cholera Research Laboratory (CRL). Its aims and objectives are to undertake and promote study, research, and dissemination of knowledge in diarrhoeal diseases and the directly related subjects of nutrition and fertility, with a view to developing improved health care methods and to prevent and control diarrhoeal diseases and improve public health programmes, especially in developing countries.

The ICDDR,B issues an annual report, working papers, scientific reports, special publications, monographs, theses, dissertations, an international journal on diarrhoeal diseases, and a bi-monthly newsletter which demonstrates the type of research activities currently in progress. The views expressed in these publications are those of the authors, and do not necessarily represent the views of the ICDDR,B.

The Centre has two major objectives:

- 1. To undertake and promote study, research, and dissemination of knowledge in diarrhoeal diseases and directly related subjects of nutrition and fertility with a view to developing improved methods of health care and the prevention and control of diarrhoeal diseases and improvement of public health programmes with special relevance to developing countries.
- 2. To provide facilities for training to Bangladeshi and other nationals in areas of the Centre's competence in collaboration with national and international institutions.

ACKNOWLEDGEMENTS

The Demographic Surveillance System (DSS) of the International Centre for Diarrhoeal Disease Research, Bangladesh (ICDDR,B) during the past few years received financial support from UNFPA, the Overseas Development Administration (ODA) of the United Kingdom, the Netherlands government, and ICDDR,B. The ICDDR,B is supported by countries and agencies which share its concern for the health problems of developing countries. Current donors include: the aid agencies of the Governments of Australia, Bangladesh, Belgium, Canada, China, Denmark, Germany, Japan, the Netherlands, Norway, Republic of Korea, Saudi Arabia, Sri Lanka, Sweden, Switzerland, Thailand, the United Kingdom, and the United States; international organizations, including Arab Gulf Fund, Asian Development Bank, European Union, the United Nations Children's Fund (UNICEF), the United Nations Development Programme (UNDP), the United Nations Population Fund (UNFPA), and the World Health Organization (WHO); private foundations including Aga Khan Foundation, Child Health Foundation, Ford Foundation, Population Council, Rockefeller Foundation and the Sasakawa Foundation; and private organizations including American Express Bank, Bayer A.G., CARE, Family Health International, Helen Keller International, the Johns Hopkins University, Macro International, New England Medical Centre, Procter Gamble, RAND Corporation, SANDOZ, Swiss Red Cross, and the University of Alabama at Birmingham, the University of Iowa, and others.

Publication of this Report was made possible by grant number 514 "Updating the Demographic Surveillance System", by the Overseas Development Administration of the United Kingdom. The ICDDR,B is grateful for this generous grant which funded the completion of this report, including the provision of the services of Dr. John Blacker, who improved the content and accuracy of this report and ensured its rapid completion. Mr. Golam Mostafa prepared the initial draft; and the layout design and desktop publishing was done by Mr. Sentu B. Gomes.

CONTENTS

| | | | | Page |
|---|-----|----------|-------|------|
| Summary | | 1333 | 2.51 | 1 |
| Chapter 1 : Introduction | | | | 2 |
| Chapter 2 : Population Changes | *** | *** | | 5 |
| Chapter 3 : Mortality | | | *** | 14 |
| Chapter 4 : Fertility | 202 | | *** | 34 |
| Chapter 5 : Marriage and Divorce | | 111 | | 45 |
| Chapter 6 : Migration | *** | *** | | 54 |
| Chapter 7 : Special Supplement: Fertility Trends in Matlab Order, Age at Birth and Birth Interval, 1980-19 | | th | 224 | 69 |
| LIST OF TABLES | | | | |
| Table 2.1 : Vital Statistics of the Matlab MCH-FP and Comparison Areas, 1982-1993 | | *** | *** | 6 |
| Table 2.2 : Mid-year Population, Events Registered, and Population Changes in 1993 | | 1,525 | 500.0 | 8 |
| Table 2.3 : Mid-year Population by Age and Sex, 1993 | | | 144 | 9 |
| Table 2.4 : Mid-year Population by Area, Age, and Sex, 19 | 993 | | | 10 |
| Table 2.5 : Mid-year Population in MCH-FP Area by Age, Sex, and Block, 1993 | *** | (2.5.5) | 222 | 11 |
| Table 3.1 : Deaths by Age and Sex, 1993 | | | | 16 |
| Table 3.2 : Deaths by Area, Age, and Sex, 1993 | | | *** | 17 |
| Table 3.3 : Deaths in MCH-FP Area by Age, Sex, and Block, 1993 | *** | | 111 | 18 |

LIST OF TABLES (continued)

| | | | | Page |
|--|------|-----|------|------|
| Table 3.4 : Death Rates by Age and Sex, 1993 | | *** | | 20 |
| Table 3.5 : Death Rates by Area, Age, and Sex, 1993 | *** | *** | *** | 21 |
| Table 3.6 : Abridged Life Table, 1993 | 22.5 | *** | 200) | 23 |
| Table 3.7 : Abridged Life Tables by Sex, 1993 | ••• | | ••• | 24 |
| Table 3.8 : Abridged Life Tables by Area, 1993 | | | | 25 |
| Table 3.9 : Abridged Life Tables for MCH-FP Area by Sex, 1993 | *** | | ••• | 26 |
| Table 3.10: Abridged Life Tables for Comparison Area by Sex, 1993 | *** | | *** | 27 |
| Table 3.11: Deaths by Age and Month, 1993 | | | | 28 |
| Table 3.12: Male Deaths by Cause and Age, 1993 | | *** | *** | 29 |
| Table 3.13: Female Deaths by Cause and Age, 1993 | ••• | 534 | 200 | 30 |
| Table 3.14: Male Deaths by Cause, Age, and Area, 1993 | | | | 31 |
| Table 3.15: Female Deaths by Cause, Age, and Area, 199 | 3 | | *** | 32 |
| Table 3.16: Age-standardized Mortality Rates by Cause of Death, 1993 | | *** | ••• | 33 |
| Table 4.1 : Number and Rates of Pregnancy Outcomes by Type and Area, 1993 | 2.2 | | | 35 |
| Table 4.2 : Pregnancy Outcomes by Month, 1993 | ••• | *** | *** | 36 |
| Table 4.3 : Age-specific Fertility Rates and Indices, 1993 | *** | *** | | 38 |
| Table 4.4 : Age-specific Fertility Rates and Indices by Area, 1993 | *** | | | 39 |

LIST OF TABLES (continued)

| | | | | | Page |
|---------------|---|-------|-------|-----|------|
| | Age-specific Fertility Rates and Indices for MCH-FP Area by Block, 1993 | | 275 | | 41 |
| | Births by Mother's Age, Live-birth Order and Area, 1993 | | *** | | 43 |
| Table 4.7 : A | Age-order-specific Fertility Rates by Area, 1993 | | **** | | 44 |
| | Groom's Age at Marriage by Previous Marital Status, 1993 | | *** | | 46 |
| | Bride's Age at Marriage by Previous Marital Status, 1993 | | *** | | 47 |
| Table 5.3 : I | Marriage Rates by Age and Sex, 1993 | | *** | *** | 48 |
| | Number of Marriages by Groom's and Bride's Age at Marriage, 1993 | 122 | | | 49 |
| | Number of Divorces by Partners' Age at Divorce, 1993 | *** | (***) | | 50 |
| Table 5.6 : I | Marriages and Divorces by Month, 1993 | 132 | *** | | 51 |
| | Number of Divorces by Sex, Age, and Duration of Marriage, 1993 | *** | *** | 737 | 53 |
| Table 6.1 : l | In- and Out-migration by Age and Sex, 1993 | *** | *** | | 56 |
| Table 6.2 : | In-migration by Age, Sex, and Area, 1993 | | | *** | 57 |
| Table 6.3 : | Out-migration by Age, Sex, and Area, 1993 | *** | | | 58 |
| | Age and Sex-specific Migration Rates by Direction, 1993 | *** | 200 | *** | 59 |
| | Male Out-migration by Cause of Movement and Age, 1993 | (555) | | *** | 61 |

LIST OF TABLES (continued)

| | | | | <u>Page</u> |
|--|-----|-------|------|-------------|
| Table 6.6 : Female Out-migration by Cause of Movement and Age, 1993 | | ••• | | 62 |
| Table 6.7 : Male In-migration by Cause of Movement and Age, 1993 | | | | 63 |
| Table 6.8 : Female In-migration by Cause of Movement and Age, 1993 | | (555) | | 64 |
| Table 6.9: In- and Out-migration by Sex and Month, 1993 | | | | 65 |
| Table 6.10: Male Migration by Destination or Origin, 1993 | *** | *** | ••• | 67 |
| Table 6.11: Female Migration by Destination or Origin, 199 | 3 | | *** | 68 |
| Table S.1 : Total Order-specific Fertility Rates by Area, 1980-1993 | *** | | ••• | 74 |
| Table S.2 : Period Parity Progression Ratios for Matlab by Area, 1980-1993 | | *** | 9990 | 75 |
| Table S.3 : Mean Age of Mothers at the Birth of their Children by Birth Order, 1980-1993 | | 224 | *** | 76 |
| Table S.4: Mean Birth Intervals in Months by Birth Order, 1984-1993 | | *** | | 77 |
| LIST OF FIGURES | | | | |
| Fig. 1.1 : Map of Bangladesh Showing the Study Area | *** | 95 | *** | 3 |
| Fig. 1.2 : Matlab Area Showing Villages of Demographic Surveillance System, 1993 | | 127 | 9000 | 4 |
| Fig. 2.1 : Trends in Fertility and Under Five Mortality by Area, 1982-1993 | *** | nu. | | 7 |
| Fig. 2.2 : Age Pyramid of the 1993 Mid-year Population | 222 | ••• | *** | 13 |

LIST OF FIGURES (continued)

| | | | | | Page |
|------------------------|---|-----------------|-----------|-------------------|------|
| Fig. 3.1 : P A | robability of Survival from Birth to ge(x) by Sex, 1993 | pim-lu 249 i | O slor | maid of brain. | 22 |
| Fig. 4.1 ; N | umber of Births and Deaths by Month, 1993 | | | | 37 |
| Fig. 4.2 : A | ge-specific Fertility Rates by Area, 1993 | £901 | AGEA | DITE. | 40 |
| Fig. 5.1 : M | arriages and Divorces by Month, 1993 | | Sec | | 52 |
| Fig. 6.1 : R | ate of In- and Out-migration by Sex and | 217-14 | | | 60 |
| Fig. 6.2 : N | umber of In- and Out-migrants by Sex and | | 실세 벨 | da sa | 00 |
| M M | | 1725 | 10,511 | e=1.58 | 66 |
| Fig. S.1 : R 19 | elative Change in PPR's (1980=1) by Area, 980-93 | iq 0-10 | 90 to: | | 78 |
| Fig. S.2 : R 19 | elative Change in Age at Birth by Area, 880-93 | | 9 kyh | | 79 |
| Fig. S.3 : Ti | rends in Birth Interval by Order by Area, 980-93 | d lo s | | M . č.) | 80 |
| | LIST OF APPENDICES | | | | |
| Appendix A: | Names and Codes of Villages in the DSS Area, 1993 | - | 620-3 | 2027 | 81 |
| Appendix B: | Mid-year Population, Births, and Deaths by Village, 1993 | usup. | | | 82 |
| Appendix C: | Life Table Equations | 1880 | MAI etti) | LUCA . | 86 |
| | New Standard Populations | 561 | (UNIVAL) | | 87 |
| | Staff of the DSS, 1993 | | Minute | | 88 |
| NO VARIO DI MANGAMBARA | | .,,, | | | 00 |

SUMMARY

This report presents the vital registration data for events taking place in 1993 in Matlab, Bangladesh. These data were collected by the Demographic Surveillance System of the International Center for Diarrhoeal Disease Research, Bangladesh. The surveillance area is divided into a Maternal and Child Health and Family Planning (MCH-FP) intervention area and a Comparison area receiving government services.

In 1993 fertility continued to fall in the MCH-FP area; the crude birth rate declined to 24.7 per thousand and the total fertility rate to 2.9 birth per women. Fertility also fell in the Comparison area, where the crude birth rate was 29.4 per thousand and total fertility rate 3.8 birth per women.

The mortality pattern showed a different picture from fertility. In the MCH-FP area crude death rate dropped to 7.7 per 1,000 and in the Comparison area the crude death rate rose to 10.2. In the MCH-FP area the infant mortality showed a sharp decline from 80.5 in 1992 to 63.1 per 1,000 live births in 1993. In the Comparison area the infant mortality rate rose from 90.2 in 1992 to 99.3. In the MCH-FP area under five mortality was 86.1 per thousand live births, while the corresponding figure for the Comparison area was 135.1 per thousand.

Rates of both in- and out-migration for the surveillance area as a whole decreased in 1993, with in-migration at 25.5 per thousand and out-migration at 36.1 per thousand, leaving a net out-migration of 10.6 per thousand, thus offsetting the rate of natural increase, and reducing the overall rate of population growth to 0.8 percent per annum.

This report also presents a Special Supplement describing fertility trends in Matlab by birth order, age at birth and birth interval.


CHAPTER 1

INTRODUCTION

Since 1963 the International Center for Diarrhoeal Disease Research, Bangladesh (ICDDR,B), formerly the Cholera Research Laboratory, has been conducting a health related research program near the town of Matlab, in rural Bangladesh. Matlab is located about 55 kilometers southeast of the country's capital, Dhaka (Figure 1.1). The Demographic Surveillance System (DSS) is one of the components of this field program. Since 1966 the DSS has maintained the registration of births, deaths, and migrations, in addition to carrying out occasional censuses. In 1975 the system was augmented to include marriages and divorces. This information is gathered by Community Health Workers and Health Assistants who visit each household in their assigned areas regularly and fill out the event registration forms. A detailed description of the DSS and its operation appears in CRL Scientific Report No. 9 (March 1978). In October 1977 the surveillance area was reduced from 233 to 149 villages and a Maternal Child Health and Family Planning (MCH-FP) Program was begun in 70 villages. The remaining 79 villages were treated as a comparison area (Figure 1.2). These changes are described in detail in the ICDDR,B Scientific Report No. 47 (May 1981).


This is the twenty-fourth volume of a series of scientific reports of the Demographic Surveillance System produced by the ICDDR,B. Presented here are results obtained from the Matlab DSS in 1993, along with brief notes and explanations of the tables. This volume also includes a special supplement describing fertility trends in Matlab by birth order, age at birth and birth interval.

Figure 1.1: Map of Bangladesh Showing the Study Area



Key: Study areas

Figure 1.2: Matlab Area Showing Villages of Demographic Surveillance System, 1993



CHAPTER 2

POPULATION CHANGES

Table 2.1 summarizes the principal vital statistics of the MCH-FP and Comparison areas from 1982 through 1993, and the basic 1993 figures, by sex, are shown in Table 2.2. The 1993 fertility rates declined in comparison with those of the preceding year in the MCH-FP area, with a total fertility rate of 2.9 and a crude birth rate of 24.7 per thousand in both years. In the Comparison area the TFR fell from 4.0 to 3.8 and the crude birth rate from 31.1 to 29.4 per thousand. The trends in the total fertility rate in both areas are illustrated in Figure 2.1(a).

Infant mortality declined sharply in the MCH-FP area, 80.5 to 63.1 from 1992 to 1993. This decrease was the net result of decrease in both the neonatal mortality and the post-neonatal mortality rate. In the Comparison area infant mortality increased from 90.2 to 99.3 per thousand from 1992 to 1993. This was the net result of an increase in the neonatal mortality and a decrease in the post-neonatal mortality rate.

Mortality of children (1-4) remained constant in comparison with that of the preceding year in MCH-FP area, but in the Comparison area showed little change, decreasing by 0.4 from 10.4 to 10.0 from 1992 to 1993.

The trends in under-five mortality are illustrated in Figure 2.1(b). Migration figures, both in and out, of the surveillance area as a whole decreased in 1993. Out- migrants continued to outnumber in-migrants, thus offsetting the rate of natural increase, which amounted to 18.1 per thousand in 1993, so that the overall rate of population growth was reduced to 0.8 percent per annum.

Tables 2.3, 2.4, and 2.5 show the age and sex distributions for the whole study area, the MCH-FP and Comparison areas, and for the four blocks of the MCH-FP area. The age-sex distribution for the study area is illustrated by the population pyramid shown in Figure 2.2. The decline of fertility in the MCH-FP area has caused a small but significant change in the age structure of the population. Children under 15 years of age constituted 43.4 percent of the population at the beginning of the MCH-FP project in 1978; by 1993 this proportion had fallen to 37.4 percent. In the Comparison area, on the other hand, the proportion under 15 showed only minimal change, from 43.3 percent in 1978 to 42.0 percent in 1993.


It may be noted that the mid-year (30th June) population estimate for the study area of 207,703 is 468 less than the population enumerated in the 1993 census, for which the reference date was 1st June and which gave a total of 208,171. The explanation lies in the heavy net out-migration during the month of June, when a net loss of 701 migrants was recorded. This feature is discussed further in Chapter 6.

Table 2.1: Vital Statistics of the Matlab MCH-FP and Comparison Areas, 1982-1993

| Vital rates (per 1000) | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 |
|--|------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|------------------------|------------------------|-------------------------|-------------------------|------------------------|
| Crude birth rate MCH-FP area Comparison area Both areas | 36.9 44.7 40.7 | 34.2 42.6 38.3 | 30.7 37.3 34.0 | 34.6 42.6 38.5 | 33.6 39.6 36.5 | 33.6 39.2 36.4 | 30.9 40.4 35.5 | 28.4 36.6 32.4 | 28.3 37.8 32.9 | 25.4 32.7 29.0 | 25.4 31.1 28.2 | 24.7 29.4 27.0 |
| Total fertility rate** MCH-FP area Comparison area Both areas | 5.0 6.3 5.6 | 4.5 6.1 5.3 | 4.0 5.1 4.5 | 4.5 6.0 5.2 | 4.3 5.5 4.9 | 4.2 5.4 4.8 | 3.8 5.4 4.5 | 3.4 4.9 4.1 | 3.4 5.0 4.1 | 3.0 4.3 3.6 | 3.0 4.0 3.5 | 2.9 3.8 3.3 |
| Crude death rate MCH-FP area Comparison area Both areas | 12.5 15.9 14.2 | 11.9 16.7 14.3 | 13.4 17.3 15.3 | 10.2 14.2 12.2 | 9.9 12.2 11.0 | 9.3 11.2 10.2 | 8.7 11.0 9.9 | 8.0 9.5 8.7 | 7.6 9.4 8.5 | 8.1 10.2 9.1 | 8.3 9.8 9.0 | 7.7 10.2 8.9 |
| Neonatal mortality* MCH-FP area Comparison area Both areas | 58.1 68.1 63.5 | 56.4 70.3 64.0 | 57.9 71.4 65.3 | 52.5 69.4 61.7 | 45.4 53.0 49.4 | 43.8 54.9 49.7 | 42.8 57.7 51.1 | 46.0 52.7 49.7 | 47.8 53.3 50.9 | 47.7 63.2 56.3 | 49.6 53.3 51.6 | 42.8 64.5 54.4 |
| Post-neonatal mortalin MCH-FP area Comparison area Both areas | 47.5 50.2 49.0 | 41.8 42.2 42.0 | 56.9 55.7 56.2 | 33.8 49.1 42.1 | 36.4 39.7 38.2 | 34.6 39.5 37.2 | 38.0 39.0 38.6 | 28.3 38.0 33.6 | 27.4 34.1 31.2 | 32.3 51.7 43.0 | 30.8 37.0 34.1 | 20.3 34.8 28.0 |
| Infant mortality* MCH-FP area Comparison area Both areas | 105.6 118.3 112.5 | 98.2 112.5 106.0 | 114.8 127.1 121.5 | 86.4 118.4 103.8 | 81.8 92.7 87.6 | 78.4 94.4 86.9 | 80.8 96.6 89.6 | 74.3 90.7 83.3 | 75.2 87.5 82.1 | 80.0 114.9 99.2 | 80.5 90.2 85.7 | 63.1 99.3 82.4 |
| Child mortality (1-4 : MCH-FP area Comparison area Both areas | yrs) 18.8 27.4 23.3 | 21.9 35.3 29.1 | 23.1 39.2 31.6 | 16.4 24.6 20.7 | 13.4 20.7 17.2 | 9.9 15.0 12.6 | 7.6 14.4 11.1 | 6.4 11.5 9.0 | 5.3 9.3 7.4 | 7.0 9.1 8.1 | 5.9 10.4 8.3 | 5.9 10.0 8.1 |
| Under five mortality* MCH-FP area Comparison area Both areas | 169.4 207.2 189.7 | 172.3 227.0 202.1 | 192.0 252.7 224.8 | 143.9 200.1 174.4 | 129.8 164.0 148.0 | 113.1 145.2 130.2 | 107.4 146.1 128.3 | 97.5 131.1 115.7 | 94.8 120.4 108.7 | 105.7 146.2 128.1 | 102.0 127.1 115.7 | 87.8 139.7 115.6 |
| Rate of natural incre MCH-FP area Comparison area Both areas | 24.3 28.8 26.5 | 22.3 25.8 24.1 | 17.3 20.0 18.6 | 24.4 28.4 26.3 | 23.7 27.4 25.5 | 24.3 28.0 26.1 | 22.1 29.4 25.7 | 20.4 27.1 23.6 | 20.7 28.4 24.4 | 17.3 22.5 19.9 | 17.1 21.2 19.1 | 17.0 19.2 18.1 |
| In-migration Out-migration Growth (%) | 24.5 26.5 2.5 | 24.6 35.8 1.3 | 24.2 42.7 0.0 | 23.9 42.1 0.8 | 28:3 41.7 1.2 | 33.6 44.3 1.5 | 26.5 41.5 1.1 | 29.3 43.9 0.9 | 26.0 42.4 0.8 | 26.9 41.9 0.5 | 33.6 48.5 0.4 | 25.5 36.1 0.8 |

^{*}Per 1000 live births. **Per woman. Note: Numbers have been rounded.

Figure 2.1:Trends in Fertility and Under Five Mortality by Area, in 1982-1993






Table 2.2: Mid-year Population, Events Registered, and Population Changes in 1993

| | | Number | | | ate per : | 1000 |
|--|-----------------------------|--------------------------|--------------------------|-------------------------------|-----------------------|----------------------|
| | Total | Males | Females | Total | Males | Females |
| Total population as of 30 June 1993: | | | | | | |
| MCH-FP area Comparison area Both areas | 105770 101933 207703 | 52699 50837 103536 | 53071 51096 104167 | : | • | • |
| Events registered (Jan - Dec 1993) | | | | | | |
| Births MCH-FP area Comparison area Both areas | 2615 2992 5607 | 1345 1485 2830 | 1270 1507 2777 | 24.7 29.4 27.0 | • | |
| Deaths -Infant* MCH-FP area Comparison area Both areas | 165 297 462 | 91 158 249 | 74 139 213 | 63.1 99.3 82.4 | 67.7 106.4 88.0 | 58.3 92.2 76.7 |
| -All deaths MCH-FP area Comparison area Both areas | 812 1037 1849 | 432 559 991 | 380 478 858 | 7.7 10.2 8.9 | 8.2 11.0 9.6 | 7.2 9.4 8.2 |
| In-migration Out-migration Marriage Divorce** | 5292 7492 2697 334 | 2416 3707 | 2876 3785 - | 25.5 36.1 13.0 123.8 | 23.3 | 27.6 36.3 |
| Population change (Jan - Dec 1993) | | | | | | |
| Net Migration | -2200 | -1291 | -909 | -10.6 | -12.5 | -8.7 |
| Natural Increase MCH-FP area Comparison area Both areas | 1803 1955 3758 | 913 926 1839 | 890 1029 1919 | 17.0 19.2 18.1 | 17.3 18.2 17.8 | 16.8 20.1 18.4 |
| Net Increase | 1558 | 548 | 1010 | 7.5 | 5.3 | 9.7 |

^{*}Rate per 1000 live births. **Rate per 1000 marriages.

Table 2.3: Mid-year Population by Age and Sex, 1993

| | | Number | | | Percent | |
|------------------|------------------------------|------------------------------|------------------------------|--------------------------|--------------------------|--------------------------|
| Age (years) | Both sexes | Males | Females | Both sexes | Males | Females |
| All ages | 207703 | 103536 | 104167 | 100.0 | 100.0 | 100.0 |
| Under 1 | 5484 | 2790 | 2694 | 2.6 | 2.7 | 2.6 |
| 1 - 4 | 22746 | 11427 | 11319 | 11.0 | 11.0 | 10.9 |
| 1 2 3 4 | 5035 5835 5723 6153 | 2550 2962 2889 3026 | 2485 2873 2834 3127 | 2.4 2.8 2.8 3.0 | 2.5 2.9 2.8 2.9 | 2.4 2.8 2.7 3.0 |
| 5 - 9 | 28612 | 14586 | 14026 | 13.8 | 14.1 | 13.5 |
| 10-14 | 25535 | 13525 | 12010 | 12.3 | 13.1 | 11.5 |
| 15-19 | 21905 | 11576 | 10329 | 10.5 | 11.2 | 9.9 |
| 20-24 | 18700 | 9068 | 9632 | 9.0 | 8.8 | 9.2 |
| 25-29 | 15693 | 6855 | 8838 | 7.6 | 6.6 | 8.5 |
| 30-34 | 14682 | 7247 | 7435 | 7.1 | 7.0 | 7.1 |
| 35-39 | 10841 | 5400 | 5441 | 5.2 | 5.2 | 5.2 |
| 40-44 | 8201 | 4003 | 4198 | 3.9 | 3.9 | 4.0 |
| 45-49 | 7765 | 3435 | 4330 | 3.7 | 3.3 | 4.2 |
| 50-54 | 7428 | 3319 | 4109 | 3.6 | 3.2 | 3.9 |
| 55-59 | 6650 | 3278 | 3372 | 3.2 | 3.2 | 3.2 |
| 60-64 | 5220 | 2651 | 2569 | 2.5 | 2.6 | 2.5 |
| 65-69 | 3503 | 1819 | 1684 | 1.7 | 1.8 | 1.6 |
| 70 - 74 | 2358 | 1242 | 1116 | 1.1 | 1.2 | 1.1 |
| 75-79 | 1336 | 722 | 614 | 0.6 | 0.7 | 0.6 |
| 80-84 | 673 | 378 | 295 | 0.3 | 0.4 | 0.3 |
| 85+ | 371 | 215 | 156 | 0.2 | 0.2 | 0.1 |

Table 2.4: Mid-year Population by Area, Age, and Sex, 1993

| A | MCH | -FP area | | Comp | arison ar | ea |
|------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|
| Age (years) | Both sexes | Males | Females | Both sexes | Males | Females |
| All ages | 105770 | 52699 | 53071 | 101933 | 50837 | 51096 |
| Under 1 | 2592 | 1323 | 1269 | 2892 | 1467 | 1425 |
| 1 - 4 | 10442 | 5303 | 5139 | 12304 | 6124 | 6180 |
| 1 2 3 4 | 2300 2664 2646 2832 | 1164 1374 1343 1422 | 1136 1290 1303 1410 | 2735 3171 3077 3321 | 1386 1588 1546 1604 | 1349 1583 1531 1717 |
| 5 - 9 | 13916 | 7023 | 6893 | 14696 | 7563 | 7133 |
| 10-14 | 12620 | 6676 | 5944 | 12915 | 6849 | 6066 |
| 15-19 | 11669 | 6195 | 5474 | 10236 | 5381 | 4855 |
| 20-24 | 9918 | 4800 | 5118 | 8782 | 4268 | 4514 |
| 25-29 | 8316 | 3575 | 4741 | 7377 | 3280 | 4097 |
| 30-34 | 7742 | 3816 | 3926 | 6940 | 3431 | 3509 |
| 35-39 | 5719 | 2872 | 2847 | 5122 | 2528 | 2594 |
| 40-44 | 4375 | 2133 | 2242 | 3826 | 1870 | 1956 |
| 45-49 | 4103 | 1818 | 2285 | 3662 | 1617 | 2045 |
| 50-54 | 3915 | 1753 | 2162 | 3513 | 1566 | 1947 |
| 55-59 | 3433 | 1696 | 1737 | 3217 | 1582 | 1635 |
| 60-64 | 2667 | 1385 | 1282 | 2553 | 1266 | 1287 |
| 65-69 | 1867 | 968 | 899 | 1636 | 851 | 785 |
| 70 - 74 | 1195 | 644 | 551 | 1163 | 598 | 565 |
| 75-79 | 714 | 394 | 320 | 622 | 328 | 294 |
| 80-84 | 347 | 193 | 154 | 326 | 185 | 141 |
| 85+ | 220 | 132 | 88 | 151 | 83 | 68 |

Table 2.5: Mid-year Population in MCH-FP Area by Age, Sex, and Block, 1993


| | B16 | ock A | | B1 | ock B | Females 13426 336 1294 272 346 334 342 1804 1552 1337 1288 1181 945 | | | |
|------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--|--|--|--|
| Age (years) | Both sexes | Males | Females | Both sexes | Males | Females | | | |
| All ages | 29382 | 14588 | 14794 | 26505 | 13079 | 13426 | | | |
| Under 1 | 729 | 376 | 353 | 682 | 346 | 336 | | | |
| 1 - 4 | 3036 | 1512 | 1524 | 2681 | 1387 | 1294 | | | |
| 1 2 3 4 | 676 752 776 832 | 335 394 367 416 | 341 358 409 416 | 574 685 705 717 | 302 339 371 375 | 346 334 | | | |
| 5 - 9 | 3896 | 1971 | 1925 | 3669 | 1865 | 1804 | | | |
| 10-14 | 3558 | 1908 | 1650 | 3261 | 1709 | 1552 | | | |
| 15-19 | 3206 | 1645 | 1561 | 2848 | 1511 | 1337 | | | |
| 20-24 | 2673 | 1266 | 1407 | 2448 | 1160 | 1288 | | | |
| 25-29 | 2453 | 1036 | 1417 | 1993 | 812 | 1181 | | | |
| 30-34 | 2149 | 1030 | 1119 | 1836 | 891 | 945 | | | |
| 35-39 | 1629 | 823 | 806 | 1338 | 640 | 698 | | | |
| 40-44 | 1199 | 587 | 612 | 1101 | 516 | 585 | | | |
| 45-49 | 1177 | 531 | 646 | 985 | 399 | 586 | | | |
| 50-54 | 1003 | 486 | 517 | 948 | 402 | 546 | | | |
| 55-59 | 863 | 471 | 392 | 879 | 412 | 467 | | | |
| 60-64 | 660 | 341 | 319 | 703 | 372 | 331 | | | |
| 65-69 | 490 | 261 | 229 | 499 | 263 | 236 | | | |
| 70 - 74 | 288 | 148 | 140 | 291 | 178 | 113 | | | |
| 75-79 | 207 | 107 | 100 | 189 | 119 | 70 | | | |
| 80-84 | 94 | 49 | 45 | 92 | 54 | 38 | | | |
| 85+ | 72 | 40 | 32 | 62 | 43 | 19 | | | |

(continued)

Table 2.5 (cont.): Mid-year Population in MCH-FP Area by Age, Sex, and Block, 1993

| Ago | | Block C | | I | Block D | |
|------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| Age (years) | Both sexes | Males | Females | Both sexes | Males | Females |
| All ages | 27575 | 13930 | 13645 | 22308 | 11102 | 11206 |
| Under 1 | 632 | 319 | 313 | 549 | 282 | 267 |
| 1 - 4 | 2573 | 1323 | 1250 | 2152 | 1081 | 1071 |
| 1 2 3 4 | 586 661 652 674 | 316 350 334 323 | 270 311 318 351 | 464 566 513 609 | 211 291 271 308 | 253 275 242 301 |
| 5 - 9 | 3556 | 1824 | 1732 | 2795 | 1363 | 1432 |
| 10-14 | 3272 | 1707 | 1565 | 2529 | 1352 | 1177 |
| 15-19 | 3158 | 1714 | 1444 | 2457 | 1325 | 1132 |
| 20-24 | 2764 | 1381 | 1383 | 2033 | 993 | 1040 |
| 25-29 | 2099 | 947 | 1152 | 1771 | 780 | 991 |
| 30-34 | 2017 | 1037 | 980 | 1740 | 858 | 882 |
| 35-39 | 1549 | 766 | 783 | 1203 | 643 | 560 |
| 40-44 | 1160 | 573 | 587 | 915 | 457 | 458 |
| 45-49 | 1063 | 491 | 572 | 878 | 397 | 481 |
| 50-54 | 1075 | 486 | 589 | 889 | 379 | 510 |
| 55-59 | 902 | 449 | 453 | 789 | 364 | 425 |
| 60-64 | 686 | 373 | 313 | 618 | 299 | 319 |
| 65-69 | 439 | 215 | 224 | 439 | 229 | 210 |
| 70-74 | 335 | 163 | 172 | 281 | 155 | 126 |
| 75-79 | 161 | 88 | 73 | 157 | 80 | 77 |
| 80-84 | 84 | 45 | 39 | 77 | 45 | 32 |
| 85+ | 50 | 29 | 21 | 36 | 20 | 16 |

Figure 2.2: Age Pyramid of the 1993 Mid-year Population



CHAPTER 3

MORTALITY

Tables 3.1 to 3.3 show the distribution of deaths by sex and age for the whole study area, for the MCH-FP and Comparison areas, and for the four blocks of the MCH-FP area. Tables 3.4 and 3.5 show the corresponding age-sex-specific mortality rates for the study area, and for the MCH-FP and Comparison areas.

Tables 3.6 to 3.10 show the abridged life tables derived from these rates. A conspicuous feature of the 1993 results was the sharp drop in infant mortality in the MCH-FP area, where it fell from 80.5 in 1992 to 63.1 in 1993. This was the result of declines in both neonatal and post-neonatal mortality. The analysis by cause of death in unilluminating: of the neonatal deaths the reduction was entirely in the category "Other Neonatal", which comprises all congenital anomalies, sudden infant death, obstetric complications of newborn, immaturity, small-for-date, birth trauma, cord hemorrhage, birth asphyxia, hypoxia, unspecified neonatal death and prematurity; among the post-neonatal deaths the largest change was in those attributed to "Nutritional" causes.

The abridged life tables in Table 3.9 and 3.10 show that the expectation of life at birth continued to increase for both sexes in the MCH-FP area and for females in the Comparison area, but for males there was sharp drop from 60.5 in 1992 to 57.8 in 1993. This reduction in life expectancy occurred not only at birth but at all ages, reflecting a generalised increase in mortality. The age-standardised mortality rates by cause of death in Table 3.16, when compared with the corresponding table for 1992, shows that the biggest increases for the Comparison area males were in the rates for cardio-vascular causes and "senility", which tells us little. The latter term can in fact be regarded as a residual category of ill-defined causes of death among old people.

The levels of adult mortality showed little change in 1993, and there was again little difference between the two areas. The probability of dying between the ages of 15 and $60~(_{45}q_{15})$ increased from 198 per thousand in 1992 to 204 in 1993, for the study area as a whole, but there were compensating movements between the two areas and between sexes; the rate rose for males and fell for females in both areas. However, male adult mortality remained consistently higher than female in both areas, and the differential appears to be increasing.

Table 3.11 shows the distribution of deaths by age and month of occurrence. Adult deaths tend to peak in the winter months.

Neonatal deaths were also most frequent in these months, doubtless reflecting the seasonal variation in births, described in Chapter 4. Post-neonatal deaths on the other hand generally tend to be highest in April, May and June.

Tables 3.12 to 3.15 show the distribution of deaths by age, sex, area and cause, and Table 3.16 gives the age-standardized mortality rates by cause of death, using the W.H.O. "World Standard" age distribution shown in Appendix-D (WHO 1992). Comparisons with the corresponding table for 1992 reveal little changes, other those noted above.

It may be noted that the W.H.O. standard age distributions are not the same for males and for females. The female age distribution comprises a higher proportion over 50 that the male. This difference could affect comparisons of the age-standard mortality rates between the sexes. Furthermore, when the total age-standardised rates are compared with those for earlier years, the female rates for 1993 in both areas are higher than the corresponding rates for 1991, despite the fact that the expectations of life were also higher in 1993 than in 1991. This apparent anomaly may be explained by compensating movements in mortality rates in infancy and old age, which have different weights in the computation of the two mortality indices. Thus while infant mortality fell between 1991 and 1993, mortality rates in the oldest age groups increased in both areas.

Table 3.1: Deaths by Age and Sex, 1993

| Age | Both sexes | Males | Females |
|--|-----------------------------|----------------------------|-----------------------------|
| All ages | 1849 | 991 | 858 |
| Under 1 year Under 1 month 1-5 months 6-11 months | 462 305 104 53 | 249 169 55 25 | 213 136 49 28 |
| 1 - 4 years 1 2 3 4 | 185 65 54 39 27 | 73 26 18 13 16 | 112 39 36 26 11 |
| 5 - 9 | 44 | 27 | 17 |
| 10-14 | 22 | 11 | 11 |
| 15-19 | 20 | 11 | 9 |
| 20-24 | 34 | 15 | 19 |
| 25-29 | 37 | 18 | 19 |
| 30-34 | 25 | 14 | 11 |
| 35-39 | 28 | 16 | 12 |
| 40-44 | 35 | 18 | 17 |
| 45-49 | 39 | 23 | 16 |
| 50-54 | 81 | 51 | 30 |
| 55-59 | 106 | 67 | 39 |
| 60-64 | 128 | 76 | 52 |
| 65-69 | 148 | 82 | 66 |
| 70-74 | 153 | 79 | 74 |
| 75-79 | 143 | 81 | 62 |
| 80-84 | 95 | 44 | 51 |
| 85+ | 64 | 36 | 28 |

Table 3.2: Deaths by Area, Age, and Sex, 1993

| Age | MCH- | FP area | | Comparison area | | | |
|--|---------------------------|-------------------------|--------------------------|-----------------------------|---------------------------|---------------------------|--|
| Age | Both sexes | Males | Females | Both sexes | Males | Females | |
| All ages | 812 | 432 | 380 | 1037 | 559 | 478 | |
| Under 1 year Under 1 month 1-5 months 6-11 months | 165 112 31 22 | 91 66 15 10 | 74 46 16 12 | 297 193 73 31 | 158 103 40 15 | 139 90 33 16 | |
| 1 - 4 years 1 2 3 4 | 62 29 16 11 6 | 24 11 3 6 4 | 38 18 13 5 2 | 123 36 38 28 21 | 49 15 15 7 12 | 74 21 23 21 9 | |
| 5 - 9 | 16 | 10 | 6 | 28 | 17 | 11 | |
| 10-14 | 13 | 8 | 5 | 9 | 3 | 6 | |
| 15-19 | 7 | 5 | 2 | 13 | 6 | 7 | |
| 20-24 | 21 | 10 | 11 | 13 | 5 | 8 | |
| 25-29 | 21 | 11 | 10 | 16 | 7 | 9 | |
| 30-34 | 12 | 8 | 4 | 13 | 6 | 7 | |
| 35-39 | 14 | 8 | 6 | 14 | 8 | 6 | |
| 40-44 | 18 | 9 | 9 | 17 | 9 | 8 | |
| 45-49 | 19 | 11 | 8 | 20 | 12 | 8 | |
| 50-54 | 44 | 22 | 22 | 37 | 29 | 8 | |
| 55-59 | 51 | 36 | 15 | 55 | 31 | 24 | |
| 60-64 | 61 | 30 | 31 | 67 | 46 | 21 | |
| 65-69 | 70 | 34 | 36 | 78 | 48 | 30 | |
| 70-74 | 73 | 39 | 34 | 80 | 40 | 40 | |
| 75-79 | 74 | 41 | 33 | 69 | 40 | 29 | |
| 80-84 | 38 | 19 | 19 | 57 | 25 | 32 | |
| 85+ | 33 | 16 | 17 | 31 | 20 | 11 | |

Table 3.3: Deaths in MCH-FP Area by Age, Sex, and Block, 1993

| | В | lock A | | Block B | | | |
|--|-------------------------|-----------------------|------------------------|------------------------|-----------------------|-----------------------|--|
| Age | Both sexes | Males | Females | Both sexes | Males | Females | |
| All ages | 214 | 110 | 104 | 208 | 106 | 102 | |
| Under 1 year Under 1 month 1-5 months 6-11 months | 54 35 11 8 | 35 24 6 5 | 19 11 5 3 | 49 35 8 6 | 23 18 3 2 | 26 17 5 4 | |
| 1 - 4 years 1 2 3 4 | 23 10 6 5 2 | 9 5 0 2 2 | 14 5 6 3 0 | 11 5 2 3 1 | 4 1 0 2 1 | 7 4 2 1 0 | |
| 5 - 9 | 4 | 2 | 2 | 6 | 4 | 2 | |
| 10-14 | 3 | 1 | 2 | 4 | 2 | 2 | |
| 15-19 | 1 | 1 | 0 | 2 | 1 | 1 | |
| 20-24 | 4 | 1 | 3 | 6 | 3 | 3 | |
| 25-29 | 6 | 2 | 4 | 7 | 4 | 3 | |
| 30-34 | 3 | 1 | 2 | 1 | 1 | 0 | |
| 35-39 | 5 | 3 | 2 | 4 | 2 | 2 | |
| 40-44 | 5 | 2 | 3 | 5 | 3 | ·///\2 | |
| 45-49 | 5 | 3 | 2 | 8 | 6 | 2 | |
| 50-54 | 6 | 4 | 2 | 10 | 4 | 6 | |
| 55-59 | 15 | 10 | 5 | 10 | 6 | 4 | |
| 60-64 | 12 | 8 | 4 | 12 | 8 | 4 | |
| 65-69 | 18 | 8 | 10 | 17 | 7 | 10 | |
| 70-74 | 17 | 8 | 9 | 19 | 10 | 9 | |
| 75-79 | 10 | 2 | 8 | 20 | 13 | 7 | |
| 80-84 | 8 | 3 | 5 | 10 | 3 | 7 | |
| 85+ | 15 | 7 | 8 | 7 | 2 | 5 | |

(continued)

Table 3.3 (cont.): Deaths in MCH-FP Area by Age, Sex, and Block, 1993

| Age | В | Block C | | Block D | | | |
|--|------------------------|-----------------------|-----------------------|------------------------|-----------------------|-----------------------|--|
| Age | Both sexes | Males | Females | Both sexes | Males | Females | |
| All ages | 213 | 121 | 92 | 177 | 95 | 82 | |
| Under 1 year Under 1 month 1-5 months 6-11 months | 34 26 4 4 | 19 15 3 1 | 15 11 1 3 | 28 16 8 4 | 14 9 3 2 | 14 7 5 2 | |
| 1 - 4 years 1 2 3 4 | 18 8 6 3 1 | 9 4 2 2 1 | 9 4 4 1 0 | 10 6 2 0 2 | 2 1 1 0 0 | 8 5 1 0 2 | |
| 5 - 9 | 3 | 3 | 0 | 3 | . 1 | 2 | |
| 10-14 | 5 | 5 | 0 | 1 | 0 | 1 | |
| 15-19 | 1 | 1 | 0 | 3 | 2 | 1 | |
| 20-24 | 7 | 3 | 4 | 4 | 3 | 1 | |
| 25-29 | 6 | 3 | 3 | 2 | 2 | 0 | |
| 30-34 | 4 | 4 | 0 | 4 | 2 | 2 | |
| 35-39 | 3 | 1 | 2 | 2 | 2 | 0 | |
| 40-44 | 1 | 1 | 0 | 7 | 3 | 4 | |
| 45-49 | 3 | 1 | 2 | 3 | 1 | 2 | |
| 50-54 | 16 | 8 | 8 | 12 | 6 | 6 | |
| 55-59 | 19 | 15 | 4 | 7 | 5 | 2 | |
| 60-64 | 23 | 11 | 12 | 14 | 3 | 11 | |
| 65-69 | 15 | 7 | 8 | 20 | 12 | 8 | |
| 70-74 | 23 | 13 | 10 | 14 | 8 | 6 | |
| 75-79 | 17 | 10 | 7 | 27 | 16 | 11 | |
| 80-84 | 10 | 4 | 6 | 10 | 9 | 1 | |
| 85+ | 5 | 3 | 2 | 6 | 4 | 2 | |

Table 3.4: Death Rates by Age and Sex, 1993

| Age | Both sexes | Males | Females |
|---|----------------------------------|----------------------------------|-----------------------------------|
| All ages | 8.9 | 9.6 | 8.2 |
| Under 1 year* Under 1 month* 1-5 months* 6-11 months* | 82.4 54.4 18.5 9.5 | 88.0 59.7 19.4 8.8 | 76.7 49.0 17.6 10.1 |
| 1 - 4 years 1 2 3 4 | 8.1 12.9 9.3 6.8 4.4 | 6.4 10.2 6.1 4.5 5.3 | 9.9 15.7 12.5 9.2 3.5 |
| 5 - 9 | 1.5 | 1.9 | 1.2 |
| 10-14 | 0.9 | 0.8 | 0.9 |
| 15-19 | 0.9 | 1.0 | 0.9 |
| 20-24 | 1.8 | 1.7 | 2.0 |
| 25-29 | 2.4 | 2.6 | 2.1 |
| 30-34 | 1.7 | 1.9 | 1.5 |
| 35-39 | 2.6 | 3.0 | 2.2 |
| 40 - 44 | 4.3 | 4.5 | 4.0 |
| 45-49 | 5.0 | 6.7 | 3.7 |
| 50-54 | 10.9 | 15.4 | 7.3 |
| 55-59 | 15.9 | 20.4 | 11.6 |
| 60-64 | 24.5 | 28.7 | 20.2 |
| 65-69 | 42.2 | 45.1 | 39.2 |
| 70-74 | 64.9 | 63.6 | 66.3 |
| 75-79 | 107.0 | 112.2 | 101.0 |
| 80-84 | 141.2 | 116.4 | 172.9 |
| 85+ | 172.5 | 167.4 | 179.5 |

^{*}Rate per 1000 live births.

Table 3.5: Death Rates by Area, Age. and Sex, 1993 (per 1000 population)

| Age | MCH | -FP area | 1 | Comparison area | | | |
|--|----------------------------------|---------------------------------|-----------------------------------|------------------------------------|----------------------------------|-------------------------------------|--|
| Age | Both sexes | Males | Females | Both sexes | Males | Females | |
| All ages | 7.7 | 8.2 | 7.2 | 10.2 | 11.0 | 9.4 | |
| Under 1 year* Under 1 month* 1-5 months* 6-11 months* | 63.1 42.8 11.9 8.4 | 67.7 49.1 11.2 7.4 | 58.3 36.2 12.6 9.4 | 99.3 64.5 24.4 10.4 | 106.4 69.4 26.9 10.1 | 92.2 59.7 21.9 10.6 | |
| 1 - 4 years 1 2 3 4 | 5.9 12.6 6.0 4.2 2.1 | 4.5 9.5 2.2 4.5 2.8 | 7.4 15.8 10.1 3.8 1.4 | 10.0 13.2 12.0 9.1 6.3 | 8.0 10.8 9.4 4.5 7.5 | 12.0 15.6 14.5 13.7 5.2 | |
| 5 - 9 | 1.1 | 1.4 | 0.9 | 1.9 | 2.2 | 1.5 | |
| 10-14 | 1.0 | 1.2 | 0.8 | 0.7 | 0.4 | 1.0 | |
| 15-19 | 0.6 | 0.8 | 0.4 | 1.3 | 1.1 | 1.4 | |
| 20-24 | 2.1 | 2.1 | 2.1 | 1.5 | 1.2 | 1.8 | |
| 25-29 | 2.5 | 3.1 | 2.1 | 2.2 | 2.1 | 2.2 | |
| 30-34 | 1.5 | 2.1 | 1.0 | 1.9 | 1.7 | 2.0 | |
| 35-39 | 2.4 | 2.8 | 2.1 | 2.7 | 3.2 | 2.3 | |
| 40-44 | 4.1 | 4.2 | 4.0 | 4.4 | 4.8 | 4.1 | |
| 45-49 | 4.6 | 6.1 | 3.5 | 5.5 | 7.4 | 3.9 | |
| 50-54 | 11.2 | 12.5 | 10.2 | 10.5 | 18.5 | 4.1 | |
| 55-59 | 14.9 | 21.2 | 8.6 | 17.1 | 19.6 | 14.7 | |
| 60 - 64 | 22.9 | 21.7 | 24.2 | 26.2 | 36.3 | 16.3 | |
| 65-69 | 37.5 | 35.1 | 40.0 | 47.7 | 56.4 | 38.2 | |
| 70 - 74 | 61.1 | 60.6 | 61.7 | 68.8 | 66.9 | 70.8 | |
| 75-79 | 103.6 | 104.1 | 103.1 | 110.9 | 122.0 | 98.6 | |
| 80-84 | 109.5 | 98.4 | 123.4 | 174.8 | 135.1 | 227.0 | |
| 85+ | 150.0 | 121.2 | 193.2 | 205.3 | 241.0 | 161.8 | |

^{*}Rate per 1000 live births.

Figure 3.1: Probability of Survival from Birth to Age (x) by Sex, 1993




Table 3.6: Abridged Life Table, 1993

| Age (years) | $_{n}Q_{n}$ | 1 _x | L _x | e ⁰ |
|----------------|-------------|----------------|----------------|----------------|
| 0 | 82.4 | 100000 | 94034 | 61.6 |
| 1 | 12.8 | 91760 | 91066 | 66.1 |
| 2 | 9.2 | 90583 | 90166 | 65.9 |
| 3 | 6.8 | 89749 | 89444 | 65.5 |
| 4 | 4.4 | 89139 | 88944 | 65.0 |
| 5 | 7.7 | 88749 | 442177 | 64.3 |
| 10 | 4.3 | 88069 | 439472 | 59.8 |
| 15 | 4.6 | 87690 | 437531 | 55.0 |
| 20 | 9.1 | 87291 | 434632 | 50.2 |
| 25 | 11.7 | 86501 | 430163 | 45.7 |
| 30 | 8.5 | 85486 | 425760 | 41.2 |
| 35 | 12.8 | 84761 | 421296 | 36.5 |
| 40 | 21.1 | 83673 | 414280 | 32.0 |
| 45 | 24.8 | 81905 | 404824 | 27.6 |
| 50 | 53.2 | 79872 | 389486 | 23.2 |
| 55 | 76.8 | 75625 | 364553 | 19.4 |
| 60 | 115.9 | 69814 | 330026 | 15.8 |
| 65 | 191.9 | 61721 | 280301 | 12.5 |
| 70 | 280.1 | 49879 | 215339 | 9.8 |
| 75 | 421.3 | 35906 | 141333 | 7.7 |
| 80 | 516.5 | 20779 | 76030 | 6.5 |
| 85+ | 1000.0 | 10046 | 57341 | 5.8 |

Table 3.7: Abridged Life Tables by Sex, 1993

| | rv. | Male | es | | Females | | | | |
|----------------|-------------|----------------|----------------|----------------|-----------------|----------------|----------------|----------------|--|
| Age (years) | $_{n}q_{x}$ | 1 _x | L _x | e ⁰ | "q _x | 1 _x | L _x | e ⁰ | |
| 0 | 88.0 | 100000 | 93630 | 60.5 | 76.7 | 100000 | 94447 | 62.6 | |
| 1 | 10.1 | 91201 | 90656 | 65.3 | 15.6 | 92330 | 91481 | 66.8 | |
| 2 | 6.1 | 90276 | 90003 | 64.9 | 12.5 | 90892 | 90326 | 66.9 | |
| 3 | 4.5 | 89729 | 89528 | 64.3 | 9.1 | 89760 | 89350 | 66.7 | |
| 4 | 5.3 | 89326 | 89091 | 63.6 | 3.5 | 88940 | 88784 | 66.3 | |
| 5 | 9.2 | 88855 | 442388 | 62.9 | 6.0 | 88628 | 441905 | 65.5 | |
| 10 | 4.1 | 88036 | 439358 | 58.5 | 4.6 | 88092 | 439534 | 60.9 | |
| 15 | 4.7 | 87679 | 437437 | 53.7 | 4.3 | 87690 | 437570 | 56.2 | |
| 20 | 8.2 | 87263 | 434659 | 49.0 | 9.8 | 87308 | 434565 | 51.4 | |
| 25 | 13.1 | 86544 | 430116 | 44.4 | 10.7 | 86451 | 430123 | 46.9 | |
| 30 | 9.6 | 85415 | 425180 | 39.9 | 7.4 | 85527 | 426179 | 42.4 | |
| 35 | 14.7 | 84594 | 420095 | 35.3 | 11.0 | 84896 | 422332 | 37.7 | |
| 40 | 22.3 | 83349 | 412457 | 30.8 | 20.1 | 83965 | 415931 | 33.1 | |
| 45 | 33.0 | 81494 | 401248 | 26.4 | 18.3 | 82280 | 407920 | 28.7 | |
| 50 | 74.2 | 78807 | 380394 | 22.2 | 35.9 | 80773 | 397146 | 24.2 | |
| 55 | 97.5 | 72962 | 348128 | 18.8 | 56.3 | 77873 | 379166 | 20.0 | |
| 60 | 134.3 | 65847 | 308356 | 15.5 | 96.6 | 73488 | 350794 | 16.0 | |
| 65 | 203.4 | 57007 | 257248 | 12.5 | 179.2 | 66387 | 303576 | 12.4 | |
| 70 | 275.4 | 45410 | 196605 | 10.0 | 285.4 | 54490 | 234504 | 9.6 | |
| 75 | 436.7 | 32905 | 128085 | 7.9 | 402.7 | 38940 | 155300 | 7.4 | |
| 80 | 449.0 | 18535 | 71498 | 7.1 | 592.0 | 23258 | 79638 | 5.7 | |
| 85+ | 1000.0 | 10213 | 59343 | 5.8 | 1000.0 | 9490 | 52875 | 5.6 | |

Table 3.8: Abridged Life Tables by Area, 1993

| Age | | MCH- | FP area | | | Comparison area | | | | |
|---------|-------------|--------|----------------|----------------|-----------------|-----------------|----------------|----------------|--|--|
| (years) | $_{n}q_{x}$ | 1, | L _x | e ⁰ | "q _x | 1 _x | L _x | e ⁰ | | |
| 0 | 63.1 | 100000 | 95432 | 64.2 | 99.3 | 100000 | 92813 | 59.3 | | |
| 1 | 12.5 | 93690 | 92998 | 67.5 | 13.1 | 90074 | 89379 | 64.8 | | |
| 2 | 6.0 | 92516 | 92239 | 67.4 | 11.9 | 88896 | 88366 | 64.7 | | |
| 3 | 4.1 | 91962 | 91771 | 66.8 | 9.1 | 87837 | 87439 | 64.5 | | |
| 4 | 2.1 | 91581 | 91484 | 66.0 | 6.3 | 87041 | 86766 | 64.0 | | |
| 5 | 5.7 | 91387 | 455727 | 65.2 | 9.5 | 86492 | 430569 | 63.4 | | |
| 10 | 5.1 | 90863 | 453239 | 60.5 | 3.5 | 85672 | 427672 | 59.0 | | |
| 15 | 3.0 | 90396 | 451356 | 55.8 | 6.3 | 85374 | 425623 | 54.2 | | |
| 20 | 10.5 | 90125 | 448436 | 51.0 | 7.4 | 84833 | 422723 | 49.5 | | |
| 25 | 12.6 | 89176 | 443296 | 46.5 | 10.8 | 84207 | 418941 | 44.9 | | |
| 30 | 7.7 | 88056 | 438714 | 42.1 | 9.3 | 83299 | 414703 | 40.4 | | |
| 35 | 12.2 | 87376 | 434428 | 37.4 | 13.6 | 82522 | 410023 | 35.7 | | |
| 40 | 20.4 | 86313 | 427500 | 32.8 | 22.0 | 81401 | 402869 | 31.2 | | |
| 45 | 22.9 | 84554 | 418292 | 28.4 | 27.0 | 79611 | 393089 | 26.8 | | |
| 50 | 54.8 | 82617 | 402564 | 24.0 | 51.4 | 77464 | 378068 | 22.5 | | |
| 55 | 71.8 | 78093 | 377383 | 20.3 | 82.2 | 73482 | 353292 | 18.5 | | |
| 60 | 108.5 | 72486 | 343947 | 16.6 | 123.6 | 67442 | 317570 | 15.0 | | |
| 65 | 172.1 | 64620 | 296626 | 13.3 | 213.9 | 59108 | 265179 | 11.7 | | |
| 70 | 266.0 | 53498 | 232923 | 10.6 | 294.4 | 46465 | 198873 | 9.2 | | |
| 75 | 411.0 | 39269 | 155710 | 8.4 | 433.0 | 32785 | 127966 | 7.0 | | |
| 80 | 428.7 | 23131 | 90562 | 7.6 | 596.3 | 18590 | 63394 | 5.4 | | |
| 85+ | 1000.0 | 13214 | 85500 | 6.5 | 1000.0 | 7505 | 36559 | 4.9 | | |

Table 3.9: Abridged Life Tables for MCH-FP Area by Sex, 1993

| Age (years) | | Ma | les | | | Females | | | | |
|----------------|-----------------------------|--------|----------------|----------------|-----------------------------|---------|----------------|----------------|--|--|
| | _n q _x | 1, | L _x | e ⁰ | _n q _x | 1, | L _x | e ⁰ | | |
| 0 | 67.7 | 100000 | 95102 | 63.5 | 58.3 | 100000 | 95781 | 64.8 | | |
| 1 | 9.4 | 93234 | 92717 | 67.1 | 15.7 | 94173 | 93300 | 67.8 | | |
| 2 | 2.2 | 92357 | 92256 | 66.8 | 10.0 | 92693 | 92228 | 67.9 | | |
| 3 | 4.5 | 92156 | 91950 | 65.9 | 3.8 | 91763 | 91587 | 67.5 | | |
| 4 | 2.8 | 91745 | 91616 | 65.2 | 1.4 | 91412 | 91347 | 66.8 | | |
| 5 | 7.1 | 91487 | 455940 | 64.4 | 4.3 | 91282 | 455497 | 65.9 | | |
| 10 | 6.0 | 90838 | 452939 | 59.8 | 4.2 | 90886 | 453549 | 61.2 | | |
| 15 | 4.0 | 90295 | 450638 | 55.2 | 1.8 | 90504 | 452140 | 56.4 | | |
| 20 | 10.4 | 89932 | 447507 | 50.4 | 10.7 | 90339 | 449466 | 51.5 | | |
| 25 | 15.3 | 88999 | 441858 | 45.9 | 10.5 | 89373 | 444701 | 47.0 | | |
| 30 | 10.4 | 87640 | 436090 | 41.5 | 5.1 | 88435 | 441139 | 42.5 | | |
| 35 | 13.8 | 86725 | 430857 | 37.0 | 10.5 | 87985 | 437799 | 37.7 | | |
| 40 | 20.9 | 85525 | 423497 | 32.4 | 19.9 | 87063 | 431314 | 33.1 | | |
| 45 | 29.8 | 83738 | 412909 | 28.1 | 17.4 | 85331 | 423235 | 28.7 | | |
| 50 | 61.0 | 81240 | 394668 | 23.8 | 49.7 | 83850 | 409565 | 24.2 | | |
| 55 | 101.1 | 76287 | 363339 | 20.2 | 42.3 | 79682 | 390584 | 20.3 | | |
| 60 | 103.1 | 68575 | 326284 | 17.2 | 114.4 | 76309 | 361009 | 16.1 | | |
| 65 | 162.1 | 61507 | 283854 | 13.9 | 182.8 | 67579 | 308434 | 12.8 | | |
| 70 | 264.0 | 51537 | 224650 | 11.0 | 268.3 | 55228 | 240127 | 10.1 | | |
| 75 | 412.2 | 37932 | 150273 | 9.1 | 409.4 | 40411 | 160418 | 7.8 | | |
| 80 | 394.8 | 22295 | 89407 | 8.7 | 468.9 | 23868 | 90702 | 6.5 | | |
| 85+ | 1000.0 | 13493 | 104770 | 7.8 | 1000.0 | 12678 | 65625 | 5.2 | | |

Table 3.10: Abridged Life Tables for Comparison Area by Sex, 1993

| Ago | | Ma | ales | | | F | emales | |
|----------------|-----------------------------|----------------|----------------|----------------|-------------|----------------|----------------|----------------|
| Age (years) | _n q _x | 1 _x | L _x | e ⁰ | $_{n}q_{x}$ | 1 _x | L _x | e ⁰ |
| 0 | 106.4 | 100000 | 92297 | 57.8 | 92.2 | 100000 | 93322 | 60.8 |
| 1 | 10.8 | 89360 | 88793 | 63.6 | 15.4 | 90776 | 89949 | 66.0 |
| 2 | 9.4 | 88398 | 87983 | 63.3 | 14.4 | 89374 | 88729 | 66.0 |
| 3 | 4.5 | 87567 | 87369 | 62.9 | 13.6 | 88085 | 87485 | 66.0 |
| 4 | 7.5 | 87172 | 86847 | 62.2 | 5.2 | 86885 | 86657 | 65.9 |
| 5 | 11.2 | 86522 | 430377 | 61.7 | 7.7 | 86430 | 430621 | 65.2 |
| 10 | 2.2 | 85554 | 427341 | 57.3 | 4.9 | 85766 | 427856 | 60.7 |
| 15 | 5.6 | 85367 | 425742 | 52.5 | 7.2 | 85343 | 425302 | 56.0 |
| 20 | 5.8 | 84893 | 423320 | 47.7 | 8.8 | 84730 | 421925 | 51.4 |
| 25 | 10.6 | 84397 | 419916 | 43.0 | 10.9 | 83982 | 417794 | 46.8 |
| 30 | 8.7 | 83500 | 415825 | 38.4 | 9.9 | 83064 | 413420 | 42.3 |
| 35 | 15.7 | 82773 | 410864 | 33.8 | 11.5 | 82240 | 409016 | 37.7 |
| 40 | 23.8 | 81473 | 402882 | 29.3 | 20.3 | 81294 | 402663 | 33.1 |
| 45 | 36.5 | 79534 | 390946 | 24.9 | 19.4 | 79647 | 394667 | 28.7 |
| 50 | 88.7 | 76633 | 367245 | 20.7 | 20.4 | 78103 | 386841 | 24.2 |
| 55 | 93.7 | 69832 | 333833 | 17.5 | 71.0 | 76513 | 369901 | 19.7 |
| 60 | 167.2 | 63290 | 291286 | 14.0 | 78.6 | 71084 | 342367 | 16.0 |
| 65 | 248.2 | 52706 | 231886 | 11.3 | 175.1 | 65497 | 300164 | 12.1 |
| 70 | 287.5 | 39627 | 170321 | 9.2 | 301.7 | 54026 | 230207 | 9.1 |
| 75 | 464.9 | 28234 | 107623 | 6.9 | 395.4 | 37728 | 151233 | 7.0 |
| 80 | 500.8 | 15110 | 55997 | 5.8 | 696.3 | 22810 | 69981 | 4.9 |
| 85+ | 1000.0 | 7542 | 31301 | 4.2 | 1000.0 | 6928 | 42829 | 6.2 |

Table 3.11: Deaths by Age and Month, 1993

| Month | | | Age | at death | |
|-----------|-------------|------------------|----------------|--------------|---------------------|
| HOHEH | All ages | Under 1 month | 1-11 months | 1-4 years | 5 years and over |
| January | 170 | 33 | 9 | 15 | 113 |
| February | 146 | 20 | 14 | 9 | 103 |
| March | 149 | 17 | 14 | 18 | 100 |
| April | 135 | 14 | 21 | 10 | 90 |
| May | 148 | 16 | 16 | 15 | 101 |
| June | 145 | 19 | 16 | 17 | 93 |
| July | 113 | 19 | 11 | 14 | 69 |
| August | 132 | 20 | 11 | 17 | 84 |
| September | 153 | 27 | 8 | 23 | 95 |
| October | 173 | 37 | 10 | 19 | 107 |
| November | 196 | 43 | 15 | 14 | 124 |
| December | 189 | 39 | 12 | 14 | 124 |
| Total | 1849 | 304 | 157 | 185 | 1203 |

Table 3.12: Male Deaths by Cause and Age, 1993

| | | | | | | | | | Age at | death | (years) | 0 | | | | | | | |
|---|------------------|-------|------|-------|---------|----------|-----------|------|----------|-----------|-----------|-----------|---------------------|---------|-------|----------|---------------|---------------|------|
| Cause | ages | 7 | 1-4 | 5-9 1 | 10-14 1 | 15-19 20 | 20-24 25- | 53 | 30-34 35 | 35-39 40- | 40-44 45- | -49 50-54 | 54 55-59 | 9 60-64 | 69-59 | 70-74 | 75-79 | 80-84 | 85+ |
| DIARRHOEAL Diarrhoea Dysentery | 91 | 29 | 13 | 23 | 00 | 10 | .00 | 20 | 0 2 | 12 | 20 | | 2 8 0 3 | 30 | 22 | 1 3 | 94 | 0 3 | 1 6 |
| INFECTIOUS Tuberculosis Tetanus (non-neonatal) Other infectious | 31 22 21 | 100 | 700 | 204 | 301 | 010 | 100 | 000 | 000 | 001 | 100 | 201 | 2 2 2 | 000 | 1 0 0 | 1001 | 000 | 011 | 000 |
| MALIGNANT NEOPLASMS NUTRITION CARDIO-VASCULAR | 33 20 71 | 0 7 0 | 100 | 000 | 100 | поп | 105 | 510 | 0003 | 503 | 1112 | 004 | 1000 | 12 | 110 | 406 | 8 0 1 | 200 | 001 |
| RESPIRATORY ARI, pneum, influenza COPD* | 58 | 42 | 7 | 40 | 0 -1 | 00 | 00 | 00 | 10 | 00 | 10 | 20 | 2 7 | H 80 | 14 | 3 | 11 | 20 | 10 |
| GASTRO-INTESTINAL DIRECT OBSTETRIC | 56 | 0 2 | 10 | 00 | 10 | 10 | 0 5 | 0 | 10 | 00 | 03 | 40 | 7 11 0 | 80 | 7 | 0 3 | 03 | 0 11 | 00 |
| NEONATAL Tetanus Other neonatal | 142 | 142 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 |
| ACCIDENTS, INJURIES Suicide Homicide Drowning Other accidents, etc. | 13 33 21 | 0041 | 2400 | 1200 | 1000 | 2002 | 2003 | 4108 | 0000 | 1000 | 0001 | 0000 | 0001 | 0001 | 0000 | 1000 | 1001 | 0000 | 0000 |
| OTHER AND UNSPECIFIED Senility Other causes n.e.c.** Unknown | 116 103 86 | 046 | 2150 | 042 | 150 | 011 | 220 | 310 | 220 | 10 | 024 | 350 | 0 0 8 7 12 11 | 4111 | 100 | 27 8 8 3 | 26 15 2 | 21 10 0 | 19 |
| TOTAL | 991 | 249 | 73 | 27 | 11 | 11 | 15 | 18 | 14 | 16 | 18 2 | 23 51 | 1 67 | 76 | 82 | 79 | 81 | 44 | 36 |
| | | | | | | | | | | | | | | | | | | | |

*Chronic obstructive pulmonary disease. **Not elsewhere classified.

Table 3.13: Female Deaths by Cause and Age, 1993

| | | | | | | | | | Age | at | death (years) | ears) | | | | | | | | |
|---|--------------------|--------|--------------|------|---------|---------|---------|---------|---------|------|---------------|----------|----------|----------------|----------|---------|---------------|----------|---------|--------------|
| Cause | ages | 4 | 1.4 | 5-9 | 10-14] | 15-19 2 | 20-24 2 | 25-29 3 | 30-34 3 | -39 | 40-44 4 | 45-49 50 | 50-54 55 | 55-59 60 | 60-64 65 | 65-69 7 | 70-74 7 | 75-79 80 | 80-84 | 85+ |
| DIARRHOEAL Diarrhoea Dysentery | 124 | 22 | 4, | നന | mo | но | 00 | m0 | 10 | 00 | 00 | 10 | 1 2 | г г | 12 | ∞ ⊣ | ∞ N | ∞ ⊢ | 7 0 | 0.22 |
| INFECTIOUS Tuberculosis Tetanus (non-neonatal) Other infectious | 111 0 0 28 | 004 | 009 | 000 | 001 | 001 | 101 | 301 | 000 | 201 | 101 | нон | 000 | 301 | 100 | 000 | 001 | 001 | 000 | 100 |
| MALIGNANT NUTRITION CARDIO-VASCULAR | 24 16 60 | 000 | 0 0 9 1 | ноо | 200 | 010 | 000 | ноо | 100 | 000 | 105 | 301 | 101 | 909 | 806 | 3 | 2 0 11 | 100 | 005 | 001 |
| RESPIRATORY ARI, pneum, influenza COPD* | 59 45 | 40 | 14 | 10 | 00 | 00 | 00 | 00 | 0 1 | 00 | 10 | 10 | 21 | 31 | 08 | 10 | 0 | 3 🗆 | 04 | 10 |
| GASTRO-INTESTINAL DIRECT OBSTETRIC | 24 | ПО | 00 | 00 | 10 | | 20 | r 7 | 3.0 | 10 | 12 | 0 2 | 90 | 03 | 00 | 0 2 | 0 | 0.02 | 00 | 00 |
| NEONATAL Tetanus (neonatal) Other neonatal | 3 | 3 | 000 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 |
| ACCIDENTS, INJURIES Suicide Homicide Drowning Other accidents, etc. | 8 1 25 12 | 1500 | 0 19 2 | 0000 | 0000 | 0000 | 00012 | 300% | 1000 | 0000 | 0000 | 0000 | 0010 | 0000 | 0000 | 0001 | 0001 | 0000 | 0000 | 0000 |
| OTHER AND UNSPECIFIED Senility Other causes n.e.c.** Unknown | 107 117 51 | 0 7 10 | 000 | 130 | 070 | 130 | 024 | 200 | 0 6 1 | 210 | 079 | 040 | 130 | 201 | 10 3 | 2 2 | 22 20 1 | 13 | 20 0 | 17 3 0 |
| TOTAL | 858 | 213 | 112 | 17 | 11 | 6 | 19 | 19 | 11 | 12 | 17 | 16 | 30 | 39 | 52 | 99 | 74 | 62 | 51 | 28 |
| | | | | | | | | | | | | | | | | | | | | Ī |

*Chronic obstructive pulmonary disease. **Not elsewhere classified.

Table 3.14: Male Deaths by Cause, Age, and Area, 1993

| | | | | | | | | Age at | death | death (years) | _ | | | | | |
|---|---------------|----------------|------|------|----------|------|------|--------|--------|---------------|--------------|------|--------------|--------------|-------|------|
| | All | ages | \ \ | | 1 | 1-4 | 5 | 5-14 | 15-44 | 44 | 45-64 | 64 | 69 | 65-84 | 00 | 85+ |
| cause | Σ | U | Σ | U | Σ | U | Σ | U | Σ | U | Σ | O | Σ | U | Σ | O |
| DIARRHOEAL Diarrhoea Dysentery | 35 | 56 15 | 12 | 17 | 3.2 | 11 2 | 1 2 | | 12 | 4 0 | 80 4 | 6.6 | 23 | 11 5 | 80 | mH |
| INFECTIOUS Tuberculosis Tetanus (non-neonatal) Other infectious. | 14 14 | 17 | 001 | 000 | 000 | 102 | 000 | 103 | 200 | 0 | 204 | 10 | o ⊢ ⊢ | 1003 | 000 | 000 |
| MALIGNANT NUTRITION CARDIO-VASCULAR | 21 6 29 | 12 14 42 | 010 | 090 | 1 4 0 | 090 | 001 | 000 | 717 | 414 | ∞ 0 ∞ | 21 | 5 0 17 | 2 0 16 | 000 | 001 |
| RESPIRATORY ARI, pneum, influenza COPD* | 31 | 30 | 13 | 50 | 00 | 7 | 10 | г П | 01 | 0 H | 7.1 | 12 | 1 22 | 3 | 10 | 00 |
| GASTRO-INTESTINAL DIRECT OBSTETRIC | 30 | 26 | 0 5 | 00 | 10 | 00 | 10 | 00 | 0 2 | 0 3 | 13 | 17 | 8 0 | 90 | 00 | 00 |
| NEONATAL Tetanus Other neonatal | 53 | 89 | 53 | 89 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 00 |
| ACCIDENTS, INJURIES Suicide Homicide Drowning Other accidents, etc. | 141 | 5 19 12 | 0010 | 0061 | 0 0 11 0 | 13 | 1000 | 1300 | . 0000 | 2012 | 7007 | 2000 | 1001 | 0001 | 0000 | 0000 |
| OTHER AND UNSPECIFIED Senility Other causes n.e.c.** | 47 49 40 | 69 54 46 | 979 | 000 | 011 | 014 | 120 | 0 4 2 | 096 | 000 | 188 | 1333 | 39 | 54 7 | 0 2 7 | 12 3 |
| TOTAL | 432 | 559 | 91 | 158 | 24 | 49 | 18 | 20 | 51 | 41 | 66 | 118 | 133 | 153 | 16 | 20 |

*Chronic obstructive pulmonary disease. **Not elsewhere classified.

Table 3.15: Female Deaths by Cause, Age, and Area, 1993

| | | | | | | | | Age a | at deat | death (years) | (5) | | | | | |
|---|--|----------------|------|------|------|---------|------|-------|----------|---------------|---------------|----------------|---------------|---------------|------|------|
| Cause | LL | All ages | V | √. | 1 | 1.4 | 5 | 5-14 | 15. | 15-44 | 45- | 45-64 | 65 | 65-84 | | 85+ |
| 27.33.33.33.33.33.33.33.33.33.33.33.33.33 | Σ | S | Σ | U | Σ | U | Σ | U | Σ | J | Σ | U | Σ | U | Σ | O |
| DIARRHOEAL Diarrhoea Dysentery | 50 | 74 | 11 | 111 | 14 | 30 | 41 | 4.5 | 0.0 | 0.5 | 4 % | ю н | 10 | 21 2 | 0.02 | 00 |
| INFECTIOUS Tuberculosis Tetanus (non-neonatal) Other infectious | 706 | 4 0 61 | 000 | 200 | 000 | 000 | 000 | 001 | 202 | 209 | 103 | 308 | 308 | 100 | 001 | 000 |
| MALIGNANT NUTRITION CARDIO-VASCULAR | 11 5 40 | 13 20 | 040 | 000 | 010 | 0 21 | 000 | 000 | 808 | | 2 0 11 | 808 | 4 0 25 | 2 0 11 | 001 | 000 |
| RESPIRATORY ARI, pneum, influenza COPD* | 16 | 43 | 12 0 | 28 | 23 | 111 | 00 | 10 | 10 | 01 | 10 | 1 | 10 | 2 12 | 0 1 | 00 |
| GASTRO-INTESTINAL DIRECT OBSTETRIC | 14 | 10 | 00 | 10 | 00 | 00 | 10 | 00 | 3.2 | 4 6 | 60 | 0 5 | 0 0 | 0 3 | 00 | 00 |
| NEONATAL Tetanus (neonatal) Other neonatal | 38 | 3 74 | 38 | 3 74 | 00 | 00 | 00 | 00 | 00 | 00 | 0 0 | 00 | 00 | 00 | 00 | 00 |
| ACCIDENTS, INJURIES Suicide Homicide Drowning Other accidents, etc. | 1 16 2 | 3000 | 0000 | 0001 | 0020 | 2 7 0 0 | 0010 | 2200 | 2 0 1 2 | 1000 | 0010 | 0000 | 0000 | 0004 | 0000 | 0000 |
| OTHER AND UNSPECIFIED Senility Other causes n.e.c.** Unknown | 49 25 | 58 61 26 | 220 | 820 | 130 | 220 | 070 | 130 | 0 0 0 10 | 020 | 10 13 8 | 10 | 31 29 4 | 42 31 0 | 810 | 940 |
| TOTAL | 380 | 478 | 74 | 139 | 38 | 74 | 11 | 17 | 42 | 45 | 9/ | 61 | 122 | 131 | 17 | 11 |
| | THE STATE OF THE S | | | | | | | | | | | | | | | ĺ |

*Chronic obstructive pulmonary disease. **Not elsewhere classified.

Table 3.16: Age-standardized Mortality Rates by Cause of Death, 1993

| | | Males | Ferr | nales |
|---------------------------|----------------|--------------------|----------------|------------|
| Cause of death | MCH-FP area | Comparison area | MCH-FP area | Comparisor |
| Diarrhoea | 70.35 | 115.54 | 129.04 | 178.20 |
| Dysentery | 24.61 | 35.10 | 21.17 | 17.62 |
| Tuberculosis | 31.33 | 37.88 | 15.11 | 9.77 |
| Tetanus (non-neonatal) | | 1.86 | 0.00 | |
| Other infectious | 27.17 | 11.69 | 28.38 | 39.98 |
| Malignant neoplasms | 42.54 | 31.10 | 26.50 | 30.36 |
| Nutritional | 11.47 | 24.93 | 9.59 | 18.12 |
| Cardio-vascular | 64.18 | 104.11 | 133.99 | 59.41 |
| ARI, pneumonia, influenza | 30.99 | 73.42 | 30.63 | 76.74 |
| C.O.P.D.* | 68.99 | 71.98 | 72.41 | 64.27 |
| Gastro-intestinal | 64.85 | 62.56 | 31.79 | 24.58 |
| Direct obstetric | 1.60 | = | 5.34 | 19.35 |
| Neonatal tetanus | | | | 5.20 |
| Other neonatal | 102.47 | 155.19 | 73.99 | 128.32 |
| Suicide | | 11.48 | 8.91 | 6.05 |
| Homicide | | | | * |
| Drowning | 22.95 | 29.72 | 28.85 | 14.89 |
| Other accidents | 18.06 | 25.71 | 3.46 | 26.01 |
| Senility | 106.98 | 180.42 | 197.24 | 273.49 |
| Other cause n.e.c.** | 104.69 | 130.27 | 163.31 | 208.35 |
| Unknown | 86.06 | 103.71 | 57.63 | 50.09 |
| Total | 904.44 | 1212.75 | 1039.15 | 1250.80 |

^{*}Chronic obstructive pulmonary disease. **Not elsewhere classified.

CHAPTER 4

FERTILITY

Table 4.1 shows the number of pregnancies and their outcomes in 1993. Compared with 1992, the number of live births fell by 189 or 3.3 percent. In the MCH-FP area the number of live births was 45 less than in 1992, and in the Comparison area they were 144 less than in 1992. In the study area as a whole, 87.4 percent of pregnancies resulted in a live birth; a proportion which remains remarkably constant from year to year.

Table 4.2 shows the distribution of pregnancies by outcome, and live births by sex and month of occurrence. The data show the usual marked seasonal variation, peaking in October-December. The sex ratio of the live births was 101.9 males per 100 females.

Table 4.3 shows the age-specific fertility rates for the study area, together with the total fertility rate, crude birth rate, general fertility rate, and gross and net reproduction rates; Table 4.4 shows the corresponding rates for the MCH-FP and Comparison areas, which are also illustrated in Figure 4.2; Table 4.5 shows the rates for the four blocks of the MCH-FP area. Although the total fertility rate in the MCH-FP area remained virtually unchanged from 1992, the shape of the age-specific fertility distribution changed. More births were concentrated in the 20-24 age group, with correspondingly fewer at older ages. In the Comparison area the age-specific rates fell for all groups except the group 45-49.

Table 4.6 shows the distribution of births by mother's age and live birth order, and Table 4.7 the age-order-specific fertility rates derived from these figures. These tables were included from 1992 in the DSS reports, and their production has been made possible by the up-dating of the pregnancy history files. The breakdown by birth order facilitates a more detailed and sensitive analysis of fertility trends and differentials. Thus the totals of the order-specific rates represent the components by birth order of the total fertility rates. In the same way as the TFR represents the average number of children which would be borne by a woman who go through life having children at the current rates, so the total for birth order N represents the proportion of women who would have at least N children.

Thus the tables highlight the differences between the MCH-FP and Comparison areas. There is comparatively little difference between the two areas for birth orders 1 to 4, but thereafter they widen dramatically: for birth orders 6 and 7, the Comparison area rates are twice as high as those of the MCH-FP area, and for birth order 8 and more, they are more than three times as great. This volume also includes, for the first time, a special supplement describing fertility trends in Matlab by birth order, age at birth and birth interval.

Table 4.1: Number and Rates of Pregnancy Outcomes by Type and Area, 1993

| Type of pregnancy | Both | areas | MCH- | FP area | Compar | ison area |
|----------------------------|------|-------|------|---------|--------|-----------|
| outcome | No. | Rate | No. | Rate | No. | Rate |
| Total pregnancies* | 6350 | 126.5 | 2913 | 109.4 | 3437 | 145.8 |
| Live birth pregnancies** | 5552 | 874.3 | 2593 | 890.1 | 2959 | 860.9 |
| Fetal wastage | 798 | 125.7 | 320 | 109.9 | 478 | 139.1 |
| Early (miscarriage) | 577 | 90.9 | 232 | 79.6 | 345 | 100.4 |
| Late (stillbirths) | 221 | 34.8 | 88 | 30.2 | 133 | 38.7 |
| Multiple birth pregnancies | | 76 | - 1 | 30 | | 46 |
| Live birth pregnancies | 1 | 65 | | 25 | | 40 |
| Three live births | | 1 | | 1 | | 0 |
| Two live births | | 53 | | 20 | | 33 |
| One live birth | | 11 | | 4 | | 7 |
| Stillbirth pregnancies | | 1 | | 0 | | 1 |
| Miscarriage pregnancies | | 10 | | 5 | | 5 |

^{*}Rates per 1000 women of age 15-49 years. **Ratio per 1000 total pregnancies.

Table 4.2: Pregnancy Outcomes by Month, 1993

| | | Pr | egnancy | outcom | ie | No.o | flive | born chil | dren |
|------------|------|---------|---------|--------|-------|-------|-------|-----------|--------|
| Months | | Miscarr | iage | Still | Live* | Both | | | |
| | All | Induced | Spon. | birth | birth | sexes | Males | Females | Ratio |
| All months | 6350 | 217 | 360 | 221 | 5552 | 5607 | 2830 | 2777 | 1.0191 |
| January | 555 | 20 | 27 | 19 | 489 | 492 | 245 | 247 | 0.9919 |
| February | 451 | 16 | 24 | 13 | 398 | 399 | 202 | 197 | 1.0254 |
| March | 572 | 25 | 27 | 19 | 501 | 506 | 252 | 254 | 0.9921 |
| April | 420 | 24 | 38 | 17 | 341 | 346 | 179 | 167 | 1.0719 |
| May | 422 | 18 | 34 | 13 | 357 | 363 | 185 | 178 | 1.0393 |
| June | 391 | 22 | 33 | 12 | 324 | 326 | 163 | 163 | 1.0000 |
| July | 465 | 26 | 35 | 15 | 389 | 392 | 211 | 181 | 1.1657 |
| August | 507 | 17 | 41 | 11 | 438 | 441 | 236 | 205 | 1.1512 |
| September | 562 | 19 | 32 | 25 | 486 | 492 | 244 | 248 | 0.9839 |
| October | 681 | 15 | 23 | 29 | 614 | 622 | 321 | 301 | 1.0664 |
| November | 642 | 5 | 32 | 31 | 574 | 580 | 282 | 298 | 0.9463 |
| December | 682 | 10 | 14 | 17 | 641 | 648 | 310 | 338 | 0.9172 |

 $[\]mbox{*For any multiple pregnancy, the outcome is recorded as live birth if at least one of the issues is live born.}$

Figure 4.1: Number of Births and Deaths by Month, 1993




Table 4.3: Age-specific Fertility Rates and Indices, 1993

| Age (years) | Number of live births | Number of women | ASFR (per 1000) |
|----------------|--------------------------|-----------------|--------------------|
| All ages | 5607 | 50203 | 111.7 |
| 15-19* | 649 | 10329 | 62.8 |
| 20-24 | 1905 | 9632 | 197.8 |
| 25-29 | 1665 | 8838 | 188.4 |
| 30-34 | 938 | 7435 | 126.2 |
| 35-39 | 347 | 5441 | 63.8 |
| 40 - 44 | 89 | 4198 | 21.2 |
| 45-49** | 14 | 4330 | 3.2 |
| Tot | al Fertility Rate (TFF | ₹) = | 3317 |
| Gen | eral Fertility Rate ((| GFR) = | 112 |
| Gro | ss Reproduction Rate | (GRR) = | 1643 |
| Net | Reproduction Rate (N | RR) = | 1413 |

*Births to mothers under age 15 were included in this group.

this group.
**Births to mothers age 50 and above were included in this group.

Table 4.4: Age-specific Fertility Rates and Indices by Area, 1993

| Age | | MCH-FP | area | Con | parison | area |
|----------|--------|--------|-------|--------|----------|-------|
| (years) | Births | Women | Rate | Births | Women | Rate |
| All ages | 2615 | 26633 | 98.2 | 2992 | 23570 | 126.9 |
| 15-19* | 303 | 5474 | 55.4 | 346 | 4855 | 71.3 |
| 20-24 | 894 | 5118 | 174.7 | 1011 | 4514 | 224.0 |
| 25-29 | 803 | 4741 | 169.4 | 862 | 4097 | 210.4 |
| 30-34 | 429 | 3926 | 109.3 | 509 | 3509 | 145.1 |
| 35-39 | 142 | 2847 | 49.9 | 205 | 2594 | 79.0 |
| 40 - 44 | 42 | 2242 | 18.7 | 47 | 1956 | 24.0 |
| 45-49** | 2 | 2285 | 0.9 | 12 | 2045 | 5.9 |
| | TFR | = | 2891 | TFR | = | 3798 |
| | GFR | = | 98 | GFR | = | 127 |
| | GRR | = | 1404 | GRR | = 0 | 1913 |
| | NRR | = | 1250 | NRR | \equiv | 1597 |

^{*}Births to mothers under age 15 were included in this group. **Births to mothers age 50 and above were included in this group.

Figure 4.2: Age-specific Fertility Rates by Area, 1993




Table 4.5: Age-specific Fertility Rates and Indices for MCH-FP Area by Block, 1993

| | | Block A | | | Block B | |
|----------------|--------|---------|-------|--------|---------|-------|
| Age (years) | Births | Women | Rate | Births | Women | Rate |
| All ages | 749 | 7568 | 99.0 | 681 | 6620 | 102.9 |
| 15-19* | 73 | 1561 | 46.8 | 73 | 1337 | 54.6 |
| 20-24 | 243 | 1407 | 172.7 | 233 | 1288 | 180.9 |
| 25-29 | 252 | 1417 | 177.8 | 212 | 1181 | 179.5 |
| 30-34 | 131 | 1119 | 117.1 | 107 | 945 | 113.2 |
| 35-39 | 38 | 806 | 47.1 | 39 | 698 | 55.9 |
| 40-44 | 12 | 612 | 19.6 | 15 | 585 | 25.6 |
| 45-49** | 0 | 646 | 0.0 | 2 | 586 | 3.4 |
| | TFR | = | 2906 | TFR | = | 3066 |
| | GFR | = | 99 | GFR | = | 103 |
| | GRR | = | 1411 | GRR | = | 1489 |

(continued)

Table 4.5 (cont.): Age-specific Fertility Rates and Indices for MCH-FP Area by Block, 1993

| | | Block C | | | Block D | |
|----------------|--------|---------|-------|--------|---------|-------|
| Age (years) | Births | Women | Rate | Births | Women | Rate |
| All ages | 636 | 6901 | 92.2 | 549 | 5544 | 99.0 |
| 15-19* | 86 | 1444 | 59.6 | 71 | 1132 | 62.7 |
| 20-24 | 225 | 1383 | 162.7 | 193 | 1040 | 185.6 |
| 25-29 | 181 | 1152 | 157.1 | 158 | 991 | 159.4 |
| 30-34 | 99 | 980 | 101.0 | 92 | 882 | 104.3 |
| 35-39 | 36 | 783 | 46.0 | 29 | 560 | 51.8 |
| 40-44 | 9 | 587 | 15.3 | 6 | 458 | 13.1 |
| 45-49** | 0 | 572 | 0.0 | 0 | 481 | 0.0 |
| D | TFR | = | 2708 | TFR | = | 2885 |
| | GFR | = | 92 | GFR | = | 99 |
| | GRR | = | 1315 | GRR | = | 1401 |
| | | | | | | |

^{*}Births to mothers under age 15 were included in this group. **Births to mothers age 50 and above were included in this group.

Table 4.6: Births by Mother's Age, Live-birth Order and Area, 1993

| Age (years) | Total | Total | - | | | | Live | e-birth | order | | | |
|----------------|-------|--------|-----|------|-----|-----|------|---------|-------|----|-----|------|
| - (Jears) | women | births | | 1 2 | 2 | 3 | 4 | 5 | 6 | 7 | 8 9 | 9 10 |
| Both ar | eas | | | | | | | | | | | |
| <15 | 12010 | 5 | į | · . | | | | | | | | |
| 15-19 | 10329 | 646 | 577 | | | | | • | | | | |
| 20-24 | 9632 | 1904 | 834 | | 287 | | | | | | | |
| 25-29 | 8838 | 1664 | 117 | | 523 | | - 15 | 9 1 | | | | - |
| 30-34 | 7435 | 938 | 12 | 77.7 | 110 | | | | | 5 | 1 | 0 |
| 35-39 | 5441 | 348 | 3 | 3 | 10 | | | | 85 | 46 | 11 | 7 |
| 40-44 | 4198 | 89 | 0 | 1 | 2 | | | | 69 | 51 | 43 | 24 |
| 45-49 | 4330 | 13 | 1 | 0 | | | | | 10 | 16 | 21 | 22 |
| MCH-FP a | 200 | | | U | 0 | 0 | 1 | 0 | 2 | 2 | 2 | 5 |
| mun-rp a | rea | | | | | | | | | | | |
| <15 | 5944 | 1 | 1 | | | | | | | | | |
| 15-19 | 5474 | 302 | 287 | 13 | | • | | | | | 21 | |
| 20-24 | 5118 | 894 | 429 | 342 | 2 | | | - | | | | |
| 25-29 | 4741 | 802 | 74 | 197 | 99 | 22 | 2 | | ** | | | |
| 30-34 | 3926 | 430 | 7 | 28 | 284 | 160 | 64 | 17 | 6 | 0 | | |
| 35-39 | 2847 | 142 | 2 | | 66 | 120 | 109 | 66 | 21 | 8 | 2 | 3 |
| 0-44 | 2242 | 42 | 0 | 1 | 6 | 27 | 25 | 32 | 24 | 10 | 11 | 4 |
| 5-49 | 2285 | 2 | 0 | | 2 | 1 | 4 | 7 | 7 | 6 | 10 | 4 |
| | | _ | U | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 0 |
| omparisor | area | | | | | | | | | | | |
| <15 | 6066 | 4 | 4 | | | | | | | | | |
| 5-19 | 4855 | 344 | 290 | E1 | | | 1.75 | | 27 | | | ¥ |
| -24 | 4514 | 1010 | | 51 | 2 | 1 | | - | | * | *1 | 2 |
| -29 | 4097 | 862 | | | 188 | 38 | 7 | 1 | • | | | |
| -34 | 3509 | 508 | 5 | | 239 | 225 | 141 | 57 | 13 | 5 | 1 | 0 |
| -39 | 2594 | 206 | 1 | 15 | 44 | | 129 | 119 | 64 | 38 | 9 | 4 |
| -44 | 1956 | 47 | 0 | 2 | 4 | 10 | 23 | 28 | 45 | 41 | 32 | 20 |
| -49 | 2045 | 11 | | 0 | 0 | 1 | 1 | 3 | 3 | 10 | 11 | 18 |
| | | 11 | 1 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 2 | 5 |

Table 4.7: Age-order-specific Fertility Rates by Area, 1993

| 1000000 | | | | | Liv | e-birth o | rder | | | | |
|----------------|---------|--------|--------|--------|--------|-----------|--------|--------|--------|--------|--------|
| Age (years) | Total | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10+ |
| Both area | s | | | | | | | | | | |
| <15 | 0.0004 | 0.0004 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 |
| 15-19 | 0.0625 | 0.0559 | 0.0062 | 0.0004 | 0.0001 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 |
| 20-24 | 0.1977 | 0.0866 | 0.0740 | 0.0298 | 0.0062 | 0.0009 | 0.0001 | 0.0000 | 0.0000 | 0.0000 | 0.0000 |
| 25-29 | 0.1883 | 0.0132 | 0.0379 | 0.0592 | 0.0436 | 0.0232 | 0.0084 | 0.0021 | 0.0006 | 0.0001 | 0.0000 |
| 30-34 | 0.1262 | 0.0016 | 0.0058 | 0.0148 | 0.0270 | 0.0320 | 0.0249 | 0.0114 | 0.0062 | 0.0015 | 0.0009 |
| 35-39 | 0.0640 | 0.0006 | 0.0006 | 0.0018 | 0.0068 | 0.0088 | 0.0110 | 0.0127 | 0.0094 | 0.0079 | 0.0044 |
| 40-44 | 0.0212 | 0.0000 | 0.0002 | 0.0005 | 0.0005 | 0.0012 | 0.0024 | 0.0024 | 0.0038 | 0.0050 | 0.0052 |
| 45-49 | 0.0030 | 0.0002 | 0.0000 | 0.0000 | 0.0000 | 0.0002 | 0.0000 | 0.0005 | 0.0005 | 0.0005 | 0.0012 |
| Total | 3.3162 | 0.7925 | 0.6235 | 0.5323 | 0.4210 | 0.3319 | 0.2338 | 0.1455 | 0.1020 | 0.0748 | 0.0587 |
| MCH-FP a | rea | | | | | | | | | | |
| <15 | 0.0002 | 0.0002 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.000 |
| 15-19 | 0.0552 | 0.0524 | 0.0024 | 0.0004 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.000 |
| 20-24 | 0.1747 | 0.0838 | 0.0668 | 0.0193 | 0.0043 | 0.0004 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.000 |
| 25-29 | 0.1692 | 0.0156 | 0.0416 | 0.0599 | 0.0337 | 0.0135 | 0.0036 | 0.0013 | 0.0000 | 0.0000 | 0.000 |
| 30-34 | 0.1095 | 0.0018 | 0.0071 | 0.0168 | 0.0306 | 0.0278 | 0.0168 | 0.0053 | 0.0020 | 0.0005 | 0.000 |
| 35-39 | 0.0499 | 0.0007 | 0.0004 | 0.0021 | 0.0095 | 0.0088 | 0.0112 | 0.0084 | 0.0035 | 0.0039 | 0.001 |
| 40-44 | 0.0187 | 0.0000 | 0.0004 | 0.0009 | 0.0004 | 0.0018 | 0.0031 | 0.0031 | 0.0027 | 0.0045 | 0.001 |
| 45-49 | 0.0009 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0004 | 0.0000 | 0.0000 | 0.0004 | 0.0000 | 0.000 |
| Total | 2.8910 | 0.7726 | 0.5934 | 0.4971 | 0.3927 | 0.2633 | 0.1738 | 0.0908 | 0.0433 | 0.0442 | 0.019 |
| Comparis | on area | | | | | | | | | | |
| <15 | 0.0007 | 0.0007 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.000 |
| 15-19 | 0.0709 | 0.0597 | 0.0105 | 0.0004 | 0.0002 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.000 |
| 20-24 | 0.2237 | 0.0897 | 0.0822 | 0.0416 | 0.0084 | 0.0016 | 0.0002 | 0.0000 | 0.0000 | 0.0000 | 0.000 |
| 25-29 | 0.2104 | 0.0105 | 0.0337 | 0.0583 | 0.0549 | 0.0344 | 0.0139 | 0.0032 | 0.0012 | 0.0002 | 0.00 |
| 30-34 | 0.1448 | 0.0014 | 0.0043 | 0.0125 | 0.0231 | 0.0368 | 0.0339 | 0.0182 | 0.0108 | 0.0026 | 0.00 |
| 35-39 | 0.0794 | 0.0004 | 0.0008 | 0.0015 | 0.0039 | 0.0089 | 0.0108 | 0.0173 | 0.0158 | 0.0123 | 0.00 |
| 40-44 | 0.0240 | 0.0000 | 0.0000 | 0.0000 | 0.0005 | 0.0005 | 0.0015 | 0.0015 | 0.0051 | 0.0056 | 0.00 |
| 45-49 | 0.0054 | 0.0005 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0000 | 0.0010 | 0.0005 | 0.0010 | 0.00 |
| Total | 3.7963 | 0.8145 | 0.6571 | 0.5724 | 0.4550 | 0.4105 | 0.3019 | 0.2064 | 0.1673 | 0.1087 | 0.102 |

CHAPTER 5

MARRIAGE AND DIVORCE

The number of marriages registered in 1993 was 2,697, giving a crude marriage rate of 13.0 per thousand. These figures show a decline on those of 1992, but are still above the low figure of 2,540, or 12.7 per thousand, recorded in 1988.

Tables 5.1 and 5.2 show the distributions of grooms and brides by age at marriage and previous marital status. The mean ages at marriage -- 27.1 and 19.4 for all grooms and brides; 25.8 and 18.6 for those marrying for the first time -- show no change on 1992. In general, however, there appears to be rise in age at marriage of females in Matlab: the mean age has been over 18 for every year since 1984, while prior to that date it was consistently below that age.

Table 5.5 shows that divorces numbered 334 in 1993, which constituted a decrease from the 1992 figure of 439. In general, the incidence of divorce in Matlab appears to have fallen: during the first four years of the present study, from 1978 to 1981, the numbers of divorces were consistently over 500; since 1981 they have been below that figure.

Table 5.5, showing the number of divorces by partner's age, indicates that the peak ages of divorce for men are 25-29, compared to 15-24 for women. This reflects the sex difference in age at marriage.

Table 5.6 shows the distributions of marriages and divorces by month. June appears to be a peak month for marriages, but no regular seasonal pattern is discernible in the number of divorces.

Table 5.7 gives data on divorces by marriage duration, sex and age. The largest percentage of divorces occurs among couples married twelve to twenty-three months.

It may be noted that the DSS definitions specify that if either partner in a marriage has been resident on the study area, the marriage should be registered. Thus a bride from the study area marries a groom from outside the area, the marriage will be included in the marriage statistics, though she immediately moves out of the area and all here subsequent childbearing goes unrecorded by the DSS. These procedures lead to an imbalance between the numbers of marriages and the numbers of births, and caution is needed if for any reason the two data sets have to be related.

Table 5.1: Groom's Age at Marriage by Previous Marital Status, 1993

| | | | | | Pre | vious mari | tal st | atus | | |
|----------------|------|---------|--------|---------|-----|------------|--------|---------|-----|---------|
| Age (years) | A11 | grooms | S | ingle | M | larried | D | ivorced | h | lidowed |
| (Jears) | No. | Percent | No. | Percent | No. | Percent | No. | Percent | No. | Percent |
| All ages | 2697 | 100.0 | 2250 | 100.0 | 85 | 100.0 | 265 | 100.0 | 97 | 100.0 |
| 10-14 | 1 | 0.0 | 1 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 117 | 4.3 | 111 | 4.9 | 1 | 1.2 | 5 | 1.9 | 0 | 0.0 |
| 20-24 | 858 | 31.8 | 778 | 34.6 | 13 | 15.3 | 65 | 24.5 | 2 | 2.1 |
| 25-29 | 1069 | 39.6 | 960 | 42.7 | 20 | 23.5 | 76 | 28.7 | 13 | 13.4 |
| 30-34 | 461 | 17.1 | 352 | 15.6 | 17 | 20.0 | 69 | 26.0 | 23 | 23.7 |
| 35-39 | 96 | 3.6 | 41 | 1.8 | 13 | 15.3 | 23 | 8.7 | 19 | 19.6 |
| 40-44 | 30 | 1.1 | 5 | 0.2 | 8 | 9.4 | 6 | 2.3 | 11 | 11.3 |
| 45-49 | 22 | 0.8 | 1 | 0.0 | 5 | 5.9 | 7 | 2.6 | 9 | 9.3 |
| 50-54 | 16 | 0.6 | 0 | 0.0 | 3 | 3.5 | 5 | 1.9 | 8 | 8.2 |
| 55-59 | 5 | 0.2 | 0 | 0.0 | 1 | 1.2 | 2 | 0.8 | 2 | 2.1 |
| 60-64 | 13 | 0.5 | 0 | 0.0 | 2 | 2.4 | 4 | 1.5 | 7 | 7.2 |
| 65+ | 8 | 0.3 | 0 | 0.0 | 2 | 2.4 | 3 | 1.1 | 3 | 3.1 |
| Unknown | 1 | 0.0 | 1 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Median age* | 2 9 | 26.0 | 9 5 | 25.0 | | 31.0 | | 28.0 | | 36.0 |
| Mean age* | 9 | 27.1 | ž R | 25.8 | | 34.8 | (| 30.4 | | 39.7 |
| Standard de | v.* | 6.4 | | 4.1 | | 11.4 | | 9.0 | | 11.1 |

^{*}Mean, median, and standard deviation were calculated from ungrouped age data.

Table 5.2: Bride's Age at Marriage by Previous Marital Status, 1993

| | | | | Previo | us mar | ital statu | IS | |
|------------|----------|---------|------|---------|--------|------------|-----|---------|
| Age | A11 | brides | S | ingle | Di | vorced | Wi | dowed |
| (years) | No. | Percent | No. | Percent | No. | Percent | No. | Percent |
| All ages | 2697 | 100.0 | 2363 | 100.0 | 299 | 100.0 | 35 | 100.0 |
| 10-14 | 112 | 4.2 | 112 | 4.7 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 1561 | 57.9 | 1485 | 62.8 | 75 | 25.1 | 1 | 2.9 |
| 20-24 | 790 | 29.3 | 661 | 28.0 | 119 | 39.8 | 10 | 28.6 |
| 25-29 | 167 | 6.2 | 90 | 3.8 | 68 | 22.7 | 9 | 25.7 |
| 30-34 | 37 | 1.4 | 12 | 0.5 | 21 | 7.0 | 4 | 11.4 |
| 35-39 | 13 | 0.5 | 0 | 0.0 | 11 | 3.7 | 2 | 5.7 |
| 40-44 | 7 | 0.3 | 0 | 0.0 | 3 | 1.0 | 4 | 11.4 |
| 45+ | 7 | 0.3 | 0 | 0.0 | 2 | 0.7 | 5 | 14.3 |
| Unknown | 3 | 0.1 | 3 | 0.1 | 0 | 0.0 | 0 | 0.0 |
| Median age | <u>*</u> | 19.0 | | 18.0 | | 23.0 | | 29.0 |
| Mean age* | | 19.4 | | 18.6 | | 23.7 | | 31.7 |
| Standard o | dev.* | 4.2 | | 3.0 | | 5.7 | | 10.5 |

^{*}Mean, median, and standard deviation were calculated from ungrouped age data.

Table 5.3: Marriage Rates by Age and Sex, 1993

| Age | | Males | | | Females | |
|---------|-----------|------------|-------|-----------|------------|-------------------|
| (years) | Marriages | Population | Rate* | Marriages | Population | Rate ² |
| 10-14 | 1 | 13525 | 0.1 | 112 | 12010 | 9.3 |
| 15-19 | 117 | 11576 | 10.1 | 1561 | 10329 | 151.1 |
| 20-24 | 858 | 9068 | 94.6 | 790 | 9632 | 82.0 |
| 25-29 | 1069 | 6855 | 155.9 | 167 | 8838 | 18.9 |
| 30-34 | 461 | 7247 | 63.6 | 37 | 7435 | 5.0 |
| 35-39 | 96 | 5400 | 17.8 | 13 | 5441 | 2.4 |
| 40-44 | 30 | 4003 | 7.5 | 7 | 4198 | 1.7 |
| 45+ | 65 | 17059 | 3.8 | 10 | 18245 | 0.5 |

^{*}Rates per 1000 population irrespective of previous marital status.

Table 5.4: Number of Marriages by Groom's and Bride's Age at Marriage, 1993

| Groom's | | | | | Bride's | age (y | ears) | | | |
|----------------|------|-------|-------|-------|---------|--------|-------|-------|-----|---------|
| age (years) | A11 | 10-14 | 15-19 | 20-24 | 25-29 | 30-34 | 35-39 | 40-44 | 45+ | Unknowr |
| All ages | 2697 | 112 | 1561 | 790 | 167 | 37 | 13 | 7 | 7 | 3 |
| 10-14 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-19 | 117 | 12 | 84 | 17 | 2 | 1 | 0 | 0 | 0 | 1 |
| 20-24 | 858 | 58 | 599 | 183 | 16 | 1 | 0 | 0 | 0 | 1 |
| 25-29 | 1069 | 32 | 635 | 351 | 45 | 5 | 1 | 0 | 0 | 0 |
| 30-34 | 461 | 9 | 208 | 180 | 56 | 7 | 1 | 0 | 0 | 0 |
| 35-39 | 96 | 1 | 29 | 38 | 21 | 3 | 4 | 0 | 0 | 0 |
| 40-44 | 30 | 0 | 0 | 9 | 12 | 6 | 2 | 1 | 0 | 0 |
| 45-49 | 22 | 0 | 3 | 5 | 7 | 5 | 1 | 0 | 1 | 0 |
| 50-54 | 16 | 0 | 1 | 4 | 4 | 4 | 1 | 1 | 1 | 0 |
| 55-59 | 5 | 0 | 0 | 1 | 2 | 0 | 0 | 1 | 1 | 0 |
| 60-64 | 13 | 0 | 1 | 1 | 2 | 4 | 3 | 2 | 0 | 0 |
| 65+ | 8 | 0 | 0 | 1 | 0 | 1 | 0 | 2 | 4 | 0 |
| Unknown | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |

Table 5.5: Number of Divorces by Partners' Age at Divorce, 1993

| Male's | | | | F | emale's | age (y | rears) | | | |
|----------------|-----|-------|-------|-------|---------|--------|--------|-------|-------|-----|
| age (years) | A11 | 10-14 | 15-19 | 20-24 | 25-29 | 30-34 | 35-39 | 40-44 | 45-49 | 50+ |
| All ages | 334 | 5 | 101 | 122 | 70 | 20 | 7 | 8 | 1 | 0 |
| 10-14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15-19 | 9 | 1 | 7 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 20-24 | 69 | 1 | 35 | 30 | 3 | 0 | 0 | 0 | 0 | 0 |
| 25-29 | 97 | 2 | 39 | 45 | 10 | 0 | 1 | 0 | 0 | 0 |
| 30-34 | 73 | 1 | 13 | 31 | 26 | 2 | 0 | 0 | 0 | 0 |
| 35-39 | 40 | 0 | 4 | 9 | 21 | 5 | 1 | 0 | 0 | 0 |
| 40-44 | 15 | 0 | 1 | 3 | 4 | 6 | 1 | 0 | 0 | 0 |
| 45-49 | 7 | 0 | 1 | 0 | 2 | 1 | 1 | 2 | 0 | 0 |
| 50-54 | 15 | 0 | 1 | 2 | 2 | 4 | 3 | 3 | 0 | 0 |
| 55-59 | 5 | 0 | 0 | 0 | 2 | 0 | 0 | 2 | 1 | 0 |
| 60-64 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| 65+ | 3 | 0 | 0 | 1 | 0 | 1 | 0 | 1 | 0 | 0 |

Table 5.6: Marriages and Divorces by Months, 1993

| March In | Marr | iage | Div | rorce |
|-----------|--------|---------|--------|---------|
| Month | Number | Percent | Number | Percent |
| January | 188 | 7.0 | 39 | 11.7 |
| February | 272 | 10.1 | 27 | 8.1 |
| March | 248 | 9.2 | 32 | 9.6 |
| April | 191 | 7.1 | 21 | 6.3 |
| May | 137 | 5.1 | 19 | 5.7 |
| June | 329 | 12.2 | 44 | 13.2 |
| July | 281 | 10.4 | 26 | 7.8 |
| August | 242 | 9.0 | 16 | 4.8 |
| September | 185 | 6.9 | 30 | 9.0 |
| October | 245 | 9.1 | 37 | 11.1 |
| November | 159 | 5.9 | 24 | 7.2 |
| December | 220 | 8.2 | 19 | 5.7 |
| Total | 2697 | 100.0 | 334 | 100.0 |

Figure 5.1: Marriages and Divorces by Month, 1993




Table 5.7: Number of Divorces by Sex, Age, and Duration of Marriage, 1993

| | | | | | | | Dur | Duration of marriage (months) | marri | age (mon | ths) | | | | | |
|----------|------|--------------|------|---------|------|--------|------|-------------------------------|-------|----------|------|--------|------|--------|------|--------|
| Age at | All | All duration | | Under 6 | 9 | 6-11 | 12 | 12-23 | 24 | 24-35 | 36 | 36-47 | 48 | 48-59 | | +09 |
| divorce | Male | Male Female | Male | Female | Male | Female | Male | Female | Male | Female | Male | Female | Male | Female | Male | Female |
| All ages | 334 | 334 | 37 | 37 | 49 | 49 | 88 | 88 | 51 | 51 | 31 | 31 | 17 | 17 | 61 | 61 |
| Under 20 | 6 | 106 | 0 | 13 | 80 | 30 | - | 41 | 0 | 14 | 0 | 7 | 0 | Н | 0 | 0 |
| 20-24 | 69 | 122 | 9 | 17 | 14 | 15 | 28 | 31 | 11 | 21 | 9 | 17 | 1 | 6 | 3 | 12 |
| 25-29 | 97 | 70 | 16 | 4 | 15 | 2 | 29 | 13 | 15 | 12 | 10 | 2 | 4 | 4 | 00 | 30 |
| 30-34 | 73 | 20 | 11 | 2 | 7 | 1 | 13 | П | 16 | 2 | 11 | 2 | 2 | Н | 10 | 11 |
| 35-39 | 40 | 7 | П | 0 | П | 0 | 12 | 2 | 3 | 2 | П | 0 | 3 | Н | 19 | 2 |
| 40-49 | 22 | 6 | 0 | г | 2 | 1 | 3 | 0 | 0 | 0 | 2 | 0 | 4 | Н | 11 | 9 |
| +09 | 24 | 0 | 3 | 0 | 2 | 0 | 2 | 0 | 9 | 0 | П | 0 | 0 | 0 | 10 | 0 |
| | | | | | | | | | | | | | | | | |

CHAPTER 6

MIGRATION

An "out-migrant" is defined as a person originally listed on a DSS census as a resident, or a person who became a resident after the census by birth or immigration, who subsequently moved out of the surveillance (DSS) area permanently. Likewise, an "inmigrant" is an individual not recorded in the last census who has permanently moved into the surveillance area. Those who stay in the area continuously for at least six months in a year or come home at least once a month to stay overnight are treated as permanent residents. It may be noted that these definitions refer to the surveillance areas as a whole. People who move from the Comparison area into the MCH-FP area, or vice versa, do not feature in the tables which show the numbers of migrants in the two areas.

Table 6.1 shows that the number of in-migrants in 1993 was 5,292 giving a crude rate of in-migration of 25.5 per thousand. Out-migrants numbered 7,492 and the out-migration rate 36.1 per thousand. Both these figures are appreciably lower than those of 1992. The net loss of migrants decreased from 3,069 in 1992 to 2,200 in 1993. Female in-migrants once again outnumbered male, both among those coming in and those going out.

The numbers for the MCH-FP and Comparison areas, shown in Tables 6.2 and 6.3 by age and sex, are fairly evenly distributed between the two areas.

Tables 6.4 shows the age-specific migration rates, which are illustrated in Figure 6.1. They show the bi-modal age distributions commonly found for migrant populations, with a primary peak of young adults and a secondary peak of young children moving with their parents. For males the ages of the out-migrants tended to be rather younger than those of the in-migrants, while for females the shapes of the distributions were similar.

Table 6.5 to 6.8 show the distributions of in- and out-migrants by age, sex and the cause of the movement.

Table 6.9 and Figure 6.2 show the numbers moving in and out by month. The apparent deficit of migrants in April and May, followed by the upsurge in June, calls for comment. Although June is often a peak month for such movements, the 1993 configuration is so pronounced that the possibility of mis-dating must be considered: entries and exits which had occurred earlier in the year may have been recorded as taking place in June. Such a situation may have arisen as a result of the additional workload imposed on the CHWs by the 1993 census.

Tables 6.10 and 6.11 show the number of males and females migrating in and out of the Matlab DSS area by location of origin or destination. For locations in Bangladesh the Division and whether the location is rural or urban is given. For numerically important origins or destinations the District is also shown. Roughly equal numbers of men and women move into and out of rural areas of Chandpur district, neighboring Matlab, probably due to marriage. There is a net loss of both men and women to urban Dhaka, primarily of young adults. There is also a net loss to India, but more evenly distributed across age groups. Migration to the Middle-East and other Asian locations is heavily concentrated among out-migrating males age 15-34.

Table 6.1: In- and Out-migration by Age and Sex, 1993

| Name of | In-m | nigration |) | Out- | migratio | n |
|-----------------------|---------------------------------|-----------------------------|-----------------------------|---------------------------------|----------------------------|-----------------------------|
| Age (years) | Both sexes | Males | Females | Both sexes | Males | Females |
| All ages | 5292 | 2416 | 2876 | 7492 | 3707 | 3785 |
| Under 5 | 835 | 436 | 399 | 890 | 428 | 462 |
| 0 1 2 3 4 | 234 169 153 125 154 | 124 82 78 69 83 | 110 87 75 56 71 | 189 183 181 172 165 | 92 89 74 89 84 | 97 94 107 83 81 |
| 5 - 9 | 545 | 277 | 268 | 611 | 304 | 307 |
| 10-14 | 394 | 173 | 221 | 579 | 288 | 291 |
| 15-19 | 869 | 164 | 705 | 1421 | 468 | 953 |
| 20-24 | 780 | 210 | 570 | 1501 | 671 | 830 |
| 25-29 | 597 | 275 | 322 | 1010 | 574 | 436 |
| 30-34 | 471 | 335 | 136 | 610 | 415 | 195 |
| 35-39 | 288 | 216 | 72 | 308 | 232 | 76 |
| 40 - 44 | 146 | 107 | 39 | 167 | 123 | 44 |
| 45-49 | 103 | 76 | 27 | 101 | 61 | 40 |
| 50-54 | 85 | 56 | 29 | 86 | 45 | 41 |
| 55-59 | 64 | 37 | 27 | 57 | 26 | 31 |
| 60-64 | 45 | 24 | 21 | 55 | 32 | 23 |
| 65+ | 70 | 30 | 40 | 96 | 40 | 56 |

Table 6.2: In-migration by Age, Sex, and Area, 1993

| Age | MCH- | FP area | | Compa | rison a | rea |
|-----------------------|-----------------------------|----------------------------|----------------------------|-----------------------------|----------------------------|----------------------------|
| (years) | Both sexes | Males | Females | Both sexes | Males | Females |
| All ages | 2667 | 1210 | 1457 | 2625 | 1206 | 1419 |
| Under 5 | 418 | 222 | 196 | 417 | 214 | 203 |
| 0 1 2 3 4 | 130 85 71 60 72 | 67 46 34 36 39 | 63 39 37 24 33 | 104 84 82 65 82 | 57 36 44 33 44 | 47 48 38 32 38 |
| 5 - 9 | 278 | 131 | 147 | 267 | 146 | 121 |
| 10-14 | 202 | 90 | 112 | 192 | 83 | 109 |
| 15-19 | 447 | 74 | 373 | 422 | 90 | 332 |
| 20-24 | 391 | 109 | 282 | 389 | 101 | 288 |
| 25-29 | 299 | 127 | 172 | 298 | 148 | 150 |
| 30-34 | 238 | 173 | 65 | 233 | 162 | 71 |
| 35-39 | 153 | 113 | 40 | 135 | 103 | 32 |
| 40 - 44 | 85 | 64 | 21 | 61 | 43 | 18 |
| 45-49 | 53 | 44 | 9 | 50 | 32 | 18 |
| 50-54 | 33 | 24 | 9 | 52 | 32 | 20 |
| 55-59 | 30 | 18 | 12 | 34 | 19 | 15 |
| 60 - 64 | 17 | 10 | 7 | 28 | 14 | 14 |
| 65+ | 23 | 11 | 12 | 47 | 19 | 28 |


Table 6.3: Out-migration by Age, Sex, and Area, 1993

| 2 | MCH- | FP area | | Compa | rison ar | 'ea |
|-----------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| Age (years) | Both sexes | Males | Females | Both sexes | Males | Females |
| All ages | 3763 | 1822 | 1941 | 3729 | 1885 | 1844 |
| Under 5 | 420 | 206 | 214 | 470 | 222 | 248 |
| 0 1 2 3 4 | 94 86 85 84 71 | 50 42 37 39 38 | 44 44 48 45 33 | 95 97 96 88 94 | 42 47 37 50 46 | 53 50 59 38 48 |
| 5 - 9 | 312 | 153 | 159 | 299 | 151 | 148 |
| 10-14 | 270 | 126 | 144 | 309 | 162 | 147 |
| 15-19 | 712 | 232 | 480 | 709 | 236 | 473 |
| 20-24 | 768 | 316 | 452 | 733 | 355 | 378 |
| 25-29 | 513 | 283 | 230 | 497 | 291 | 206 |
| 30-34 | 311 | 207 | 104 | 299 | 208 | 91 |
| 35-39 | 165 | 123 | 42 | 143 | 109 | 34 |
| 40-44 | 95 | 76 | 19 | 72 | 47 | 25 |
| 45-49 | 52 | 33 | 19 | 49 | 28 | 21 |
| 50-54 | 40 | 15 | 25 | 46 | 30 | 16 |
| 55-59 | 28 | 15 | 13 | 29 | 11 | 18 |
| 60-64 | 26 | 15 | 11 | 29 | 17 | 12 |
| 65+ | 51 | 22 | 29 | 45 | 18 | 27 |

Table 6.4: Age and Sex-specific Migration Rates by Direction, 1993 (per 1000 population)

| | Both | sexes | Ma | les | Fe | males |
|-----------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|
| Age (years) | In | Out | In | Out | In | Out |
| All ages | 25.5 | 36.1 | 23.3 | 35.8 | 27.6 | 36.3 |
| Under 5 | 29.6 | 31.5 | 30.7 | 30.1 | 28.5 | 33.0 |
| 0 1 2 3 4 | 42.7 33.6 26.2 21.8 25.0 | 34.5 36.3 31.0 30.1 26.8 | 44.4 32.2 26.3 23.9 27.4 | 33.0 34.9 25.0 30.8 27.8 | 40.8 35.0 26.1 19.8 22.7 | 36.0 37.8 37.2 29.3 25.9 |
| 5 - 9 | 19.0 | 21.4 | 19.0 | 20.8 | 19.1 | 21.9 |
| 10-14 | 15.4 | 22.7 | 12.8 | 21.3 | 18.4 | 24.2 |
| 15-19 | 39.7 | 64.9 | 14.2 | 40.4 | 68.3 | 92.3 |
| 20-24 | 41.7 | 80.3 | 23.2 | 74.0 | 59.2 | 86.2 |
| 25-29 | 38.0 | 64.4 | 40.1 | 83.7 | 36.4 | 49.3 |
| 30-34 | 32.1 | 41.5 | 46.2 | 57.3 | 18.3 | 26.2 |
| 35-39 | 26.6 | 28.4 | 40.0 | 43.0 | 13.2 | 14.0 |
| 40-44 | 17.8 | 20.4 | 26.7 | 30.7 | 9.3 | 10.5 |
| 45-49 | 13.3 | 13.0 | 22.1 | 17.8 | 6.2 | 9.2 |
| 50-54 | 11.4 | 11.6 | 16.9 | 13.6 | 7.1 | 10.0 |
| 55-59 | 9.6 | 8.6 | 11.3 | 7.9 | 8.0 | 9.2 |
| 60-64 | 8.6 | 10.5 | 9.1 | 12.1 | 8.2 | 9.0 |
| 65+ | 8.5 | 11.6 | 6.9 | 9.1 | 10.3 | 14.5 |

Figure 6.1: Rate of In- and Out-migration by Sex and Age, 1993






Table 6.5: Male Out-migration by Cause of Movement and Age, 1993

| All migrants 3707 Work/Economic/Educational 2064 -acquired/seeking job job completion/retirement 17 -to acquire education 2264 | | | | | | | | | | | | | | |
|--|------------------------------------|--------|--------------------|---------------------|--------------------------|---------------------|-------------------------|--------------------|---------------|-------------|---------|-------------------|--------|--------------|
| 2000 | 11 <5 | 5-9 | 10-14 | 15-19 | 20-24 | 25-29 | 30-34 | 35-39 | 40-44 | 45-49 | 50-54 | 55-59 | 60-64 | 65+ |
| | 7 428 | 304 | 288 | 468 | 671 | 574 | 415 | 232 | 123 | 61 | 45 | 56 | 32 | 40 |
| nterrupt | 2064 17 235 235 1 1 | mom000 | 51 41 0 | 263 3 82 0 | 517 2 66 0 0 | 483 27 0 0 | 350 3 0 0 1 | 197 0 0 0 | 0000 | 9400000 | 7000108 | 15 0 0 0 | 000000 | 910000 |
| Housing/Environmental -acquired/seeking new 110 land/house 2 | 5.0 | 1 0 | 0 33 | 40 | 10 | 14 | 14 | 15 | 16 | 80 O | 9 1 | 4 1 | 90 | 90 |
| e/widow | 3 0 | 00 | 00 | 00 | 01 | 00 | 2 0 | 00 | 00 | 00 | 00 | 00 | 00 | 00 |
| Figure 1202 Figure | 2 425 1 1 8 0 | 286 | 189 0 0 1 | 110 0 2 0 | 0300 | 0 0 0 0 | 0 4 0 | 11 0 0 0 | 8000 | 4000 | m000 | 4000 | 0014 | 19 0 7 |
| Legal problems 12 | 2 0 | 0 | 0 | 0 | 2 | 0 | 4 | П | П | П | 1 | П | 1 | 0 |
| Other and not stated 25 -other n.e.c.* 25 -unknown or not stated 1 | 1 0 | 10 | 0.0 | 00 | e 0 | 10 | , 40 | 40 | \rightarrow | 0 5 | 0.0 | 00 | 10 | 10 |

Table 6.6: Female Out-migration by Cause of Movement and Age, 1993

| Total <pre></pre> | Carica of movement | | | | | | | A | Age (years | rs) | | | | | | |
|---|--|------------------------|----------------|--------|------------------------|--------------------------|--------------------------|---------------|---------------|---------------|--------|----------|--------|--------|--------|--|
| terrupt | בממפה כו וויסאבוויבורב | Total | <5× | 6-9 | 10-14 | 15-19 | 20-24 | 25-29 | 30-34 | 35-39 | 40-44 | 45-49 | 50-54 | 55-59 | 60-64 | |
| Tement 100 4 43 0 4 61 181 102 55 18 12 1 1 1 1 2 1 1 1 1 1 1 1 1 1 1 1 1 | All migrants | 3785 | 462 | 307 | 291 | 953 | 830 | 436 | 195 | 76 | 44 | 40 | 41 | 31 | 23 | |
| Hardow 702 0 0 19 377 242 49 14 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | Work/Economic/Educational -acquired/seeking job -job completion/retirement -to acquire education -educ. completed/interrupt -student lodging | 443 10 96 0 | 000000 | 400000 | 26 0 0 0 0 | 181 1 34 0 0 | 102 2 18 0 0 | 00000 | 180000 | 1200000 | HOHOOO | нооооо | N00000 | нооооо | нонооо | |
| Avidow 702 0 0 19 377 242 49 14 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | | 43 | 10 | 0.0 | Н0 | 00 | 110 | 7 0 | 10 | \bowtie | 10 | 23 | 00 | 10 | 10 | |
| 2337 448 292 175 326 402 296 146 58 40 32 37 24 13 13 13 0 0 0 2 9 1 1 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | 2. | 702 | 00 | 00 | 19 | 377 | 242 | 18 | 14 | \rightarrow | 0 H | 00 | 00 | 00 | 00 | |
| ted $\begin{array}{cccccccccccccccccccccccccccccccccccc$ | -move with or join spouse/parents -adoption -family friction/breakdown -health or old age care | 2337 13 13 14 | 448 13 0 | 292 | 175 0 0 | 326 0 2 0 | 402 | 296 0 1 | 146 0 1 | 8000 | 0000 | 32 0 0 1 | 37 | 24 | 18 | |
| ted 17 0 0 2 3 4 4 1 0 0 1 0 0 1 0 0 1 0 0 0 0 0 0 0 0 | Legal problems | T | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | Other and not stated other n.e.c.* -unknown or not stated | 17 | 00 | 00 | 0 2 | 0 33 | 40 | 40 | 10 | 00 | 00 | но | 00 | 00 | 00 | |

*Not elsewhere classified.

Table 6.7: Male In-migration by Cause of Movement and Age, 1993

| 4 | | | | | | | A | Age (years) | (8) | | | | | | |
|--|------------------------|--------------------|--------------------|--------------------|-------------------------|--------------------|---------------|---------------------|------------------------|-------|-------------------|-------|----------|-------|-------|
| cduse of movement | Total | \$ | 6-9 | 10-14 | 15-19 | 20-24 | 25-29 | 30-34 | 35-39 | 40-44 | 45-49 | 50-54 | 55-59 | 60-64 | +59 |
| All migrants | 2416 | 436 | 277 | 173 | 164 | 210 | 275 | 335 | 216 | 107 | 9/ | 26 | 37 | 24 | 30 |
| Work/Economic/Educational -acquired/seeking job -job completion/retirement -to acquire education -educ. completed/interrupt -student lodging | 408 385 132 0 | 000000 | 201200 | 2227 | 114 0 0 0 0 | 47 19 0 0 | 72 75 0 | 102 92 4 0 | 74 0 0 0 0 | 0000 | 32 0 0 1 | 119 | m 570000 | 70000 | 00000 |
| Housing/Environmental -acquired/seeking new land/house -river erosion | 160 | 00 | 00 | 00 | 0.22 | 15 | 30 | 33 | 26 | 60 | 60 | 00 | 11 0 | 7 | 9 0 |
| Marriage/Familial -marriage -separation/divorce/widow | 7 | 00 | 00 | 00 | 00 | 10 | 00 | 0 П | 1 | 10 | 00 | 00 | 00 | 00 | 00 |
| -move with of John spouse/parents -adoption -family friction/breakdown -health or old age care | 1252 9 10 15 | 427 6 0 0 | 253 1 0 0 | 121 1 0 0 | 83000 | 78 0 0 2 | 82 0 1 | 040 | 130 | 23 | 0000 | 5010 | 8000 | HH02 | 8009 |
| Legal problems | 6 | 0 | 0 | 0 | 0 | 1 | 2 | Н | 2 | 3 | 0 | 0 | 0 | 0 | 0 |
| Other and not stated -other n.e.c,* -unknown or not stated | 17 0 | но | 00 | 10 | 0 0 | 0 3 | 0 3 | 10 | 0 3 | 00 | 00 | 10 | 00 | 10 | 00 |

*Not elsewhere classified.

Table 6.8: Female In-migration by Cause of Movement and Age, 1993


| Cause of movement | | | | | | | A | Age (years) | rs) | | | | | | |
|--|----------------------------|---------------------|-------|---------------|--------------------|------------------------|--------------------|--------------------|--------|--------------|--------|--------------|--------|--------------|----------|
| ממסכ כן וווסגפוופור | Total | 2 | 6-9 | 10-14 | 15-19 | 20-24 | 25-29 | 30-34 | 35-39 | 40-44 | 45-49 | 50-54 | 55-59 | 60-64 | 65+ |
| All migrants | 2876 | 399 | 268 | 221 | 705 | 570 | 322 | 136 | 72 | 39 | 27 | 53 | 27 | 21 | 40 |
| Work/Economic/Educational -acquired/seeking job -job completion/retirement -to acquire education -educ. completed/interrupt -student lodging | 33 74 74 00 00 | 00000 | 40000 | 328 | 133 | 10 8 0 0 0 | 000329 | 410000 | V80010 | 00000 | 000000 | 040000 | 000000 | 100000 | 000000 |
| Housing/Environmental -acquired/seeking new land/housing -river erosion | 33 | 00 | 0.0 | 00 | 0.0 | 40 | . 41 | 40 | 0.0 | 0 0 | 22 | 10 | 40 | 1 | 0 1 |
| Marriage/Familial -marriage -separation/divorce/widow | 673 | 00 | 00 | 41 | 415 | 204 | 26 | 117 | V 89 | | 0 0 5 | 10 | 0 0 | 00 | 10 |
| spouse will of Join spouse/parents -adoption -family friction/breakdown -health or old age care | 1797 16 15 11 | 384 12 0 1 | 239 | 172 1 0 | 215 0 1 0 | 279 1 5 | 221 0 6 2 | 105 0 1 3 | 0000 | 29 0 1 | 27000 | 25 0 0 | 19 | 16 0 1 | 32 0 0 3 |
| Legal problems | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other and not stated .other n.e.c.* .unknown or not stated | 19 | 10 | 00 | e 0 | 40 | 90 | 0 | 00 | 0 | 00 | 00 | 00 | 0 5 | 00 | 0 |
| | | | | | | | | | | | | | | | |

*Not elsewhere classified.

Table 6.9: In- and Out-migration by Sex and Month, 1993

| A | In-m | igration | | Out- | migratio | n |
|----------------|------------|----------|---------|------------|----------|---------|
| Age (years) | Both sexes | Males | Females | Both sexes | Males | Females |
| January | 1058 | 559 | 499 | 852 | 463 | 389 |
| February | 451 | 190 | 261 | 449 | 199 | 250 |
| March | 327 | 133 | 194 | 293 | 148 | 145 |
| April | 221 | 95 | 126 | 206 | 113 | 93 |
| May | 173 | 80 | 93 | 149 | 70 | 79 |
| June | 785 | 362 | 423 | 1486 | 718 | 768 |
| July | 486 | 217 | 269 | 924 | 448 | 476 |
| August | 507 | 218 | 289 | 857 | 414 | 443 |
| September | 396 | 183 | 213 | 634 | 302 | 332 |
| October | 392 | 170 | 222 | 780 | 385 | 395 |
| November | 279 | 121 | 158 | 516 | 288 | 228 |
| December | 217 | 88 | 129 | 346 | 159 | 187 |
| All months | 5292 | 2416 | 2876 | 7492 | 3707 | 3785 |

Figure 6.2: Number of In- and Out-migrants by Sex and Month, 1993






Table 6.10: Male Migration by Destination or Origin, 1993

| D Li Li | . / | | | 0ι | ıt-Mig | ation | | | | | In-Mi | gration | | |
|-------------|---|---|---------------------------------------|--------------------------|-------------------------------|------------------------------|--|--|---|-------------------------------------|--|-----------------------------------|--|--|
| Destination | | 1 - | | 1 | Age (ye | ears) | | | | | Age | (years) | | |
| | rural/ urban | District | 0-14 | 15-24 | 25-34 | 35-44 | 45+ | Total | 0-14 | 15-24 | 25-34 | 35-44 | 45+ | Total |
| All migrant | s | | 1020 | 1139 | 989 | 355 | 204 | 3707 | 886 | 374 | 610 | 323 | 223 | 2416 |
| Rajshahi | rural urban | all all | 0 14 | 0 5 | 0 | 0 | 0 1 | 0 27 | 3 | 0 6 | 4 5 | | 0 5 | 8 29 |
| Khulna | rural urban | all all | 6 23 | 6 32 | 3 10 | 0 | 0 | 15 67 | 6 27 | 6 5 | 4 11 | | 2 14 | 18 65 |
| Dhaka | rural rural rurai urban urban urban urban urban urban | Dhaka Narayangan Narsingdi rest Dhaka Narayangan Narsingdi Gazipur rest | 0 18 331 | 9 | 1 7 237 52 3 2 | 0 2 70 16 2 0 | 1 0 0 2 47 13 3 3 | 17 15 1 42 1098 215 28 26 33 | 13 13 9 11 190 53 2 12 25 | 0 1 5 103 35 1 6 | 7 4 7 5 186 52 5 11 | 3 4 75 18 5 3 | 2 4 0 2 46 20 4 2 10 | 33 27 20 27 600 178 17 34 |
| Chittagong | rural rural rural urban urban urban urban urban urban | Comilla Chandpur rest Sylhet Comilla Chandpur Chittgong rest | 51 256 8 45 17 8 52 | 97 3 40 13 7 | 82 3 11 7 8 43 | 67 2 5 4 4 8 | 9 53 1 5 3 3 5 2 | 104 555 17 106 44 30 171 32 | 55 308 6 29 14 9 60 | 93 0 16 11 2 2 28 | 14 | 79 0 6 10 3 7 1 | 3 57 1 10 3 3 15 2 | 29 156 |
| India | | | 48 | 37 | 27 | 15 | 19 | 146 | 15 | 5 13 | 1 | 1 3 | 3 | 45 |
| Other A | sia | | 0 | 143 | 20: | 52 | 12 | 408 | (| 7 | 36 | 5 19 | 6 | 68 |
| Middle- | East | | 1 | 132 | 2 250 | 89 | 13 | 485 | (| 0 4 | 4 | 4 24 | 8 | 80 |
| Other | | | (|) (|) (| 1 | 0 | 1 | (| 0 0 | | 0 0 | 0 | 0 |
| Unknowr | î | | 6 | 5 6 | 3 ! | 5 2 | 3 | 24 | | 0 | 0 | 0 0 | 1 | . 1 |

Table 6.11: Female Migration by Destination or Origin, 1993

| D | . I and and | | | Ou | ut-Mig | ration | | | | | In-Mig | gration | | |
|-------------|--|---|--|-----------------------------------|---|--|--|--|--|---------------------------------|---------------------------------|------------------------------|--|--|
| Destination | - | n – | | 1 | Age (ye | ears) | | | | | Age | (years) | | |
| Division | rural/ urban | District | 0-14 | 15-24 | 25-34 | 35-44 | 45+ | Total | 0-14 | 15-24 | 25-34 | 35-44 | 45+ | Total |
| All migrant | s | | 1060 | 1782 | 631 | 120 | 192 | 3785 | 888 | 1275 | 458 | 110 | 145 | 2876 |
| Rajshahi | rural urban | all all | 0 6 | 2 8 | 0 | 0 | 1 | 3 22 | 2 11 | 3 | 1 5 | 0 | 0 | 6 22 |
| Khulna | rural urban | all | 2 14 | 8 17 | 2 11 | 1 | 3 2 | 16 44 | 10 20 | 5 13 | 3 16 | 2 | 0 3 | 20 55 |
| Dhaka | rural rural rural rural urban urban urban urban urban urban | Dhaka Narayangan; Narsingdi rest Dhaka Narayangan; Narsingdi Gazipur rest | 23 353 | 10 19 | 5 5 0 9 202 46 4 4 | 0 0 0 1 36 6 1 1 3 | 1 0 1 66 17 2 2 | 23 26 3 65 1160 235 25 39 51 | 16 11 14 22 208 59 3 10 | 11 4 22 216 50 4 | 5 8 119 30 1 8 | 8 1 0 | 2 0 0 3 36 8 2 0 5 | 40 29 23 56 604 155 11 27 44 |
| Chittagong | rural rural rural urban urban urban urban urban urban | Comilla Chandpur rest Sylhet Comilla Chandpur Chittgong rest | 52 283 10 31 22 8 49 11 | 665 12 35 15 8 119 | 7 9 31 | 37 0 5 5 2 5 | 8 37 1 7 3 3 11 2 | 229 1203 27 94 52 30 215 40 | 42 293 6 27 21 7 59 | 671 21 19 12 6 | 145 5 11 10 5 30 | 35 0 13 3 0 5 | 8 53 0 8 2 1 5 | 188 1197 32 78 48 19 150 37 |
| India | | | 58 | 29 | 28 | 10 | 22 | 147 | 12 | 7 | 5 | 1 | 7 | 32 |
| Other Asia | | | 0 | 4 | 2 | 1 | 0 | 7 | C |) 0 | 0 | 1 | 0 | 1 |
| Middle-Eas | t | | 6 | 8 | 6 | 1 | 0 | 21 | 0 |) 1 | . 0 | 0 | 0 | 1 |
| Other | | | 0 | 0 | (| 0 | 0 | 0 | 1 | |) 0 | 0 | 0 | 1 |
| Unknown | | | 3 | 3 | . 1 | . 0 | 1 | 8 | (|) (|) (| 0 | 0 | (|

CHAPTER 7

Special Supplement

FERTILITY TRENDS IN MATLAB BY BIRTH ORDER, AGE AT BIRTH AND BIRTH INTERVAL, 1980-1993

The rapid decline in fertility in Matlab during the 1980's and early 1990's is perhaps the most important and striking feature to emerge from the Demographic Surveillance System. Between 1980 and 1993 total fertility fell from 5.1 to 2.9 births per woman in the MCH-FP area and from 6.7 to 3.9 in the Comparison area. The bulk of the "fertility transition" was completed in both areas in a remarkably short space of time. In this issue of the DSS report we take a closer look at this transition, and we analyse its components in terms of birth order, mothers' age at birth, and the length of the intervals between births.

Birth Order Fertility Rates and Parity Progression Ratios

If the births occurring in each year are tabulated, not only by mother's age, but also by birth order (i.e. whether it is her first, second, third...etc.) and the numbers in each age-group-birth-order cell are divided by the total number of women in the age group, age-order-specific fertility rates are obtained. The rates for 1993 are shown in Table 4.7.

Now each age-order fertility rate represents the proportion of women in the age group who had their nth birth during the year in question. Thus if the rates are summed across the child-bearing age groups (and multiplied by 5 if 5-year age groups are used), the total order rates so obtained will represent the proportion of women who will have at least n births by the age of 50 if they go through life having children at the current rates. Table S.1 shows the total order rates for both the Matlab areas from 1980 to 1993.

Despite substantial year-by-year fluctuations, the total first-birth rates show no consistent change over the period. But as we move up the birth orders, bigger and bigger declines are apparent, so that the 1993 rates for birth orders 5 and over were less than half of the 1980 rates in both areas. The contrast between the two areas is also striking. The fertility transition began earlier in the MCH-FP area: in the mid-1970's total fertility in the two areas was virtually identical (LeGrand and Phillips 1996), but by 1980 a substantial differential had already opened up. Thus the relative difference between the MCH-FP area and Comparison area rates for 1980 tend to increase with birth order. All these trends and differentials thus reflect the reduction in the proportions of women having large numbers of children which is the basic mechanism of the fail in fertility.

However, order-specific fertility rates are unsatisfactory indices largely because the denominators (all women in the age group) are not those who are truly "exposed to risk". Fourth-order births can only by borne by women who have already had three. Thus "true" birth-order rates should be obtained by dividing the number of births of order n by the number of women who, at the beginning of the year, had had n-1 (twins apart). With such rates, it is possible to construct "parity progression ratios" (PPR's) - the proportion of women with n children who ultimately go on to have n+1. These ratios are commonly regarded as the most powerful and sensitive measures of fertility change. As with other measures of fertility and mortality, they may be calculated on either a cohort or a period basis; we have used the period approach which is better adapted for measuring changes in the recent past.

Table S.2 shows period PPR's, calculated by Feeney's indirect method (Feeney 1985, 1986, 1992). Further details of this method are described in the Technical Note at the end of the article. For the graphic presentation of the main features of these results we have averaged successive pairs of parity progressions (1 to 2 and 2 to 3; 3 to 4 and 4 to 5,...etc.), and we have shown the values of the PPR's in each year relative to 1980. The results are shown in Figure S.1. While there was virtually no decline in either area in the PPR's at 1-2/2-3, the falls increase systematically with increasing birth order. In the Comparison area the pace of decline was slow up to 1990 and then steepened. Thus the contrast between the two areas is clearly apparent from the two graphs.

Age at Birth and Birth Order

From tabulations of births by mother's age and birth order, we have calculated the average age of mothers at the birth of their children by birth order. The results are shown in Table S.3.

There was a general rise in age at birth, particularly of women having their first, second, third and fourth births in both areas; for those having fifth and higher order births, the rise in average age was less pronounced in the MCH-FP area, and in the Comparison area there was no systematic change at all. These features are illustrated in Figure S.2., where we have again shown the changes relative to 1980.

Birth Intervals and Birth Order

From tabulations of births by birth order and the interval since the preceding birth we have calculated the average interval between births for each birth order. The tabulations could not be made for years prior to 1983, as the information on the date of the preceding birth had to be extracted from the data base, which was constructed in 1982. We have also omitted the figures for 1983 which appeared suspect with large numbers of unknown values. The results are shown in Table S.4, and are illustrated in Figure S.3.

There was a general lengthening of birth intervals in Matlab during the period under review. They increased by 9.5 months between 1984 and 1993 in the MCH-FP area, and by 3.6 months in the Comparison area. These changes were more pronounced at lower birth orders - particularly the intervals between second and third, and between third and fourth, births. At higher birth orders an upward trend is still discernible, but it is less steep.

The intervals between first and second births were consistently shorter than those between higher order births. Otherwise there appears to be no association between the length of the interval and birth order, except perhaps at the highest open-ended category, which tended to be shorter.

The birth intervals in the Comparison area were consistently shorter than those in the MCH-FP area, and as the increase in the intervals over time were also less, the differences between the two areas became increasingly accentuated.

Discussion

The dramatic falls in the proportions of women having large numbers of children revealed by the parity progression ratios were the direct result of the rapid uptake of family planning by the people of Matlab. In the MCH-FP area contraceptive prevalence had been about 10% before the start of the project, and increased to 28% by 1978, 46% by 1984, and 63% by 1993. Annual data for the Comparison area are not available, but surveys conducted in 1984 and 1992 gave figures of 16% and 30% respectively (LeGrand and Phillips 1996; ICDDR,B 1995).

The rise in age at first birth was doubtless the result, in part at least, of the rise in age at marriage described in Chapter 5. The singulate mean ages at marriage for females derived from the distributions of the population by age and marital status from the Matlab censuses of 1974, 1982 and 1993 were 17.0, 18.6 and 20.3 respectively. There may also have been some increase in the interval between marriage and first birth, but changes in this respect were small and erratic. The corresponding rises in ages at second, third and fourth births can thus be seen as knock-on effect. However the lengthening of the birth intervals strongly suggests that contraception was being used for spacing as well as stopping. It cannot be attributed to longer breast-feeding and post-partum amenorrhoea as these were in fact shortening during the period under review (Salway, Roy and Koenig 1992).

In this article we have been able to do no more than show the bare bones of the fertility transition in Matlab, but we hope that these will stimulate further in-depth studies and research.

Technical Note

Feeney's method for the indirect estimation of parity progression ratios requires information simply on the annual numbers of births by birth order. Such data are available for the two areas of Matlab for the period covered by this study from 1980 to 1993, but information on birth intervals, which would be required for direct estimates of PPR's, cannot be calculated for years prior to 1983, as the data base was only constructed in 1982. Feeney's indirect method overcomes this deficiency by using models or standard sets of progression rates by duration in parity. We have constructed such models from the data on intervals since the preceding birth for the years 1985-88, and we have used separate models for the intervals between first and second births and those between second and higher order births.

Feeney's method also requires data on the open birth interval distribution for the initial year of the study. As such data are rarely available, he suggests that a "stationary population" based on the standard progression rates by duration may be used. However he notes that this procedure may lead to biases in the estimated PPR's for the first few years of the study. We believe that in the case of the Matlab data such biases may have tended to under-estimate the PPR's in the early years. In other words the decline in the PPR's may in reality have been even steeper than those shown by our series.

It may be noted that we have not attempted to show the progression ratios from marriage to first birth. This is because the numbers of marriages recorded as taking place in the study area have been inflated by the inclusion of the marriages of brides who had been living in the area up to the time of their marriage but who married grooms from outside the area. As their subsequent childbearing occurred outside the area the ratios of first births to marriages were biased downwards.

References

- Feeney, G., 1985, "Parity Progression Projection", International Population Conference Florence 1985, Vol.4, pp.125-136.
- Feeney, G.,1986, "Period Parity Progression Measures of Fertility in Japan", NUPRI Research Paper Series No. 35.
- Feeney, G., 1992, "Indirect Estimation of Period Parity Progression Ratios from Time Series of Births distributed by Order", Unpublished Note.
- ICDDR,B, 1995, Demographic Surveillance System Early Indicators, Matlab 1994.
- LeGrand, T.K., and Phillips, J.F., 1996, "The Effect of Fertility Reduction on Infant and Child Mortality: Evidence from Matlab in Rural Bangladesh", Population Studies Vol.50 No.1 pp.51-68
- Salway, S., Roy N.Ch., and Koenig M.A., "Levels and Trends in Postpartum Amenorrhoea, Breast-Feeding and Birth Intervals in Matlab, Bangladesh 1978-1989", ICDDR,B Working Paper No.69.

Table S.1: Total Order-specific Fertility Rates by Area, 1980-1993

| Year | Total | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10+ |
|------|------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| (a) | MCH-FP are | ea | | | | | | | | | |
| 1980 | 5.0995 | 0.7687 | 0.7820 | 0.6982 | 0.7020 | 0.6069 | 0.5584 | 0.3652 | 0.2615 | 0.1800 | 0.1766 |
| 1981 | 4.8118 | 0.7715 | 0.7624 | 0.6548 | 0.6514 | 0.6092 | 0.4424 | 0.3439 | 0.2474 | 0.1688 | 0.1587 |
| 1982 | 4.9602 | 0.8407 | 0.7454 | 0.7221 | 0.7003 | 0.6340 | 0.4462 | 0.3739 | 0.2212 | 0.1389 | 0.1356 |
| 1983 | 4.4982 | 0.7411 | 0.7521 | 0.6491 | 0.5616 | 0.5194 | 0.4118 | 0.3525 | 0.2354 | 0.1355 | 0.136 |
| 1984 | 3.9292 | 0.6711 | 0.6643 | 0.5690 | 0.5254 | 0.4441 | 0.3399 | 0.2556 | 0.1886 | 0.1294 | 0.1399 |
| 1985 | 4.4496 | 0.7213 | 0.7490 | 0.6634 | 0.5891 | 0.5530 | 0.4057 | 0.3063 | 0.2099 | 0.1335 | 0.117 |
| 1986 | 4.2407 | 0.7971 | 0.6795 | 0.6765 | 0.5325 | 0.4473 | 0.3832 | 0.2984 | 0.2086 | 0.1146 | 0.103 |
| 1987 | 4.1543 | 0.7596 | 0.6710 | 0.7142 | 0.5609 | 0.4607 | 0.3552 | 0.2453 | 0.1496 | 0.1199 | 0.118 |
| 1988 | 3.7733 | 0.7560 | 0.7029 | 0.6171 | 0.5140 | 0.3706 | 0.2920 | 0.2091 | 0.1419 | 0.0913 | 0.077 |
| 1989 | 3.4198 | 0.6417 | 0.6362 | 0.6111 | 0.4723 | 0.3426 | 0.2564 | 0.2099 | 0.1161 | 0.0645 | 0.069 |
| 1990 | 3.3662 | 0.7842 | 0.6385 | 0.5473 | 0.4407 | 0.3061 | 0.2142 | 0.1679 | 0.1249 | 0.0661 | 0.076 |
| 1991 | 3.0048 | 0.7489 | 0.5620 | 0.4922 | 0.4409 | 0.2881 | 0.1820 | 0.1212 | 0.0741 | 0.0488 | 0.046 |
| 1992 | 2.9569 | 0.7745 | 0.6286 | 0.4953 | 0.4087 | 0.2572 | 0.1542 | 0.1051 | 0.0500 | 0.0366 | 0.046 |
| 1993 | 2.8910 | 0.7726 | 0.5934 | 0.4971 | 0.3927 | 0.2633 | 0.1738 | 0.0908 | 0.0433 | 0.0442 | 0.019 |
| (b) | Comparison | n area | | | | | | | | | |
| 1980 | 6.6741 | 0.8428 | 0.7780 | 0.8780 | 0.9048 | 0.9339 | 0.7105 | 0.5794 | 0.4536 | 0.2820 | 0.310 |
| 1981 | 6.2777 | 0.8513 | 0.8531 | 0.8040 | 0.8114 | 0.7720 | 0.6642 | 0.5644 | 0.4035 | 0.2630 | 0.290 |
| 1982 | 6.3206 | 0.8439 | 0.7925 | 0.8157 | 0.8689 | 0.7751 | 0.7226 | 0.5278 | 0.4147 | 0.2763 | 0.282 |
| 1983 | 5.9964 | 0.8335 | 0.8187 | 0.7144 | 0.7372 | 0.7561 | 0.6890 | 0.5649 | 0.3649 | 0.2349 | 0.280 |
| 1984 | 5.0667 | 0.7360 | 0.7166 | 0.6590 | 0.6329 | 0.5864 | 0.5412 | 0.4459 | 0.3235 | 0.2064 | 0.216 |
| 1985 | 5.8991 | 0.8209 | 0.8046 | 0.7926 | 0.7356 | 0.7065 | 0.6465 | 0.4857 | 0.3570 | 0.2760 | 0.271 |
| 1986 | 5.4449 | 0.7452 | 0.7965 | 0.7877 | 0.6738 | 0.6247 | 0.5455 | 0.4794 | 0.2832 | 0.2479 | 0.260 |
| 1987 | 5.3062 | 0.8160 | 0.6989 | 0.7021 | 0.7100 | 0.6303 | 0.5529 | 0.4107 | 0.3487 | 0.2011 | 0.233 |
| 1988 | 5.4001 | 0.8673 | 0.7336 | 0.7444 | 0.7457 | 0.6000 | 0.5655 | 0.3945 | 0.3168 | 0.1790 | 0.253 |
| 1989 | 4.8904 | 0.7398 | 0.6909 | 0.7624 | 0.6210 | 0.5283 | 0.4978 | 0.3721 | 0.2851 | 0.1745 | 0.218 |
| 1990 | 5.0327 | 0.7638 | 0.8228 | 0.6313 | 0.6840 | 0.5915 | 0.4592 | 0.3966 | 0.2817 | 0.1928 | 0.208 |
| 1991 | 4.3265 | 0.8742 | 0.6792 | 0.5979 | 0.5855 | 0.4607 | 0.3370 | 0.3300 | 0.1976 | 0.1368 | 0.127 |
| 1992 | 4.0343 | 0.8488 | 0.6958 | 0.6342 | 0.4884 | 0.4082 | 0.3087 | 0.2143 | 0.1710 | 0.1243 | 0.140 |
| | | | | | | | | | | 0.1087 | 0.102 |

Table S.2: Period Parity Progression Ratios for Matlab by Area, 1980-1993

| Year | 1-2 | 2-3 | 3-4 | 4-5 | 5-6 | 6-7 | 7-8 | 8-9 |
|--------|------------|--------|-------|-------|-------|-------|-------|-------|
| (a) MC | H-FP area | 1 | | | | | | |
| 1980 | 0.814 | 0.732 | 0.852 | 0.827 | 0.835 | 0.699 | 0.715 | 0.669 |
| 1981 | 0.802 | 0.725 | 0.850 | 0.833 | 0.771 | 0.667 | 0.698 | 0.659 |
| 1982 | 0.814 | 0.776 | 0.879 | 0.859 | 0.768 | 0.702 | 0.656 | 0.582 |
| 1983 | 0.834 | 0.769 | 0.846 | 0.810 | 0.736 | 0.696 | 0.673 | 0.565 |
| 1984 | 0.812 | 0.738 | 0.806 | 0.747 | 0.655 | 0.596 | 0.590 | 0.556 |
| 1985 | 0.859 | 0.787 | 0.817 | 0.806 | 0.694 | 0.659 | 0.605 | 0.562 |
| 1986 | 0.852 | 0.817 | 0.804 | 0.755 | 0.686 | 0.676 | 0.612 | 0.511 |
| 1987 | 0.862 | 0.854 | 0.813 | 0.778 | 0.676 | 0.620 | 0.510 | 0.546 |
| 1988 | 0.872 | 0.812 | 0.788 | 0.703 | 0.632 | 0.573 | 0.489 | 0.458 |
| 1989 | 0.836 | 0.818 | 0.739 | 0.664 | 0.591 | 0.583 | 0.454 | 0.371 |
| 1990 | 0.836 | 0.782 | 0.696 | 0.605 | 0.535 | 0.529 | 0.487 | 0.391 |
| 1991 | 0.779 | 0.739 | 0.691 | 0.592 | 0.485 | 0.427 | 0.357 | 0.343 |
| 1992 | 0.803 | 0.734 | 0.671 | 0.550 | 0.438 | 0.407 | 0.252 | 0.302 |
| 1993 | 0.774 | 0.738 | 0.657 | 0.559 | 0.513 | 0.399 | 0.259 | 0.374 |
| (b) C | comparison | ı area | | | | | | |
| 1980 | 0.806 | 0.870 | 0.874 | 0.870 | 0.768 | 0.781 | 0.754 | 0.617 |
| 1981 | 0.831 | 0.872 | 0.857 | 0.840 | 0.750 | 0.780 | 0.720 | 0.603 |
| 1982 | 0.832 | 0.899 | 0.898 | 0.849 | 0.789 | 0.769 | 0.733 | 0.619 |
| 1983 | 0.859 | 0.881 | 0.882 | 0.852 | 0.783 | 0.784 | 0.671 | 0.559 |
| 1984 | 0.834 | 0.846 | 0.829 | 0.776 | 0.728 | 0.692 | 0.629 | 0.511 |
| 1985 | 0.868 | 0.877 | 0.863 | 0.817 | 0.781 | 0.703 | 0.641 | 0.633 |
| 1986 | 0.879 | 0.890 | 0.849 | 0.793 | 0.741 | 0.713 | 0.566 | 0.627 |
| 1987 | 0.859 | 0.868 | 0.866 | 0.824 | 0.780 | 0.672 | 0.667 | 0.579 |
| 1988 | 0.870 | 0.881 | 0.883 | 0.811 | 0.797 | 0.675 | 0.647 | 0.548 |
| 1989 | 0.849 | 0.897 | 0.829 | 0.767 | 0.758 | 0.660 | 0.631 | 0.537 |
| 1990 | 0.889 | 0.857 | 0.856 | 0.787 | 0.741 | 0.684 | 0.642 | 0.566 |
| 1991 | 0.838 | 0.817 | 0.804 | 0.710 | 0.630 | 0.616 | 0.514 | 0.454 |
| 1992 | 0.837 | 0.814 | 0.739 | 0.644 | 0.583 | 0.497 | 0.490 | 0.449 |
| 1992 | | | | | | | | |


Table S.3: Mean Age of Mothers at the Birth of their Children by Birth Order, 1980-1993

| | | | | | Birth or | der | | | | |
|------|------------|-------|-------|-------|----------|-----------|-------|-------|-------|------|
| /ear | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10+ |
| (a) | MCH-FP are | a | | | | | | | | |
| 1980 | 19.01 | 21.79 | 24.39 | 27.58 | 30.01 | 33.00 | 34.83 | 36.58 | 37.50 | 39.8 |
| 1981 | 19.54 | 22.28 | 24.67 | 27.21 | 30.05 | 32.34 | 34.49 | 36.46 | 38.22 | 38.4 |
| 1982 | 19.63 | 22.24 | 24.80 | 27.45 | 30.10 | 33.07 | 34.63 | 36.44 | 37.72 | 40.5 |
| 1983 | 19.33 | 22.22 | 24.91 | 26.81 | 29.69 | 32.18 | 34.47 | 35.82 | 38.71 | 38.9 |
| 1984 | 19.90 | 22.35 | 24.72 | 27.33 | 30.19 | 32.31 | 34.73 | 36.35 | 37.58 | 39.4 |
| 1985 | 20.14 | 22.55 | 24.84 | 27.87 | 30.18 | 32.90 | 34.92 | 36.33 | 37.89 | 40.2 |
| 1986 | 20.40 | 22.83 | 24.97 | 27.53 | 30.06 | 32.84 | 34.52 | 37.10 | 38.70 | 40.3 |
| 1987 | 20.29 | 23.09 | 25.11 | 28.02 | 29.97 | 32.85 | 35.40 | 36.42 | 38.28 | 39.1 |
| 1988 | 21.01 | 23.25 | 25.87 | 27.78 | 30.48 | 32.30 | 35.22 | 37.93 | 39.43 | 40.4 |
| 1989 | 20.91 | 23.70 | 25.95 | 28.22 | 30.67 | 32.44 | 35.30 | 36.13 | 38.75 | 40.8 |
| 1990 | 21.07 | 23.75 | 26.50 | 28.78 | 31.18 | 32.50 | 34.95 | 37.12 | 39.84 | 41.1 |
| 1991 | 21.02 | 24.29 | 26.77 | 29.16 | 30.94 | 33.30 | 35.46 | 36.40 | 38.50 | 41.6 |
| 1992 | | 24.25 | 26.97 | 29.23 | 31.06 | 33.99 | 35.31 | 38.27 | 37.00 | 40.7 |
| 1993 | | 24.62 | 27.29 | 29.85 | 31.72 | 33.69 | 35.26 | 37.50 | 39.24 | 37.9 |
| (b) | Comparisor | area | | | | | | | | |
| 1980 | 19.33 | 21.54 | 24.29 | 26.88 | 30.17 | 32.32 | 34.66 | 36.33 | 38.18 | 39.5 |
| 1981 | | 22.09 | 24.38 | 27.11 | 29.46 | 31.87 | 35.03 | 35.78 | 37.58 | 38.8 |
| 1982 | | 22.32 | 24.30 | 26.71 | 29.79 | 32.31 | 34.60 | 36.21 | 37.70 | 39.4 |
| 1983 | | 22.09 | 24.13 | 26.39 | 29.47 | 32.00 | 34.20 | 35.78 | 37.59 | 40.0 |
| 1984 | | 22.26 | 24.53 | 26.83 | 29.05 | 31.50 | 34.29 | 35.77 | 37.82 | 39. |
| 1985 | | 22.14 | 24.63 | 26.60 | 29.26 | 31.73 | 33.85 | 36.69 | 37.95 | 39.9 |
| 1986 | | 22.58 | 24.45 | 26.77 | 29.49 | 31.89 | 33.82 | 36.14 | 37.73 | 39. |
| 1987 | | 22.69 | 24.65 | 27.06 | 29.13 | 31.26 | 34.01 | 35.98 | 37.33 | 40. |
| 1988 | | 22.90 | 24.85 | 27.07 | 29.35 | 31.73 | 33.32 | 35.83 | 38.00 | 39. |
| 1989 | | 23.04 | 25.39 | 27.55 | 29.42 | 32.08 | 33.45 | 35.47 | 36.99 | 40. |
| 1990 | | 23.38 | 25.28 | 27.47 | 29.62 | 31.70 | 33.73 | 35.89 | 38.13 | 39. |
| 1991 | | 23.72 | 25.92 | 27.46 | 29.39 | 31.75 | 34.72 | 36.62 | 38.31 | 39. |
| | | 23.70 | 26.00 | 27.99 | 30.38 | 32.18 | 33.83 | 35.13 | 37.75 | 39. |
| 1992 | 21.08 | 23.70 | 20.00 | 61.33 | 30.00 | 0 = 1 = 0 | | | | |

Table S.4. Mean Birth Intervals in Months by Birth Order, 1984-1993

| | | | | | В | irth orde | r | | | |
|------|------------|--------|-------|-------|-------|-----------|-------|-------|-------|--------|
| Year | Total | 1-2 | 2-3 | 3-4 | 4-5 | 5-6 | 6-7 | 7-8 | 8-9 | 9+-10+ |
| (a) | MCH-FP are | a | | | | | | | | |
| 1984 | 39.38 | 37.02 | 37.77 | 39.35 | 43.00 | 40.39 | 40.54 | 45.87 | 41.31 | 43.94 |
| 1985 | 39.08 | 36.30 | 38.92 | 41.33 | 40.19 | 41.17 | 39.37 | 42.17 | 36.22 | 42.57 |
| 1986 | 40.46 | 37.68 | 39.65 | 41.14 | 42.30 | 44.40 | 44.86 | 42.65 | 39.78 | 35.74 |
| 1987 | 41.15 | 38.41 | 40.65 | 42.82 | 42.45 | 42.67 | 53.60 | 38.00 | 41.79 | 29.79 |
| 1988 | 40.91 | 37.85 | 41.79 | 42.19 | 43.67 | 41.26 | 41.51 | 42.00 | 41.30 | 45.94 |
| 1989 | 42.30 | 40.15 | 43.76 | 41.79 | 43.14 | 44.08 | 45.56 | 42.89 | 39.10 | 40.59 |
| 1990 | 44.34 | 41.12 | 46.49 | 47.76 | 44.76 | 43.04 | 42.50 | 45.38 | 40.50 | 46.67 |
| 1991 | 45.95 | 42.74 | 47.87 | 48.84 | 47.34 | 46.58 | 42.96 | 42.15 | 39.36 | 43.30 |
| 1992 | 47.56 | 44.91 | 48.38 | 49.81 | 49.79 | 49.32 | 50.90 | 49.38 | 34.80 | 39.65 |
| 1993 | 48.89 | 44.89 | 51.92 | 52.81 | 48.42 | 48.39 | 50.57 | 45.12 | 44.35 | 41.45 |
| Mean | 42.72 | 39.90 | 43.44 | 44.46 | 44.20 | 44.06 | 45.16 | 43.08 | 39.11 | 40.53 |
| (b) | Comparison | n area | | | | | | | | |
| 1984 | 34.70 | 33.77 | 34.44 | 34.04 | 34.61 | 36.79 | 34.73 | 34.48 | 35.74 | 38.50 |
| 1985 | 34.30 | 32.92 | 34.29 | 35.10 | 36.34 | 33.60 | 33.03 | 34.49 | 36.71 | 34.05 |
| 1986 | 34.99 | 33.80 | 35.37 | 34.03 | 36.16 | 34.59 | 37.52 | 37.02 | 35.21 | 33.84 |
| 1987 | 34.18 | 32.89 | 34.39 | 34.91 | 35.05 | 35.26 | 34.29 | 32.98 | 33.87 | 32.67 |
| 1988 | 34.09 | 33.24 | 34.27 | 34.20 | 35.19 | 33.27 | 34.15 | 35.34 | 36.30 | 32.81 |
| 1989 | 35.44 | 34.06 | 36.66 | 35.92 | 34.85 | 35.11 | 35.57 | 36.71 | 34.94 | 35.74 |
| 1990 | 35.58 | 33.62 | 37.47 | 37.24 | 34.86 | 36.26 | 34.69 | 35.82 | 37.41 | 31.82 |
| 1991 | 36.56 | 35.32 | 37.08 | 36.92 | 37.19 | 36.56 | 38.22 | 37.36 | 35.81 | 32.79 |
| 1992 | 37.51 | 35.73 | 37.87 | 38.31 | 38.03 | 40.46 | 38.72 | 35.11 | 39.25 | 34.29 |
| 1993 | 38.29 | 35.39 | 41.45 | 38.64 | 38.64 | 38.46 | 40.13 | 37.66 | 39.62 | 39.83 |
| Mean | 35.33 | 34.10 | 36.36 | 35.91 | 36.25 | 36.33 | 36.39 | 36.22 | 36.83 | 34.92 |

Figure S1: Relative Change in PPR's (1980=1) by Area, 1980-93







Figure S2: Relative Change in Age at Birth by Area, 1980-93








Figure S3: Trends in Birth Interval by Order by Area, 1984-93





Appendix A Names and Codes of Villages in the DSS Area, 1993

| | | MCH-FP are | ea | | | Comparison | area | |
|--------|---|--|--|---|--|--|--|---|
| Block* | Village code | | Village code | Village name | Village code | Village name | Village code | Village name |
| A | D W V10 V11 V31 V32 | Kaladi Dhakirgaon Nabakalash | V59 V60 V61 V62 V72 | Doshpara Suvankardi Munsabdi Shilmondi Upadi | A B C F G J U | Uddamdi Charmasua Sarderkandi Sepoykandi Thatalia Char Harigope Baispur | V78 V79 V80 V90 V95 V96 V97 | Soladana Pitambordi Daribond Narinda Baluchar Rampur Dhanagoda |
| В | H V12 V13 V19 V20 V21 V22 V23 V24 | Bhangerpar Baburpara Lakshmipur Dagorpur Khadergaon Beloti | V26 V56 V82 V83 V85 V87 VB12 VB13 | Narayanpur Palipara Dhanarpar Padmapal Bhanurpara Hurmaisha Nagda Naogaon | V01 V02 V03 V04 V05 V06 V07 V08 V09 V14 | Kadamtali Nilokhi Char Nilokhi Char Pathalia Gazipur Fatepur Nayakandi Goalbhar Naburkandi Enayetnagar | V98 V99 VB1 VB2 VB3 VB4 VB5 VB6 VB7 VB8 | Santoshpur Baluakandi Taltoli Sree Rayerchar Rayerkandi Ramdaspur Thakurpara Sarkerpara Mirpur Farazikandi |
| С | K L M N O P Q V27 V28 V30 V39 | Shahpur Tatkhana Char Nayergaon Aswinpur Nayergaon Titerkandi Char Shibpur Panchghoria Khidirpur Harion Gobindapur | V40 V41 V42 V43 V44 V64 V86 V88 VB11 D100 D101 | Masunda Paton Adhara (South) Kanachak Panchdona Kawadi Adhara Datikara Mehron Barogaon Naojan | V35 V36 V37** V38 V45 V46 V47 V48 V49 V50 V51 V53 | Durgapur Ludhua Charputia Galimkha Bakchar Silinda Tulatali Gangkandi Harina Bhabanipara Bakharpur Induriakandi Chhoto Haldia | VB9** VB10 D28 D29 D30 D31 D32 D33 D34 D35 D88 D89 | Ramanathgonj South Rampur Bazarkhola Kirtonkhola Banuakandi Harina Bazarkho Khalisha Nayanagar Saidkharkandi Molla Kandi Sankibhanga Sankibangha Namapara |
| D | R S T V15 V16 V17 V18 V25 V29 V33 V34 | Nandalalpur Tatua Amuakanda Bhati Rasulpur Binandapur Hatighata Torkey Char Pathalia Shibpur(South) Shibpur(North) | V63 V67 V81) V84 | Nayakandi Balakandi Induria Baluchar Islamabad (East) Majlishpur Sonaterkandi Shanbajkandi Islamabad (Middle) | V58** V65 V66 V68 V69** V70** V71 V73 V74 V75 V76 | Mohishmari Nayachar Thatalia Sobahan Naobangha South Joypur Khamarpara Sadardia Ketundia Mukundia Chosoi | D90 D91** D92** D93 D94 | Zahirabaj North Joypur West Joypur Maizkandi Hazipur Tapaderpara Rampur Nayakandi Bara Haldia Mandertoli |

^{*}Division by block applies only to the MCH-FP area. **Lost due to river erosion.

Appendix B
Mid-year Population, Birth, and Deaths by Village, 1993

| Village code* | Popula- tion | Live births | Deaths | Birth rate | Death |
|------------------|-----------------|----------------|------------------|---------------|---------------------------|
| D | 1800 | 52 | 10 | 28.9 | 5.6 |
| W | 3651 | 75 | 20 | 20.5 | 5.5 7.1 |
| V10 | 1548 | 36 | 11 | 23.3 | 7.1 |
| V11 | 1867 | 46 | 15 | 24.6 | 8.0 |
| V31 V32 | 8527 2646 | 197 53 | 60 20 | 23.1 | 7.0 |
| V52 V59 | 1024 | 27 | 10 | 20.0 26.4 | 7.6 |
| V60 | 896 | 18 | 5 | 20.4 | 9.8 5.6 7.4 11.6 |
| V61 | 673 | 22 | 5 | 32.7 | 7.4 |
| V62 | 859 | 15 | 10 | 17.5 | 11.6 |
| V72 | 5891 | 208 | 48 | 35.3 | 8.1 |
| Block A | 29382 | 749 | 214 | 25.5 | 7.3 |
| Н | 1178 | 35 | 9 7 | 29.7 | 7.6 |
| V12 | 530 | 12 | 7 | 22.6 | 13.2 |
| V13 | 748 | 22 | 5 | 29.4 | 6.7 7.9 |
| V19 | 2918 | 57 | 23 | 19.5 | 7.9 |
| V20 V21 | 1176 486 | 28 12 | 9 | 23.8 24.7 | 7.7 |
| V21 V22 | 563 | 10 | 6 | 17.8 | 4.1 |
| V23 | 507 | 12 | 9 2 6 1 | 23.7 | 2.0 |
| V24 | 2631 | 84 | 30 | 31.9 | 2.0 11.4 |
| V26 | 2601 | 56 | 24 | 21.5 | 9.2 |
| V56 | 1467 | 34 | 4 | 23.2 | 9.2 2.7 7.6 |
| V82 | 1453 | 40 | 11 | 27.5 | 7.6 |
| V83 | 522 | 11 | 5 | 21.1 | 9.6 |
| V85 | 434 | 12 | 5 4 1 | 27.6 | 9.2 1.7 |
| V87 | 594 | 20 | | 33.7 | 1.7 |
| VB12 VB13 | 4097 4600 | 134 102 | 24 43 | 32.7 22.2 | 5.9 9.3 |
| Block B | 26505 | 681 | 208 | 25.7 | 7.8 |

(continued)

Appendix B (cont.)

| Village code | Popula- tion | Live births | Deaths | Birth rate | Death rate |
|-----------------|-----------------|----------------|--|---------------|---------------|
| K | 892 | 19 | 9 | 21.3 | 10.1 |
| Ľ | 506 | 16 | 3 2 | 31.6 | 5.9 |
| М | 173 | 4 | 2 | 23.1 | 11.6 |
| N | 2020 | 48 | 15 | 23.8 | 7.4 |
| 0 | 1442 | 36 | 7 | 25.0 | 4.9 |
| P | 1899 | 46 | 11 | 24.2 | 5.8 |
| Q | 325 | 5 | 1 | 15.4 | 3.1 |
| V27 | 866 | 18 | 7 | 20.8 | 8.1 |
| V28 | 1401 | 39 | 11 | 27.8 | 7.9 |
| V30 | 543 | 7 | 6 | 12.9 | 11.0 |
| V39 | 331 | 2 | 6 | 6.0 | 3.0 |
| V40 | 755 | 22 | 7 15 | 29.1 | 3.0 9.3 |
| V41 | 1528 | 41 | 15 | 26.8 | 9.8 |
| V42 | 719 | 17 | 3 | 23.6 | 4.2 |
| V43 | 854 | 20 | 3 9 4 | 23.4 | 10.5 |
| V44 | 605 | 9 | 4 | 14.9 | 6.6 |
| V64 | 4488 | 96 | 33 | 21.4 | 7.4 |
| V86 | 786 | 15 | 4 | 19.1 | 5.1 |
| V88 | 486 | 11 | 1 | 22.6 | 2 1 |
| VB11 | 2486 | 56 | 24 | 22.5 | 2.1 9.7 |
| D100 | 3196 | 82 | 30 | 25.7 | 9.4 |
| D101 | 1274 | 27 | 10 | 21.2 | 7.8 |
| Block C | 27575 | 636 | 213 | 23.1 | 7.7 |
| R S T | 1344 | 30 | 12 | 22.3 | 8.9 |
| 2 | 956 | 18 | 6 | 18.8 | 6.3 |
| 1 | 1534 | 39 | 17 | 25.4 | 11.1 |
| V15 | 598 | 16 | 5 | 26.8 | 8.4 |
| V16 | 769 | 10 | 5 | 13.0 | 6.5 |
| V17 | 1081 | 25 | 5 5 6 25 | 23.1 | 5.6 |
| V18 | 3718 | 111 | 25 | 29.9 | 6.7 |
| V25 | 1220 | 25 | 5 | 20.5 | 4.1 |
| V29 | 438 | 9 | 6 | 20.5 | 13.7 |
| V33 | 506 | 10 | 0 | 19.8 | 0.0 |
| V34 | 808 | 16 | 8 | 19.8 | 9.9 |
| V52 | 236 | 6 | 2 | 25.4 | 8.5 |
| V54 | 637 | 16 | 7 | 25.1 | 11.0 |
| V55 | 536 | 13 | 7 | 24.3 | 13.1 |
| V57 | 1068 | 26 | 5 6 0 8 2 7 7 9 21 | 24.3 | 8.4 |
| V63 | 2046 | 45 | 21 | 22.0 | 10.3 |
| V67 | 606 | 14 | 6 | 23.1 | 9.9 |
| V81 | 642 | 19 | 4 | 29.6 | 6.2 |
| V84 | 2181 | 68 | 12 | 31.2 | 5.5 |
| V89 | 1384 | 33 | 14 | 23.8 | 10.1 |
| Block D | 22308 | 549 | 177 | 24.6 | 7.9 |
| MCH-FP Area | 105770 | 2615 | 812 | 24.7 | 7.7 |

(continued)

Appendix B (cont.)

| Village | Popula- | Live | Deaths | Birth | Death |
|---|--|---|--|---|---|
| code | tion | births | | rate | rate |
| A B C F G J V001 V002 V004 V005 V006 V007 V008 V009 V14 V356 V37 V38 V49 V40 V47 V48 V49 V50 V50 V66 V68 V69 V70 V71 V73 V74 V75 V76 V77 V76 V77 V77 V77 V78 V79 V79 V79 V79 V79 V79 V79 V79 V79 V79 | 2862 2072 3779 1421 2520 543 8005 626 527 659 303 3373 2364 409 1281 1144 834 3610 5102 0 1693 1090 403 1814 605 1326 134 931 3230 0 740 804 890 0 457 784 1387 389 1670 247 354 1135 1169 1712 622 438 174 677 | 84 66 116 41 76 11 25 19 17 12 60 60 60 60 60 60 60 60 60 60 60 60 60 | 17 23 36 15 24 47 77 35 92 38 28 22 12 12 42 90 18 85 18 31 59 65 29 12 13 16 59 16 59 16 59 17 26 59 18 59 19 59 19 59 19 59 19 59 19 59 19 59 19 59 19 59 19 59 19 59 19 59 19 59 19 59 19 59 19 19 19 19 19 19 19 19 19 19 19 19 19 | 29.4 31.9 30.7 28.9 30.3 31.4 14.4 32.3 919.8 27.3 28.8 19.6 24.8 22.8 30.7 26.7 24.8 22.8 34.7 25.3 36.5 36.5 36.5 36.0 36.3 36.0 36.3 36.0 36.0 36.0 36.0 | 5.9 11.1 9.5 10.6 9.5 7.4 9.6 4.8 9.5 13.7 6.1 11.8 9.4 11.6 11.6 11.6 11.6 11.6 11.6 11.6 11 |

(continued)

Appendix B (cont.)

| Village code | Popula- tion | Live births | Deaths | Birth rate | Death |
|---|--|---|--|---|--|
| VB1 VB2 VB3 VB4 VB5 VB6 VB7 VB8 VB9 VB10 D28 D29 D30 D31 D32 D33 D34 D35 D89 D90 D91 D92 D93 D99 D99 D99 D99 D99 D99 D99 | 1159 990 3002 3742 956 632 279 1375 0 2619 1212 176 720 1049 723 1127 1411 666 1463 1152 1075 0 0 1173 1364 467 819 816 3366 2091 | 35 33 91 93 26 8 6 39 0 84 46 4 27 39 25 29 38 8 30 33 24 0 0 39 52 17 21 26 84 54 | 13 14 30 23 11 5 2 13 0 26 9 1 14 12 7 12 12 6 6 0 0 9 21 4 11 10 28 29 | 30.2 33.3 30.3 24.9 27.2 12.7 21.5 28.4 0 32.1 38.0 22.7 37.5 37.2 34.6 25.7 26.9 12.0 20.5 28.6 22.3 0 0 33.2 38.1 38.1 36.4 25.6 31.9 25.0 25.8 | 11.2 14.1 10.0 6.1 11.5 7.9 7.2 9.5 0 9.9 7.4 5.7 19.4 11.4 9.7 10.6 8.5 9.0 14.4 5.2 5.6 0 0 7.7 15.4 8.6 13.4 12.3 8.3 13.9 |
| omparison rea | 101933 | 2992 | 1037 | | 10.5 |

^{*}See village name in Appendix A.

Appendix C

Life Table Equations

1.
$$_{n}q_{x} = \frac{_{n}m_{x}}{\frac{1}{n} + _{n}m_{x} \left[\frac{1}{2} + ^{n}/_{12} \left(_{n}m_{x} - \ln C\right)\right]}$$

2.
$$\ell_0 = 100,000$$

 $\ell_x = (1 - {}_{n}q_{x-1}) \ell_{x-n}$

3.
$$L_0 = 0.276\ell_0 + 0.724\ell_1$$

$$L_1 = 0.410\ell_1 + 0.590\ell_2$$

$$L_i = \frac{1}{2} (\ell_i + \ell_{i+1}), \quad i = 2, 3, 4$$

$$_nL_x = \frac{_nd_x}{_nm_x} \quad \text{for } 5 \le x \le 80$$

$$_L_{85} = \frac{\ell_{85}}{_L_{85}} \quad \text{for the last age group } 85 + 1$$

4.
$$\dot{e}_x = \frac{T_x}{\ell_x} \quad \text{where } T_x = \frac{\overline{\Sigma}}{2} L_y$$

<u>Note:</u> Greville's method, as suggested in Shryock, H.S., Seigel, J.S., and Associates, <u>The Methods and Materials of Demography</u> (revised), U.S. Dept. of Commerce, Bureau of the Census, 1975, Vol. II p.414 and pp. 444-5.

⁽ $ln\ C$ assumed to be 0.095; separation factors in Equation 3 correspond to an infant mortality rate of 100.)

Appendix D
New Standard Populations

| Age group (years) | Males | Females | Both sexes combined |
|----------------------|---------|---------|---------------------|
| 0 | 2,558 | 2,471 | 2,396 |
| 1-4 | 9,513 | 9,231 | 9,490 |
| 5-9 | 10,824 | 10,472 | 10,649 |
| 10-14 | 9,954 | 9,609 | 9,783 |
| 15-19 | 9,989 | 9,627 | 9,809 |
| 20-24 | 9,477 | 9.137 | 9,308 |
| 25-29 | 8,458 | 8,204 | 8,332 |
| 30-34 | 7,355 | 7,175 | 7,266 |
| 35-39 | 6,585 | 6,476 | 6,531 |
| 40 - 44 | 5,326 | 5,253 | 5,290 |
| 45-49 | 4,341 | 4,335 | 4,338 |
| 50-54 | 3,994 | 4,061 | 4,027 |
| 55-59 | 3,486 | 3,604 | 3,544 |
| 60-64 | 2,912 | 3,179 | 3,045 |
| 65-69 | 2,167 | 2,591 | 2,378 |
| 70-74 | 1,424 | 1,837 | 1,629 |
| 75-79 | 958 | 1,406 | 1,181 |
| 80-84 | 429 | 814 | 602 |
| 85+ | 250 | 518 | 402 |
| Total | 100,000 | 100,000 | 100,000 |

Appendix E

Staff of the DSS, 1993

Dr. Michael A. Strong, DSS Project Director

Matlab Field Station

Supervisory Staff:

Mr. A.M. Sarder, Manager Mr. A.K.M. Nurul Islam, SFRO Mr. Liaquat Ali Mondal, FRO Mr. Md. Ismail, FRO Mr. Md. Khalilur Rahman I, Asst. Supvr.

Senior Health Assistants:

Mr. Md. Sirajul Hoque Mr. K.J.M. Mannan Pathan Mr. M. Abdur Rashid Mia Mr. M.A. Latif Patwari Mr. A.F.M. Aminul Islam Khan

Mr. M.A. Mannan Bakaul Mr. Monoranjan Das Mr. Md. Aftekharuzzaman Mr. Md. Mozammel Haque

Paramedic:

Mr. M. Monirul Alam Bhuiyan

Admin. Assistant:

Mr. A.K.M. Mozibul Hoque

Health Assistants:

Mr. M. Idris Ali Miah I Mr. M. Abul Kashem Mr. M. Idris Ali Miah II Mr. Zahirul Hogue Mr. Md. Nurul Hague Mr. Golam Hossain Mr. P.C. Chakraborty Mr. Md. Jasimuddin Mr. Nasir Ahmed Mr. Md. Shahidur Rahman

Mr. Alfaz Uddin A. Chowdhury

Mr. Md. Sadiquzzaman Mr. Shah Mostafa Kamal Mr. Sheikh Abdul Jabber Mr. Md. A. Malek Patwari Mr. Md. Monirul Hoque

Mr. Jabed Ali

Recorders:

Ms. Shahana Ahmed, HA Ms. Monowara Begum HA

Note: Besides these, 110 CHWs contributed for the DSS data collection.

Dhaka-based Staff

Mr. Saker A. Chowdhury Ms. Lutfun Nahar Mr. Md. Golam Mostafa Mr. Sentu B. Gomes Mr. M.A. Jalil Sarker Ms. Rahima Mazhar Mr. A.B.M. Delwar Hossain

Mr. Md. Kapil Ahmed Mr. Sajal K. Saha Mrs. Habiba Rahman Mr. Md. Khayrul Alam Khan Mr. Md. Arifur Rahim Ms. Nasrin Aktar

Mr. Birendra Nath Adhikary

Dr. Michael A. Strong (left the Project in June/95) Dr. Jeroen K. van Ginneken (joined as Project Director as of Nov/95) Dr. R. Bairagi (acted as Project Director during July-Oct/95) Dr. M.A. Kashem Shaikh (joined as Dhaka-based Manager as of April/96)



Each year, ICDDR,B treats over 70,000 patients attending its two hospitals, one in urban Dhaka, the other in rural Matlab. Though they are planted in Bangladeshi soil, they grow because of the dedication of thousands of concerned people throughout the world. The patients are mostly children with diarrhoea and associated illnesses and the services are offered free to the poorer section of the community

Since these services are entirely dependent on financial support from a number of donors, now we at the ICDDR,B are establishing an entirely new endeavour: an ENDOWMENT FUND. We feel

that, given securely implanted roots, the future of the hospitals can confidently depend upon the harvest of fruit from perpetually bearing vines.

To generate enough income to cover most of the patient costs of the hospitals, the fund will need about five million dollars. That's a lot of money, but look at it this way:

JUST \$150 IN THE FUND WILL COVER THE COST OF TREATMENT FOR ONE CHILD EVERY YEAR FOREVER!

We hope you will come forward with your contribution so that we can keep this effort growing forever or until the world is free of life-threatening diarrhoea. IT IS NOT AN IMPOSSIBLE GOAL.

Cheques may be made out to: ICDDR,B Hospital Endowment Fund.

For more information please call or write to: Chairman, Hospital Endowment Fund Committee GPO Box 128 - Dhaka, 1000, Bangladesh

Telephone: 600-171 through 600-178 Fax: (880-2)-883116