# Chakaria Health and Demographic Surveillance System Focusing on the Poor and Vulnerable

Demographic Events, Safe Motherhood, and Water and Sanitation Practices - 2009

Scientific Report No. 110



# Chakaria Health and Demographic Surveillance System Focusing on the Poor and Vulnerable

Demographic Events, Safe Motherhood, and Water and Sanitation Practices - 2009

S. M. A. Hanifi Farhana Urni Mohammad Iqbal Shahidul Hoque Abbas Bhuiya



The annual reports of the Chakaria Health and Demographic Surveillance System (Chakaria HDSS) are not copyrighted and may be freely quoted as long as the source is properly indicated.

All staff members of the Chakaria HDSS, Dhaka and Chakaria, have contributed to the preparation of this report.

ISBN 978-984-551-313-5

Scientific Report No. 110

September 2010

# Cover and layout design by

Md. Abdur Razzaque

#### Published by

ICDDR,B GPO Box 128, Dhaka 1000 Mohakhali, Dhaka 1212, Bangladesh

Telephone: 8860523-32 (10 lines); Fax: (880-2)-8826050

Email: msik@icddrb.org URL: http://www.icddrb.org

#### **Printed by**

**Print Link Printers** 

#### **ACKNOWLEDGEMENTS**

Data presented in this report were collected through the Chakaria Health and Demographic Surveillance System, one of the activities of the Chakaria Field site of ICDDR,B. The analysis of data for this report was possible with the support extended by the Department for International Development (DFID), UK, through the 'Future Health Systems: Innovation for Equity' – a Research Programme Consortium, grant number GR-00445 and by the Government of Bangladesh through the 'Improved Health for the Poor: Health Nutrition and Population Research' project, grant number GR-00410.

ICDDR,B acknowledges with gratitude the commitment of the Government of Bangladesh and the above development partner to the Centre's research. ICDDR,B also gratefully acknowledges the following donors which provide unrestricted support to the Centre's research efforts: Australian Agency for International Development (AusAID), Government of the People's Republic of Bangladesh, Canadian International Development Agency (CIDA), Embassy of the Kingdom of the Netherlands (EKN), Swedish International Development Cooperation Agency (Sida), Swiss Agency for Development and Cooperation (SDC), and Department for International Development, UK (DFID).

The project team is grateful to the villagers for their cooperation in providing invaluable information. The team is also grateful to Dr. Abdur Razzaque and Mr. Masud Reza for reviewing an earlier version of the report. The untiring efforts of the team members of the Chakaria Field site in maintaining the surveillance system are gratefully acknowledged. The team also acknowledges the efforts of Abdullah Al Mamun in finalizing the report.

# **CONTENTS**

#### **CHAPTER I**

INTRODU	CTION	7
	CHAPTER 2	
METHODS	AND MATERIALS	10
	CHAPTER 3	
POPULATION	ON AND POPULATION CHANGES	12
	CHAPTER 4	
MORTALIT	Y	15
	CHAPTER 5	
FERTILITY .		22
	CHAPTER 6	
MIGRATIO	N	25
	CHAPTER 7	
MARRIAGE	-	29
	CHAPTER 8	
SAFE MOT	HERHOOD PRACTICES	30
	CHAPTER 9	
WATER AN	ID SANITATION PRACTICES	35
TABLES		
Table 1.	Demographic and health indicators, Chakaria HDSS, 1999-2009	1.3
Table 2.	Age-specific death rate (per 1,000 population) by sex, Chakaria HDSS, 2009	
Table 3.	Abridged Life Table, Chakaria HDSS, 2009	
Table 4.	Under-5 mortality rate per 1,000 live births by asset quintile, Chakaria HDSS, 2009	17

Table 5.	Causes of death, Chakaria HDSS, 2004-09
Table 6.	Age-specific fertility rate per 1,000 women aged 15-49 years, Chakaria HDSS, 2009
Table 7.	Crude birth rate per 1,000 population by asset quintile, Chakaria HDSS, 200923
Table 8.	Pregnancy outcome, Chakaria HDSS, 2009
Table 9.	Migration rate per 1,000 population by asset quintile, Chakaria HDSS, 200925
Table 10	). Number of migrants by sex and month, Chakaria HDSS, 200925
Table 1	. Number of migrants by sex and month, intervention area, Chakaria HDSS, 200926
Table 12	2. Number of migrants by sex and month, comparison area, Chakaria HDSS, 200926
Table 13	3. Origin and destination of migrants by sex, Chakaria HDSS, 200927
Table 14	4. Reasons for migration, Chakaria HDSS, 200928
Table 15	5. Age at marriage by sex, Chakaria HDSS, 200929
Table 16	6. Antenatal care by type of sources and asset quintile, Chakaria HDSS, 200931
Table 17	7. Postnatal care by type of sources and asset quintile, Chakaria HDSS, 200932
Table 18	3. Assistance during delivery by asset quintile, Chakaria HDSS, 200933
Table 19	Place of delivery by asset quintile, Chakaria HDSS, 200934
Table 20	). Proportion of caesarean-section delivery by asset quintile, Chakaria HDSS, 200934
Table 2	1. Sources of drinking water by asset quintile, Chakaria HDSS, 1994 and 200935
Table 22	2. Sources of water for bathing by asset quintile, Chakaria HDSS, 1994 and 200935
Table 23	3. Latrine use by asset quintile, Chakaria HDSS, 1994 and 200936
FIGURE	es es estado
Fig. 1.	Map of Chakaria showing intervention and comparison areas9
Fig. 2.	Male and female population by age and sex, Chakaria HDSS, 200912
Fig. 3.	Probability of survival by age and sex, Chakaria HDSS, 2009
Fig. 4.	Age-specific fertility rate, Chakaria HDSS, 200922

Fig. 5. Numb	per of births and deaths by month, Chakaria HDSS, 200924
Fig. 6. Numb	per of births and deaths by month, Intervention area, Chakaria HDSS, 200924
Fig. 7. Numb	per of births and deaths by month Comparison area, Chakaria HDSS, 200924
Fig. 8. Numb	per of marriages by month, Chakaria HDSS, 200929
AUTHORS' C	OMMENTS37
REFERENCES	38
ADDITIONAL	_ READINGS39
Appendix A:	Midyear population by age and sex in the intervention and comparison areas, Chakaria, HDSS, 2009
Appendix B:	Percentage distribution of midyear population by age and sex in the intervention and comparison areas, Chakaria HDSS, 200943
Appendix C:	Number of births by age of mother, Chakaria HDSS, 200944
Appendix D:	Number of deaths by age and sex, Chakaria HDSS, 200945
Appendix E:	Causes of deaths by age and sex, Chakaria HDSS, 200946
Appendix F:	Number of migrants by age and sex, Chakaria HDSS, 200948
Appendix G:	Migration rate per 1,000 population by age and sex, Chakaria HDSS, 2009
Appendix H:	Number of migrants by origin or destination, Chakaria HDSS, 200950
Appendix I:	Number of in-migrants by reasons for migration, Chakaria HDSS, 200951
Appendix J:	Number of out-migrants by reasons for migration, Chakaria HDSS, 200952
Appendix K:	Percentage of population by age and marital status, Chakaria HDSS, 200953
Appendix L:	Percentage of population by age and marital status, intervention area, Chakaria HDSS, 2009
Appendix M:	Percentage of population by age and marital status, comparison area, Chakaria HDSS, 2009
Appendix N:	Chakaria HDSS project team, 200956

# **CHAPTER I**


#### Introduction

Chakaria is one of the 481 *upazilas* (sub-districts) in Bangladesh. It is located between latitudes 21°34′ and 21°55′ North and longitudes 91°54′ and 92°13′ East in the southeastern coast of the Bay of Bengal. Administratively, it is under Cox's Bazar district with an estimated population of 418,607 in 2009. The highway from Chittagong to Cox's Bazar passes through Chakaria. The east side of Chakaria is hilly, while on the west side towards the Bay of Bengal is lowland. A map showing the location of Chakaria is presented in Figure 1.

ICDDR,B started its activities in Chakaria in 1994. The focus of the activities has been to facilitate local initiatives for the improvement of health of the villagers in general and of children, women, and the poor in particular. Thus, the activities of the project have been participatory with emphasis on empowering the people by raising awareness about health, inducing positive preventive behaviour through health education, and providing technical assistance to any health initiatives taken by the village-based indigenous self-help organizations. Some major initiatives taken by the villagers included assessment of health needs, defining actions for health, implementing them, and monitoring their implementation and outputs. Among the health-related activities, identification of volunteers for health education, mobilizing local resources for the establishment of village health posts and their management, introduction of a pre-paid family health card, and establishment of health cooperatives have been the major ones. Details of the activities of the project and the outcomes have been reported elsewhere (1;2). Health services that are currently available in the intervention and comparison areas are presented in the box below. Collection of data from sample households on a quarterly basis, referred hitherto as Chakaria Health and Demographic Surveillance System (Chakaria HDSS), has been initiated in both the areas since 1999. The primary purpose of this surveillance system is to monitor the impact of interventions with equity focus and generate relevant health, demographic and socioeconomic information for policies and programmes, and further research. This report presents data collected through the Chakaria HDSS during 2009.

# Existing health services in the intervention and comparison areas, Chakaria Health and Demographic Surveillance System, 2009

	ipine surveinance system, 2005	
	Comparison area (Two unions with 39,329 population)	
No.	Healthcare facility/provider	No.
	ICDDR,B facilitated and Community initiated	
7	Village health post	0
12	Trained midwife	0
1	Qualified physician	0
10	Male paramedic	0
	Government	
6	Union Health and Family Welfare Centre (UHFWC)	1
216	EPI centre	38
0	Rural dispensary	1
6	Family Welfare Visitor (FWV)	2
3	Sub-Assistant Community Medical Officer (SACMO)/Medical assistant	2
23	Family Welfare Assistant (skilled birth attendant)	1
	Private	
186	Village doctor (allopathic)	54
78	Village doctor (homeopathic)	24
142	Allopathic pharmacy	35
13	Homeopathic pharmacy	2
3	Diagnostic centre	0
	NGO	
4	Health and development activities	4
	7 12 1 10 6 216 0 6 3 23 186 78 142 13 3	No. Healthcare facility/provider  ICDDR,B facilitated and Community initiated  7 Village health post  12 Trained midwife  1 Qualified physician  10 Male paramedic  Government  6 Union Health and Family Welfare Centre (UHFWC)  216 EPI centre  0 Rural dispensary  6 Family Welfare Visitor (FWV)  3 Sub-Assistant Community Medical Officer (SACMO)/Medical assistant  23 Family Welfare Assistant (skilled birth attendant)  Private  186 Village doctor (allopathic)  78 Village doctor (homeopathic)  142 Allopathic pharmacy  13 Homeopathic pharmacy  14 Diagnostic centre  NGO



#### **Methods and Materials**

The Chakaria HDSS covered 8 unions<sup>1</sup>, namely Baraitali, Kayerbil, Bheola Manik Char, Paschim Boro Bheola, Shaharbil, Kakara, Harbang, and Purba Boro Bheola. Of these, the last two unions formed the comparison area, and the first 6 formed the intervention area. In 1999, 106,320 people were living in 20,252 households in the intervention area and 34,418 people were living in 6,727 households in the comparison area (3). A household is defined as blood or otherwise related group of members and unrelated individuals living in the same compound at least once a month and sharing the food from the same kitchen. A household member is considered to have migrated out if s/he has left the household and does not intend to come back within six months of the time s/he left. A person is considered to have migrated in if s/he was not previously included in the list of household members and intends to live in the household for at least once in a month for the next six months.

Although Chakaria HDSS started in 1999 covering all the households in 8 unions, data collection was interrupted during 2001-2003. Since 2004, quarterly data collection has resumed, and data are being collected from 3,727 and 3,315 systematically randomly-chosen households in the intervention and comparison areas respectively. 27 field-trained workers collected data during 2009. The data collectors were also provided with written instructions for specific questions that required added explanations.

Six supervisors supervised the data-collection process. To detect any anomalies, the supervisors re-visited 5% of the households, chosen randomly, within 2 days of data collection by the field workers. Later on, the supervisors and the relevant field workers together sorted out any inconsistencies in the collected data. All the filled-up questionnaires were manually checked for completeness and for any inconsistencies. Subsequently, computer-based data-editing procedures were applied to ensure the quality of data.

Asset quintiles based on ownership of various assets by any member of the households were used to examine differences in various dependant variables. The list of assets included almirah, table/chair, van/rickshaw, *choki/khat*, radio, television, cycle, motorcycle, fridge, sofa, electric fan, sewing machine, telephone and electricity. The principal component analytical technique was used for calculating weights of the assets to derive household asset index scores

10


<sup>&</sup>lt;sup>1</sup> Government has restructured the existing 8 unions into 11 in 2005.

(4). The major demographic indicators, safe motherhood and water and sanitation related practices have been tabulated for the various asset quintiles.

It should be mentioned that the number of observations in the tables presented in this report differ in some instances due to missing information for some variables.

# **Population and Population Changes**

The population pyramid based on the sample households is presented in Figure 2. The shape of the pyramid is typical of a developing country with declining mortality and fertility. The sex ratio (male per 100 females) was 104 in 2009. The age dependency ratio<sup>2</sup> was 75 in 2009 (see Appendix A).



The major demographic and health indicators in the intervention and comparison areas during 1999, and 2004-2009 are presented in Table 1. A declining trend in the fertility indicators and natural rate of population increase has been observed during 1999-2009. Most of the rates in Chakaria HDSS area are much higher than those in the Matlab HDSS area, another rural field site of ICDDR,B (5). In 2009, the rate of natural increase and the annual population growth rate in the surveillance area was 1.6 % and 0.6% respectively (Table 1).

**12** 

<sup>&</sup>lt;sup>2</sup> The age dependency ratio represents the ratio of the combined child population (under 15) and aged population (65 and over) to the population of intermediate age (15 to 64).

12.0% of births in Chakaria were delivered at facilities (Hospital or Clinic) in 2009. The percentage of births at facilities decreased from 14.4 in 2008 to 12.0 in 2009. 10.9% of the births were attended by Skilled Birth Attendant (SBA). There has been a decrease in deliveries by SBAs from 16.2% in 2008 to 10.9% in 2009 (Table 1).

The legal age of marriage is 18 years for female and 21 years for male in Bangladesh. In 2009, 39.3% of the women married before reaching their 18th birth day. The percentage of underage female marriage decreased to 39.3% in 2009 from 47.5% in 2008. 24.8% of the males were married before the age of 21 years in 2009. The proportion of male marriages before 21 years has remained similar between 2008 and 2009. The percentage of underage marriage for females remained higher than males during 2004 to 2009.

			Cha	akaria H	DSS			Matlab HDSS
Rates per 1,000	1999	2004	2005	2006	2007	2008	2009	2008
Crude birth rate								
Intervention area	33.8	30.6	29.8	25.8	26 .9	24.7	23.7	23.5
Comparison area	33.9	28.8	27.4	25.3	27.2	26.5	21.9	22.1
Both areas	33.9	29.7	28.7	25.6	26.6	25.5	22.9	22.9
Total fertility rate*								
Intervention area	5.1	4.6	4.4	3.5	3.6	3.3	2.9	2.7
Comparison area	4.9	4.4	4.0	3.3	3.4	3.4	2.6	2.7
Both areas	5.1	4.5	4.2	3.4	3.5	3.3	2.8	2.7
Neonatal mortality**								
Intervention area	40.0	24.8	25.2	33.7	27.0	25.0	29.1	15.8
Comparison area	47.3	40.8	35.9	42.3	44.3	33.5	46.8	26.1
Both areas	41.7	31.9	31.5	37.6	34.8	29.0	36.8	20.7
Post-neonatal mortality**	ŧ .							
Intervention area	21.2	15.5	14.1	17.7	18.0	23.3	25.7	4.9
Comparison area	22.4	19.7	25.1	15.4	7.4	5.6	15.6	10.4
Both areas	21.4	17.4	17.4	16.6	13.3	14.9	21.3	7.5
Infant mortality rate**								
Intervention area	61.2	40.3	39.3	51.4	45.0	48.3	54.8	20.6
Comparison area	69.7	60.5	61.0	57.7	51.7	39.0	62.5	36.4
Both areas	63.2	49.3	48.9	54.2	48.0	43.9	58.1	28.1
Child mortality rate (1-4	yrs)							
Intervention area	9.0	8.1	7.5	6.2	4.7	3.6	5.2	3.0
Comparison area	10.6	5.5	5.3	2.4	4.4	6.2	4.1	2.9
Both areas	9.4	6.9	6.5	4.5	4.6	4.7	4.7	3.0

Table 1. (Contd...)

Rates per 1,000			Ch	akaria F	HDSS			Matlal HDS
	1999	2004	2005	2006	2007	2008	2009	2008
Crude death rate								
Intervention area	6.7	5.9	5.8	5.4	5.4	5.7	6.8	6.4
Comparison area	7.9	7.0	6.5	5.7	6.8	6.7	6.1	7.
Both areas	7.0	6.3	6.1	5.6	6.1	6.1	6.5	6.
Rate of natural increase								
Intervention area	27.1	24.7	24.0	20.4	21.7	19.4	16.9	17.
Comparison area	26.0	21.8	20.8	19.6	19.2	21.0	15.8	14.
Both areas	26.9	23.4	22.5	20.0	20.6	20.2	16.4	16.
In-migration rate								
Intervention area	-	17.1	24.5	29.7	23.4	27.1	32.0	
Comparison area	-	16.6	23.7	30.0	26.0	26.0	27.1	
Both areas	-	16.9	24.1	29.9	24.6	26.6	29.8	44.
Out-migration rate								
Intervention area	-	22.2	23.8	33.8	31.0	36.2	38.8	
Comparison area	-	19.5	25.9	34.3	33.2	34.7	42.9	
Both areas	-	21.0	24.8	34.0	32.0	35.5	40.6	65
Growth rate (%)		2.0	2.5	1.6	4.4	1.0	1.0	
Intervention area	-	2.0	2.5	1.6	1.4	1.0	1.0	
Comparison area Both areas	-	1.9 1.9	2.0 2.1	1.5 1.6	1.3 1.3	1.2 1.1	$0.4 \\ 0.6$	- 0
Facility-based delivery (%)		1.7	2.1	1.0	1.5	1.1	0.0	- 0
Intervention area	_	6.8	6.4	6.2	3.8	18.3	14.3	
Comparison area	_	4.4	3.8	4.5	6.8	9.5	9.2	
Both areas	_	5.4	4.9	5.4	5.1	14.4	12.0	
Received assistance from SBA during delivery (%)								
Intervention area	_	14.3	9.2	16.5	20.4	18.0	10.7	
Comparison area	-	14.8	11.6	13.8	18.2	12.8	11.1	
Both areas	-	14.5	10.3	15.3	19.1	16.2	10.9	
Male marriage at ages ınder 21 years (%)								
Intervention area	-	23.4	25.6	26.3	25.2	25.6	21.8	
Comparison area	-	23.3	23.8	29.7	26.0	23.8	28.1	
Both areas	-	23.3	24.7	27.9	25.6	24.7	24.8	
Female marriage at ages ander18 years (%)								
Intervention area	-	51.4	43.1	51.2	40.4	46.0	40.2	
Comparison area	-	56.6	52.0	48.4	46.7	49.0	38.5	
Both areas	-	53.6	47.3	49.8	43.2	47.5	39.3	

# **Mortality**

Age-specific mortality rates by area and sex are presented in Table 2. The crude death rate for the intervention and comparison areas in Chakaria, when considered together, was 6.5 per 1,000 populations in 2009. The rate was slightly higher in the intervention area than in the comparison area. Infant mortality rate for all the villages in the intervention and comparison areas was 58.1 per 1,000 live births with a lower rate in the intervention area than in the comparison area. Child mortality rate was 4.7 per 1,000 children aged 1-4 years in the intervention and comparison areas combined. The rate was higher in the intervention area than in the comparison area (Table 2).

Abridged Life Table for males and females are presented in Table 3. Life expectancy at birth was 67.4 years for males and 66.7 years for females. The rate of mortality of children aged less than 5 years (under-five mortality) was 76.8 per 1,000 live births in Chakaria in 2009 (Table 4). Figure 3 shows the probability of survival by sex during various age groups. The probability of survival of females remained same males up to age 45 years, but after age 45 probability of survival oscillated.

Table 2		pecific of 5, 2009	leath ra	ite (per 1	l,000 pc	pulatio	on) by se	x, Chaka	aria
Age	Inte	rvention a	rea	Con	nparison a	area	]	Both areas	
(years)	Male	Female	Both	Male	Female	Both	Male	Female	Both
<1*	66.7	42.3	54.8	51.7	74.1	62.5	60.2	56.0	58.1
1-4	4.7	5.7	5.2	2.0	6.5	4.1	3.5	6.1	4.7
5-9	0.6	1.3	0.9	0.8	0.8	0.8	0.7	1.1	0.9
10-14	1.1	0.6	0.9	0.7	0.0	0.3	0.9	0.3	0.6
15-19	1.2	1.8	1.5	0.7	2.0	1.4	1.0	1.9	1.4
20-24	3.1	1.7	2.4	0.9	3.0	1.9	2.1	2.3	2.2
25-29	0.0	1.4	0.6	1.3	0.0	0.7	0.6	0.7	0.7
30-34	6.6	0.0	3.0	0.0	0.0	0.0	3.6	0.0	1.7
35-39	0.0	3.4	1.7	0.0	1.8	1.0	0.0	2.6	1.3
40-44	1.9	5.6	3.8	2.5	2.3	2.4	2.2	4.1	3.2
45-49	16.4	8.8	12.7	6.7	4.6	5.7	11.8	6.8	9.3
50-54	13.2	5.0	9.0	6.2	8.0	7.1	10.0	6.5	8.1
55-59	18.2	6.9	12.9	17.2	16.6	16.9	17.8	11.8	15.0
60-64	18.1	19.4	18.7	9.8	13.9	11.5	14.6	17.2	15.7
65-69	38.7	46.1	42.0	32.8	32.5	32.6	35.7	39.1	37.3
70-74	14.9	29.4	21.2	27.8	85.4	52.6	20.6	54.6	35.2
75-79	48.8	51.5	50.0	78.9	68.5	73.8	60.3	58.8	59.6
80-84	115.4	45.5	83.3	81.1	32.3	58.8	101.1	40.0	73.2
85+	147.5	314.3	208.3	102.6	236.8	168.8	131.3	274.0	191.9
All	7.5	6.1	6.8	5.3	6.9	6.1	6.5	6.5	6.5
*Per 1,000	live birth	s; HDSS=He	alth and I	Demographi	c Surveilla	nce Syster	n.		

Table	3. Abr	idged I	ife Tabl	le, Chak	aria H	IDSS, 20	009			
Age			Male					Female		
(years)	$_{n}m_{x}$	$_{n}q_{x}$	$l_x$	$_{n}L_{x}$	$e_{x}$	$_{n}m_{x}$	$_{n}q_{x}$	$_{n}l_{x}$	$_{n}L_{x}$	$e_{x}$
0	0.0626	0.0626	100,000	94,990	67.4	0.0554	0.0554	100,000	95,564	66.7
1	0.0035	0.0138	93,738	372,522	70.9	0.0061	0.0240	94,455	373,550	69.7
5	0.0007	0.0034	92,444	461,506	67.9	0.0011	0.0053	92,185	459,793	67.3
10	0.0009	0.0046	92,134	459,690	63.1	0.0003	0.0016	91,694	458,128	62.7
15	0.0010	0.0049	91,709	457,505	58.4	0.0019	0.0093	91,546	455,774	57.8
20	0.0021	0.0104	91,258	454,098	53.6	0.0023	0.0113	90,698	451,132	53.3
25	0.0006	0.0030	90,308	450,907	49.2	0.0008	0.0037	89,676	447,605	48.9
30	0.0036	0.0180	90,033	446,420	44.3	0.0000	0.0000	89,340	446,699	44.0
35	0.0000	0.0000	88,411	442,056	40.1	0.0026	0.0129	89,340	444,040	39.0
40	0.0022	0.0107	88,411	439,872	35.1	0.0041	0.0204	88,187	436,780	34.5
45	0.0118	0.0573	87,464	425,658	30.4	0.0068	0.0333	86,388	425,285	30.2
50	0.0100	0.0488	82,451	402,910	27.1	0.0064	0.0317	83,515	411,434	26.1
55	0.0177	0.0851	78,428	376,528	23.4	0.0118	0.0576	80,864	393,474	21.9
60	0.0146	0.0704	71,754	346,991	20.3	0.0171	0.0822	76,203	366,375	18.1
65	0.0357	0.1646	66,704	307,427	16.6	0.0392	0.1793	69,940	319,806	14.4
70	0.0207	0.0985	55,724	265,746	14.4	0.0543	0.2402	57,399	253,693	12.0
75	0.0603	0.2630	50,234	219,095	10.7	0.0588	0.2574	43,611	190,838	10.0
80	0.1011	0.4032	37,022	147,604	8.6	0.0395	0.1804	32,385	147,998	7.6
85+	0.1300	1.000	22,096	169,968	7.7	0.2740	1.000	26,543	96,884	3.6

HDSS = Health and Demographic Surveillance System.

The Abridged life table is constructed applying the Greville's method illustrated in "The Methods and Materials of Demography", edited by Jacob S. Siegel and David A. Swanson, Second edition; Elsevier Academic Press, 2004: 301-40.

 $_{n}m_{x}$  = Central mortality rate

 $_{n}q_{x}$  = Probability of dying between the ages x and x+n;

 $<sup>\</sup>begin{array}{ll} {}_{n}q_{x} & = {}_{n}m_{x}/[(1/n) + {}_{n}m_{x}\{1/2 + n/12({}_{n}m_{x} \text{-logec})\}]; \\ & \log {}_{e}c\text{=}.095 \end{array}$ 

 $l_x$  = Survivors to exact age x

 $_{n}L_{x}$  = Numbers of years lived by the total of the cohort of 100,000 births in the interval;  $L_{0}$ =.20 $l_{0}$ +.80 $l_{1}$ ,  $L_{85+}$ = $l_{85+}/m_{85+}$ 

 $e_x$  = Life expectancy at age x




Table 4 presents under-5 mortality rates by household asset quintiles. Under-5 mortality rate was inversely correlated with household asset scores. The mortality rate of children from the lowest quintile was nearly 7 times of children from the highest quintile.

Cilakaila	HDSS, 2009		
Asset quintile	Number of births	Number of under-5 deaths	Under-5 mortality rate
Lowest	233	27	115.9
Second	170	15	88.2
Medium	153	16	104.6
Fourth	197	14	71.1
Highest	237	4	16.9
All	990	76	76.8

#### Causes of death

Causes of death were recorded as reported by the informed household members. A physician classified the reported causes of death with medical synonyms. Table 5 presents the number of deaths from various causes in the year 2004-2009. Stroke, senility, asthma, neoplasm, neonatal, respiratory infections, drowning, hepatitis, accident, and diabetes were the 10 leading causes of death in Chakaria in 2009.

		No. of deaths	42	37	31	29	26
	2009	Cause	Stroke	Senility	Asthma/ Bronchitis	Neoplasm (Benign and Malignant)	Neonatal (Premature and LBW, Birth asphyxia, Bone trauma, Sepsis and infection)
		No. of deaths	33	33	26	22	19
	2008	Cause	Stroke	Neoplasm (Benign and Malignant)	Asthma/ Bronchitis	Respiratory infections	Senility
		No. of deaths	30	29	26	25	25
	2007	Cause	Asthma/ Bronchitis	Neoplasm (Benign and Malignant)	Respiratory infections	Senility	Stroke
6		No. of deaths	31	28	26	26	21
SS, 2004-0	2006	Cause	Stroke	Senility	Asthma/ Bronchitis	Respiratory infections	Neoplasm (Benign and Malignant)
ria HD		No. of deaths	29	28	28	23	19
Table 5. Causes of Death, Chakaria HDSS, 2004-09	2005	Cause	Stroke	Respiratory infections	Senility	Neoplasm Benign and Malignant)	Asthma Bronchitis
s of Do		No. of deaths	39	30	26	17	15
le 5. Cause	2004	Cause	Respiratory infections	Senility	Asthma/ Bronchitis	Neonatal (Premature and LBW, Birth asphyxia, Birth trauma, Sepsis and infection	Diarrheal disease
Tab	Pant	Nallh	1	2	3	4	v

Table 5. (Contd...)

$\Box$	of hs	2	- <del> </del>		 	 		 
	No. of deaths	22	14	6	<b>∞</b>	7	7	5
2009	Cause	Respiratory infections	Drowning	Hepatitis	Accident	Diabetes	Diarrheal	Cardiovas- cular other than stroke and hype- rtension
	No. of deaths	13	6	6	10	7	_	S
2008	Cause	Hepatitis	Accident	Drowning	Cardiovas- cular other than stroke and hyper- tension	Diarrheal diseases	Hyper- tension	Maternal death
	No. of deaths	24	16	11	6	6	∞	ν
2007	Cause	Neonatal (Premature and LBW, Birth asphyxia, Birth trauma, Sepsis and infection)	Accident	Cardiovas- cular other than stroke and hyper- tension	Diarrheal Diseases	Hepatitis	Drowning	Nutritional diseases
	No. of deaths	15	11	_	9	9	3	ε
2006	Cause	Neonatal (Premature and LBW, Birth asphyxia, Birth trauma, Sepsis and	Drowning	Hepatitis	Accident	Diarrheal diseases	Diabetes	Hyper- tension
	No. of deaths	41	12	10	∞	7	9	9
2005	Cause	Neonatal (Premature and LBW, Birth asphyxia, Bone trauma, Sepsis and infection)	Drowning	Accident	Cardiovascul ar other than stroke and hypertension	Nutritional	Diarrheal diseases	Hepatitis
	No. of deaths	41	14	14	12	11	∞	œ
2004	Cause	Hepatitis	Neoplasm (Benign and Malignant)	Cardiovasc ular other than stroke and hyper- tension	Stroke	Accident	Malaria	Drowning
ı	Rank	9		∞	6	10	11	12

Table 5. (Contd...)

deaths a culosis   7 Typhoid 4 Cardiovascular 3 Hyper- 3 troke and hypertension diseases 3 Rabies 2 Rabies 3 Rabi		2004		2005		2006		2007	,	2008		2009	
Tuber- 4 Malaria 3 Diabetes 3 culosis  Typhoid 4 Cardiovascular 3 Hyper- 3 other than stroke and hypertension  Urinary 4 Urinary 3 Urinary 3 diseases 3 Rabies 3 Epilepsy 2  Rabies 3 Rabies 3 Malaria 2  Maternal 3 Tuberculosis 3 Maternal 2  Diabetes 3 Burn 2 Geath  Hyper- 3 Digestive 2 Suicide 2 Homicide 3 Nutritional 2 Tuber- 2 Homicide 3 Nutritional 2 Tuber- 2 Homicide 3 Nutritional 2 Tuber- 2	Cause		No. of deaths	Cause	No. of deaths	Cause	No. of deaths	Cause	No. of deaths	Cause	No. of deaths	Cause	No. of deaths
Typhoid 4 Cardiovascular 3 Hyper- 3 other than stroke and hypertension  Urinary 4 Urinary 3 Urinary 3 diseases  Rabies 3 Rabies 3 Epilepsy 2  Maternal 3 Tuberculosis 3 Malaria 2  Diabetes 3 Burn 2 Maternal 2  Hyper- 3 Digestive 2 Suicide 2  Homicide 3 Nutritional 2 Tuber- 2  Homicide 3 Nutritional 2 Tuber- 2  Homicide 3 Nutritional 2 Tuber- 2	Nutritional		7	Tuber- culosis	4	Malaria	3	Diabetes	3	Diabetes	4	Suicide	8
Urinary4Urinary3Urinary3Rabies3Rabies3Epilepsy2Maternal death3Tuberculosis death3Maternal death2Diabetes3Burn2Maternal death2Hyper- tension3Digestive diseases2Suicide dath2Homicide3Nutritional diseases2Tuber- culosis2	Homicide		7	Typhoid	4	Cardiovascular other than stroke and hypertension	က	Hyper- tension	æ	Digestive disease	က	Epilepsy	7
Rabies3Rabiesysy2Maternal death3Tuberculosis Amalaria2Diabetes tension3Burn death2Maternal death2Hyper- tension3Digestive diseases2Suicide 22Homicide diseases3Nutritional diseases2Tuber- 22	Hyper- tension		9	Urinary	4	Urinary diseases	m	Urinary diseases	m	Neonatal (Premature and LBW, Birth asphyxia, Bone trauma, Sepsis and infection)	13	Hyper- tension	2
Maternal3Tuberculosis3Malaria2Diabetes3Burn2Maternal2Hyper-3Digestive2Suicide2tensiondiseases2Tuber-2Homicide3Nutritional2Tuber-2diseasesculosis	Diabetes	i	S	Rabies	3	Rabies	8	Epilepsy	7	Tuber- culosis	4	Nutri- tional	2
Diabetes 3 Burn 2 Maternal 2 Hyper- 3 Digestive 2 Suicide 2 tension diseases Homicide 3 Nutritional 2 Tuber- 2 diseases	Urinary diseases	i	S	Maternal death	3	Tuberculosis	3	Malaria	7	Burn	2	Other urinary	2
Hyper-3Digestive2Suicide2tensiondiseasesHomicide3Nutritional2Tuber-2diseasesculosis	Typhoid	i	2	Diabetes	3	Burn	2	Maternal death	2	other urinary	2	Burn	
3 Nutritional 2 Tuber- 2 diseases culosis	Digestive disease		2	Hyper- tension	8	Digestive diseases	2	Suicide	2	homicide	Н	Digestive disease	н
	Maternal death		П	Homicide	3	Nutritional diseases	7	Tuber- culosis	7	nutritional	1	Disease of uterus	1

Table 5. (Contd...)

-	2004		2005		2006		2007		2008		2009	
Kank	Cause	No. of Cause deaths	Cause	No. of deaths	Cause	No. of Cause deaths	Cause	No. of Cause deaths	Cause	No. of deaths	Cause	No. of deaths
21	Suicide	1	Burn	2	Congenital anomalies	1	Typhoid	2	rabies	1	Dysentery	
22	Unknown	36	Malaria	T	Leprosy	T	Dysentery		snake bite	T	Epestaxis	-
23	1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1	Congenital anomalies		Tetanus		Digestive disease		suicide		Homicide	
24	1	1 1 1 1 1 1 1	Digestive disease	-	Unknown	42	Homicide	1	typhoid	-	Malaria	-
25		1 1 1 1 1 1 1	Suicide			1 1 1 1 1 1 1	Rabies	1	malaria	1	Maternal death	-
26		 	Snake bite			1 1 1 1 1 1 1	Unknown	32	unknown	46	Tuber- culosis	-
27		1 1 1 1 1 1 1 1	Epilepsy	Π	1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1 1 1		Unknown	35
			Unknown	46								
Total		280		271		249		274		274		292
HDSS	HDSS = Health and Demographic Surveillance System.	emograț	ohic Surveillan	ce Syster	n.							

# **Fertility**

The crude birth rate in 2009 was 22.9 per 1,000 populations, which was lower than the rate in 2008 (25.5 per 1,000 population) (Table 1). Total fertility rates per woman also showed a downward trend during 1999-2009 with a value of 2.8 in 2009 (Table 1). The fertility rate was highest among women of age-group of 20-24 years (Fig. 4 and Table 6).

Table	Table 6. Age-specific fertility rate per 1,000 women aged 15-49 years, Chakaria HDSS, 2009									
	Inter	vention a	rea	Com	parison a	ırea	Во	oth areas		
Age	No. of	No. of	Birth	No. of	No. of	Birth	No. of	No. of	Birth	
(years)	females	births	rate	females	births	rate	females	births	rate	
15-19	1,699	150	88.3	1,525	134	87.9	3,224	284	88.1	
20-24	1,206	218	180.8	999	148	148.1	2,205	366	166.0	
25-29	736	100	135.9	598	88	147.2	1,334	188	140.9	
30-34	706	75	106.2	554	55	99.3	1,260	130	103.2	
35-39	599	30	50.1	560	20	35.7	1,159	50	43.1	
40-44	533	9	16.9	437	3	6.9	970	12	12.4	
45-49	456	2	4.4	431	0	0.0	887	2	2.3	
Total	5,935	584	582.5	5,104	448	525.0	11,039	1,032	555.9	
TFR			2,912			2,625			2,780	
TFR = To	tal fertility 1	rate per 1,0	000 wome	en; HDSS = Hea	alth and I	Demograp	hic Surveilland	ce System		





Table 7. Crude h HDSS, 2	oirth rate per 1,000 po 2009	pulation by asset qui	ntile, Chakaria
Asset quintile	Midyear population	Number of births	Birthrate
Lowest	8,969	233	26.0
Second	9,459	170	18.0
Medium	6,464	153	23.7
Fourth	7,818	197	25.2
Highest	9,160	237	25.9
All	41,870	990	23.6
HDSS = Health and De	emographic Surveillance System	ı.	


Table 7 presents the crude birth rates by household asset quintiles. The crude birth rate showed a 'U' shaped relationship with household socioeconomic status measured by asset quintiles.


Of the pregnancies in 2009, 12.5% of 1,268 were terminated prematurely and spontaneously, 3.7% were terminated through induction, and 2.4% resulted in stillbirths (Table 8).

Table 8. Pregnancy outco	ome, Cha	Karia Hi	<b>733</b> , 2009			
, , , , , , , , , , , , , , , , , , ,	Intervent	tion area	Compar	rison area	Both	1 areas
Pregnancy outcome	No.	%	No.	%	No.	%
Spontaneous abortion	87	12.3	71	12.7	158	12.5
Induced abortion	27	3.8	20	3.6	47	3.7
Stillbirth	11	1.6	20	3.6	31	2.4
Live birth*	584	82.4	448	80.1	1,032	81.4
Total number of pregnancies	709	100.0	559	100.0	1,268	100.0
*Multiple live births included HDSS = Health and Demographic	Surveillance S	System.				

Distribution of births and deaths by month did not show any distinct seasonal pattern (Fig. 5). The seasonal patterns of birth and death were similar in the intervention and comparison areas (Fig. 6 and 7).







# **M**igration

In 2009, the rate of out-migration was higher at 41.3 per 1,000 population than that of in-migration at 28.3 per 1,000 population (Table 9). The rates were higher than in 2008. Monthly data on migration are presented in Tables 10, 11 and 12. Data showed that the number of in-migrants was lower than that of out-migrants during 2009 in both the areas. The sex differential in migration was also not prominent. The rate of in-migration among males was highest in January, and the rate was highest among female in May. The rate of out-migration was highest among the males in December and females was highest in May.

	ation rate per 1,000 p S, 2009	opulation by asset	quintile, Chakaria
Asset quintile	Midyear population	In-migration rate	Out-migration rate
Lowest	8,969	23.1	29.4
Second	9,459	21.2	35.1
Medium	6,464	22.1	34.3
Fourth	7,818	31.7	43.5
Highest	9,160	42.1	62.3
All	45,098	29.8	40.6

Month	Ir	n-migration		Ou	t-migration	
Month	Male	Female	Both	Male	Female	Both
January	61	84	145	89	97	186
February	33	79	112	56	69	125
March	38	65	103	60	76	136
April	30	42	72	63	70	133
May	46	98	144	64	136	200
June	48	86	134	67	110	177
July	27	63	90	57	93	150
August	37	47	84	57	78	135
September	49	55	104	70	60	130
October	47	74	121	52	71	123
November	39	62	101	54	76	130
December	52	80	132	94	114	208
All	507	835	1,342	783	1,050	1,833

Table 11. Number of migrants by sex and month, intervention area, Chakaria HDSS, 2009 In-migration Out-migration Month Male Female Both Male Female Both January February March April May June July August September October November December All 

HDSS = Health and Demographic Surveillance System.

Table 12. Nur Cha	mber of mig karia HDSS,		and mont	th, compa	arison area,	
Month	In	-migration		Ου	ıt-migration	
MOIIII	Male	Female	Both	Male	Female	Both
January	22	33	55	50	54	104
February	14	35	49	26	30	56
March	15	25	40	31	45	76
April	7	13	20	33	33	66
May	19	47	66	34	68	102
June	21	36	57	31	44	75
July	12	30	42	27	41	68
August	16	26	42	26	34	60
September	20	26	46	30	29	59
October	17	34	51	26	38	64
November	8	22	30	21	28	49
December	18	39	57	47	53	100
All	189	366	555	382	497	879

#### Origin and destination of migrants

During 2009, 4.8% of 1,342 in-migrants moved into Chakaria HDSS households from outside of Bangladesh whereas 9.3% of 1,831 out-migrants moved out of Bangladesh from Chakaria HDSS area. The proportion of migrants that moved out of Bangladesh was higher than the proportion of migrants that moved into Bangladesh. Overall, the rates of movement of people to and from Chakaria were similar (Table 13).

Table 13. Origin and of HDSS, 2009	destina	tion of n	nigrants b	y sex, Chal	karia	
,	I	n-migratio	n	Οι	ıt-migration	
Origin or destination	Male (%)	Female (%)	Both (%)	Male (%)	Female (%)	Both (%)
Inside Bangladesh	88.0	99.5	95.2	79.3	99.1	90.7
Outside Bangladesh	12.0	0.5	4.8	20.7	0.9	9.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
Total number of migrants	507	835	1,342	781	1,050	1,831
Inside Chakaria	79.4	73.7	75.6	74.2	73.1	73.5
Outside Chakaria	20.6	26.3	24.4	25.8	26.9	26.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Total number of migrants	412	791	1,203	493	877	1,370
Inside HDSS area	58.3	54.1	55.5	59.1	55.3	56.6
Outside HDSS area	41.7	45.9	44.5	40.9	44.7	43.4
Total	100.0	100.0	100.0	100.0	100.0	100.0
Total number of migrants	367	714	1,081	421	760	1,181
HDSS = Health and Demograph	nic Surveil	lance Syster	n.			

# Reasons for migration

Table 14 presents the reasons of migration by sex. 39.6% of the migrants moved out due to family-related issues - mostly marriage, followed by housing (28.5%), work (26.9%), and education (5.0%). Reasons for moving out for

males were different from those of females. 51.5% of male in-migrants moved due to work related issues whereas only 9.1% of the females moved due to that reason. On the other hand, 59.1% of female in-migrants moved due to family related issues - mostly marriage, while only 12.6% of males moved due to family related reasons (Table 14). The reasons of movement for out-migration were mostly similar to the reasons for in-migration.

Table 14. Reasons for m	igration,	Chakaria	HDSS, 2	2009		
	In	ı-migration		Οι	ıt-migration	
Reasons for migration	Male (%)	Female (%)	Both (%)	Male (%)	Female (%)	Both (%)
Family-related	12.6	59.1	41.8	15.7	58.1	39.6
Work-related	51.5	9.1	24.9	48.8	9.6	26.9
Housing-related	26.6	25.6	26.0	28.5	28.2	28.5
Education	9.3	6.2	7.3	7.0	4.1	5.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Total number of migrants	507	835	1,342	783	1,050	1,833
HDSS = Health and Demographic	Surveillance	System.				

### **Marriage**


In total 861 marriages took place in the surveillance households in Chakaria during 2009. The highest number of marriages took place in May and the lowest in September. The number of marriages showed a downward trend from May to September (Fig. 8).

Table 15 presents singulate mean age at marriage (SMAM) and median age at first marriage. The SMAM was 27 years for males and 20 years for females. The SMAM in 2009 remained same as of 2008 for males and females. The median age at first marriage for males and females were 27 and 20 years. Both the indicators for males and females were almost positively associated with household socioeconomic status.

Table 1	5. Age <i>a</i>	nt marriage by sex, Chakar	ia HDSS,	2009
Asset		Male		Female
quintile	SMAM*	Median age at first marriage*	SMAM*	Median age at first marriage
Lowest	23.7	22.8	18.8	19.9
Second	26.1	25.3	19.8	20.5
Medium	27.1	26.1	19.4	19.7
Fourth	28.1	27.8	19.7	20.4
Highest	29.5	28.8	20.6	21.0
All	27.3	26.6	19.7	20.3

HDSS = Health and Demographic Surveillance System. SMAM = Singulate mean age at marriage

<sup>\*</sup> The SMAM and median age at first marriage are calculated by applying indirect methods illustrated in "The Methods and Materials of Demography", edited by Jacob S. Siegel and David A. Swanson, Second edition; Elsevier Academic Press, 2004: 196-202.



#### **Safe Motherhood Practices**

The health-related activities of ICDDR,B in Chakaria included facilitation of provision of safe motherhood services (e.g. antenatal care, postnatal care, and delivery services) by the trained midwives who were based in the seven village health posts that had been established and managed by the villagers since the late nineties. The services provided by these midwives were not strictly restricted to the intervention area. The women from the comparison area also availed their services to some extent. Apart from this, the physicians employed by ICDDR,B with financial support from the community, also provided healthcare services once a week to the villagers from these village health posts during 1998 and 2005.

At present, the Upazila Health Complex of the government and four private hospitals provide healthcare services at the headquarters of Chakaria. At the union level, 6 Union Health and Family Welfare Centres (UHFWCs) of the government and 7 village health posts which were initiated by the community members provide healthcare services in the intervention area. At the same level, one UHFWC and one Rural Dispensary (RD) of the government provide health services in the comparison area. The Family Development Services and Research (FDSR), an NGO, also provides healthcare services both in intervention and comparison areas.

# Use of antenatal care services

During 2009, 65.0% of 987 pregnant women in Chakaria received at least one antenatal check-up (ANC). The percentage of women receiving at least one ANC was higher in the intervention area (69.4%) than in the comparison area (59.5%). The women in the intervention area received services from various sources. Among these sources, the nurses/doctors were dominant, followed by midwives and then FDSR and BRAC. In the comparison area, the dominant source was also Nurse/doctor, but was followed by FDSR and BRAC and then midwives. (Table 16).

The use of ANC was very inequitable in both the intervention and the comparison areas. Of the various sources, services from nurse and doctor has been the most inequitable (Table 16).

Table 16. A	Antenata HDSS, 20		type of s	sources	and asset	quintile	e, Chak	aria
Area	Asset quintile	Received any ANC	Midwife*	FWV*	Nurse/ doctor*	FDSR/ CMH*	None	No. of women
		(%)	(%)	(%)	(%)	(%)	(%)	
	Lowest	56.1	24.4	4.9	14.6	12.2	43.9	123
	Second	67.3	21.4	8.2	19.4	18.4	32.7	98
Intervention	Middle	70.5	28.4	4.5	27.3	10.2	29.5	88
area	Fourth	74.0	15.4	6.7	42.3	9.6	26.0	104
	Highest	78.2	7.7	0.7	66.9	2.8	21.8	142
	Total	69.4	18.6	4.7	36.0	10.1	30.6	555
	Lowest	45.4	7.4	7.4	3.7	26.9	54.6	108
	Second	52.8	13.9	2.8	11.1	25.0	47.2	72
Comparison	Middle	61.5	6.2	20.0	26.2	9.2	38.5	65
area	Fourth	69.9	8.6	6.5	40.9	14.0	30.1	93
	Highest	69.1	7.4	2.1	55.3	4.3	30.9	94
	Total	59.5	8.6	7.2	27.5	16.2	40.5	432
	Lowest	51.1	16.5	6.1	9.5	19.0	48.9	231
	Second	61.2	18.2	5.9	15.9	21.2	38.8	170
	Middle	66.7	19.0	11.1	26.8	9.8	33.3	153
Both areas	Fourth	72.1	12.2	6.6	41.6	11.7	27.9	197
	Highest	74.6	7.6	1.3	62.3	3.4	25.4	236
	Total	65.0	14.2	5.8	32.3	12.8	35.0	987

<sup>\*</sup>Multiple responses recorded

ANC = Antenatal care

FWV = Family welfare visitor

FDSR = Family Development Services and Research

CMH = Christian Memorial Hospital

HDSS = Health and Demographic Surveillance System

# Use of postnatal care services

It was observed that only 30.0% of the pregnant women received at least one postnatal care (PNC) during 2009. This percentage was higher in the intervention area (32.8%) than the comparison area (26.4%). The nurses, doctors and midwifes were the dominant sources for PNC in both the areas, and the utilization of services was characterized by large inequities (Table 17).

Table 17. F	Postnatal HDSS, 200		pe of sou	rces an	d asset q	<b>uintile</b>	, Chak	aria
Area	Asset	Received	Midwife*	FWV*	Nurse/	FDSR/	None	No. of
	quintile	any PNC (%)	(%)	(%)	doctor* (%)	CMH* (%)	(%)	women
	Lowest	17.9	4.9	0.0	11.4	1.6	82.1	123
	Second	22.4	5.1	0.0	15.3	2.0	77.6	98
Intervention	Middle	30.7	17.0	1.1	12.5	0.0	69.3	88
area	Fourth	46.2	5.8	1.9	37.5	1.0	53.8	104
	Highest	44.4	7.7	0.0	36.6	0.0	55.6	142
	Total	32.8	7.7	0.5	23.6	0.9	67.2	555
	Lowest	16.7	2.8	0.0	12.0	1.9	83.3	108
	Second	20.8	4.2	1.4	15.3	0.0	79.2	72
Comparison area	Middle	18.5	6.2	0.0	12.3	0.0	81.5	65
	Fourth	31.2	9.7	0.0	21.5	0.0	68.8	93
	Highest	42.6	10.6	0.0	30.9	1.1	57.4	94
	Total	26.4	6.7	0.2	18.8	0.7	73.6	432
	Lowest	17.3	3.9	0.0	11.7	1.7	82.7	231
	Second	21.8	4.7	0.6	15.3	1.2	78.2	170
	Middle	25.5	12.4	0.7	12.4	0.0	74.5	153
Both areas	Fourth	39.1	7.6	1.0	29.9	0.5	60.9	197
	Highest	43.6	8.9	0.0	34.3	0.4	56.4	236
	Total	30.0	7.3	0.4	21.5	0.8	70.0	987

\*Multiple responses recorded

PNC = Postnatal care

FWV = Family welfare visitor

FDSR = Family Development Services and Research

CMH = Christian Memorial Hospital

HDSS = Health and Demographic Surveillance System.

# Assistance during delivery

In Chakaria, the traditional birth attendants (TBAs) were more popular than the skilled birth attendants (SBAs) for assisting deliveries. 73.5 percent of 930 deliveries in Chakaria were assisted by the TBAs as opposed to 26.5 of the deliveries assisted by the SBAs (e.g. nurses/doctors, FWVs, midwives). The percentage of deliveries assisted by the TBAs was slightly higher in the comparison area (76.3%) than the intervention area (71.4%) (Table 18).

Despite the fact that the services provided by the midwives of the Chakaria project were also available to some parts of the comparison area, the use of these trained midwives was similar in the intervention area compared to the comparison area (10.7% vs. 11.1%) (Table 18). At the same time, the overall use of SBAs that comprised nurses, doctors, FWVs, and midwives was higher in the intervention area (28.6%) than the comparison area (23.7%) (Table 18). The use rate of nurse/doctors by the women from the highest quintile was much higher than those by women from the lowest quintiles.

Table 18. As	ssistance durin	ig delivery l	y asset qu	intile, Chal	karia HD	SS, 2009
Area	Asset quintile	Midwife	FWV	Nurse/ doctor	TBA	No. of women
		(%)	(%)	(%)	(%)	
	Lowest	5.2	0.9	6.1	87.8	115
	Second	9.0	1.1	5.6	84.3	89
Intervention	Middle	17.2	1.1	9.2	72.4	87
area	Fourth	12.4	3.1	15.5	69.1	97
	Highest	10.9	2.2	36.5	50.4	137
	Total	10.7	1.7	16.2	71.4	525
	Lowest	5.0	0.0	3.0	92.0	100
	Second	10.1	1.4	4.3	84.1	69
Comparison	Middle	8.3	1.7	8.3	81.7	60
area	Fourth	12.4	3.4	7.9	76.4	89
	Highest	19.5	1.1	31.0	48.3	87
	Total	11.1	1.5	11.1	76.3	405
Both areas	Lowest	5.1	0.5	4.7	89.8	215
	Second	9.5	1.3	5.1	84.2	158
	Middle	13.6	1.4	8.8	76.2	147
	Fourth	12.4	3.2	11.8	72.6	186
	Highest	14.3	1.8	34.4	49.6	224
	Total	10.9	1.6	14.0	73.5	930

FWV = Family Welfare Visitor

HDSS = Health and Demographic Surveillance System.

# Place of delivery

Eighty eight percent of the deliveries took place at home. Only 12.0% of 988 deliveries took place either at hospitals or at clinics. The percentage of deliveries taking place at the hospitals was higher in the intervention area (14.3%) compared to the comparison area (9.2%) (Table 19). The women from the households in the highest asset quintile had a much higher rate of facility based delivery than those from the lowest quintile.

Table 19. I	Tace of delivery	y by asset quintile, Cha	akaria HDSS,	2009
Area	Asset quintile	Hospital/Clinic (%)	Home (%)	No. of women
	Lowest	5.7	94.3	123
	Second	5.1	94.9	98
Intervention	Middle	8.0	92.0	88
area	Fourth	14.4	85.6	104
	Highest	31.9	68.1	141
	Total	14.3	85.7	554
	Lowest	3.6	96.4	110
	Second	4.2	95.8	72
Comparison	Middle	3.1	96.9	65
area	Fourth	7.5	92.5	93
	Highest	25.5	74.5	94
	Total	9.2	90.8	434
Both areas	Lowest	4.7	95.3	233
	Second	4.7	95.3	170
	Middle	5.9	94.1	153
	Fourth	11.2	88.8	197
	Highest	29.4	70.6	235
	Total	12.0	88.0	988

Table 20 shows caesarean-section delivery by household asset quintile in 2009. Caesarean-section delivery accounted for 4% of the deliveries in the Chakaria HDSS area in 2009. Although the number of caesarean sections was small, the number of women with caesarean sections from the highest quintile was 4 times the number of women from the lowest quintile.

Table 20. Proportion of caesarean-section delivery by asset quintile, Chakaria HDSS, 2009						
Asset quintile	No. of caesarean- section delivery	Caesarean-section delivery (%)	Total number of deliveries			
Lowest	5	2.1	233			
Second	1	0.6	170			
Middle	2	1.3	153			
Fourth	9	4.6	197			
Highest	19	8.0	237			
Total	36	3.6	990			

#### **Water and Sanitation Practices**

Tubewell was the universal source of drinking water in Chakaria. As a source it increased from 99.3% in 1994 to 99.9% in 2009. No variation persists in drinking water sources among the people belonging to the five asset quintiles (Table 21).

Table 21. Sources of drinking water by asset quintile, Chakaria HDSS, 1994 and 2009								
	Sources							
Asset - quintile	1994				2009			
	Tube	Well/Ditch/	No. of	Tube	Well/ Ditch/	No. of		
	Well	River/ Pond	Households	Well	River/ Pond	Households		
	%	(%)		%	(%)			
Lowest	99.7	0.3	369	100	0.0	336		
Second	99.7	0.3	367	100	0.0	336		
Medium	98.4	0.6	374	99.7	0.3	330		
Fourth	99.5	0.5	353	100	0.0	328		
Highest	99.2	0.8	365	100	0.0	339		
All	99.3	0.7	1,828	99.9	0.1	1,669		

The major source of water for bathing had shifted between 1994 to 2009. River, pond, ditch, and canal were the dominant sources of water for bathing (77.8%) in 1994 opposed to 37.9% in 2009. The sources varied by asset quintile both in 1994 and in 2009. Tubewell water was more popular in the highest quintile (Table 22).

Table 22. Sources of water for bathing by asset quintile, Chakaria HDSS, 1994 and 2009								
	Sources							
Asset -	1994				2009			
	Tube	Well/Ditch/	No. of	Tube	Well/Ditch/	No. of		
quintile	Well	River/Pond	Households	Well	River/Pond	Households		
	(%)	(%)		%	(%)			
Lowest	18.4	81.6	369	56.8	42.2	336		
Second	19.3	80.7	367	54.4	45.6	336		
Medium	17.4	82.6	374	59.4	39.6	330		
Fourth	22.9	77.1	353	64.3	35.7	328		
Highest	32.9	67.1	365	75.8	24.2	339		
All	22.2	77.8	1,828	62.1	37.9	1,669		

Table 2	3. Latri	ine use b	y asset	Table 23. Latrine use by asset quintile, Chakaria HDSS, 1994 and 2009	Chakari	a HDSS,	1994 a	nd 2009				
			15	1994					20	2009		
•			Fixed	Fixed place					Fixed	Fixed place		
Asset	oN fixed	Faeces drained	Faeces draine	Faeces	Faeces drained	No. of house	No fixed	Faeces	Faeces drained	Faeces	Faeces	No. of house-
quintile	place	into	d into	into	into	-holds	place	into	into	into	d into	holds
		surface	simple	concrete	septic		1	surface	simple	concrete	septic	
		water	pit	pit	tank			water	pit	pit	tank	
		bodies	latrine	latrine				bodies	latrine	latrine		
	%	%	%	%	%		%	%	%	%	%	
Lowest	35.0	26.3	34.4	4.3	0.0	369	9.2	20.2	31.6	38.4	9.0	336
Second	25.3	30.2	33.2	11.2	0.0	367	4.5	19.1	33.3	42.9	0.3	336
Medium	26.7	25.9	39.0	8.3	0.0	374	5.5	16.1	27.9	47.6	3.0	330
Fourth	18.1	24.9	38.0	17.3	1.7	353	3.1	22.6	24.7	45.1	4.6	328
Highest	0.9	18.4	30.4	34.0	11.2	365	2.4	13.9	16.8	49.0	18.0	339
All	22.3	25.2	35.0	14.9	2.6	1,828	4.9	18.3	26.8	44.6	5.3	1,669

2009, there was a substantial improvement in all the households irrespective of asset quintiles. Also at the Table 23 presents the percentage of households using the various types of latrine by asset quintile in 1994 and in 2009. The use of sanitary latrine increased from 17.5% in 1994 to 49.9% in 2009 in Chakaria HDSS area. Although, a variation in using sanitary latrine between the five asset quintiles persists in 1994 and same time, the open place for defecation decreased significantly, from 22.3% in 1994 to 4.9% in 2009.

Note: In this chapter, to compare the water and sanitation practices between 1994 and 2009, the asset quintile measurement tool was applied, which was derived from a common list of assets including radio, television, bicycle, motor cycle, land ownership, occupation and education of household head, which were available in both years.

#### **AUTHORS' COMMENTS**

The report revealed that the rates for crude birth and total fertility have decreased more than expected, between 2008 and 2009. In trying to understand the factors associated with this, we examined the midyear population, number of women of reproductive age, and number of births for both years. We found the numerator (number of birth) to be the main factor for lower rates in 2009. Moreover, when checking the number of births by villages for 2008 and 2009, a discrepancy was found in the number of births in 12 villages. In these 12 villages, there were 56 births in 2009, compared to 110 births in 2008. To understand this drastic decrease we re-checked all of the conception records from those villages for that period. However, no inconsistencies were found in that data. Furthermore, we also compared the number of pregnant women for the entire DSS area for 2008 and 2009. Here we found that the number of pregnant women had decreased in 2009 from 2008. We also found that the rate of reported spontaneous abortion had increased from 8.8% in 2008 to 12.5% in 2009.

Simultaneously, we also noticed an increase in neonatal, post-neonatal and infant mortality rates in 2009, which is dependent on the denominator (number of births).

We think that the low number of conception and high rate of spontaneous abortion, contributed towards decreased birth numbers in 2009 than in 2008.

Receiving assistance from SBA decreased in the intervention area significantly. One of the reasons may be the cessation of the SBA voucher programme in May 2009.

#### **REFERENCES**

- 1. Bhuiya A, Ribaux C, and Eppler P. Community-led primary healthcare initiatives: Lessons learned from a project in rural Bangladesh. In J Rohde and J Wyon (eds.) Community-Based Health Care: Lessons from Bangladesh to Boston. Boston: Management Sciences for Health. 2002. pp 87-111.
- 2. Eppler P, Bhuiya A, and Hossain M. A Process-oriented approach to the establishment of Community-based Village Health Posts. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 1996. 37p.
- 3. Bhuiya A, Hanifi SMA, and Mahmood SS. Chakaria health and demographic surveillance system: focusing on the poor and vulnerable. Socioeconomic, Health and Demographic Profile, 1999–2000. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 2006. 56p. (Scientific report no. 94).
- 4. Filmer D, Pritchett LH. Estimating wealth effects without expenditure dataor tears: an application to educational enrollments in states of India. Demography. 2001;38(1):115-32.
- 5. International Centre for Diarrhoeal Disease Research, Bangladesh. Health and Demographic Surveillance System-Matlab: Registration of health and demographic events 2008. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 2010. 71 p. (Scientific report no. 109).

#### **ADDITIONAL READINGS**

Bhuiya A. Health for the Rural Masses: Insights from Chakaria. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 2009.

Bhuiya A, Hanifi SMA, and Urni F., Mahmood SS. Three methods to monitor utilization of healthcare services by the poor.International Journal for Equity in Health 2009, 8:29

Bhuiya A, Hanifi SMA, and Urni F. Chakaria health and demographic surveillance system: focusing on the poor and vulnerable. Demographic Events and SAFE Motherhood Practices - 2008. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 2009. 52 p. (Scientific report no. 108).

Bhuiya A, Hanifi SMA, and Urni F., Iqbal M. Chakaria health and demographic surveillance system: focusing on the poor and vulnerable. Demographic Events and Safe Motherhood Practices - 2007. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 2008. 47 p. (Scientific report no. 105).

Bhuiya A, Hanifi SMA, and Urni F. Chakaria health and demographic surveillance system: focusing on the poor and vulnerable. Socioeconomic, Health and Demographic Profile and Utilization of Healthcare Services - 2006. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 2008. 51 p. (Scientific report no. 102).

Bhuiya A., Hanifi SMA, Mahmood SS. Chakaria health and demographic surveillance system: focusing on the poor and vulnerable. Demographic profile, family-planning use, and safe motherhood practices, 2005. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 2007. 50 p. (Scientific report no. 100).

Bhuiya A, Hanifi SMA, Chowdhury M, Jahangir M, and Gwatkin DR. Rapid methods for monitoring the utilization of healthcare facilities by the poor: Findings from a pilot project in rural Bangladesh. FHS Working paper 2, Bangladesh series, October 2007. http://www.futurehealthsystems.org/publications/WP%202%20final.pdf.

Bhuiya A, Hanifi SMA, Roy N, Streatfield PK. Performance of the Lot Quality Assurance Sampling Method Compared to surveillance for Identifying Inadequately- performing Areas in Matlab, Bangladesh. J Health Popul Nutr 2007 March; 25(1):37-46.

Bhuiya A, Aziz A, Hanifi SMA. Reproductive and sexual health problems as perceived by women and men in a rural area of Bangladesh. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh; 1997. 24 p.

Bhuiya A, Hanifi SMA, Hossain M, Aziz A. Effects of an AIDS awareness campaign about AIDS in a remote rural area of Bangladesh. Int Q Community Health Educ 2000; 19(1): 51-63.

Bhuiya A, Ribaux C, Eppler P. Community-led primary healthcare initiatives: lessons learned from a project in rural Bangladesh. In: Rohde J, Wyon J, editors. Community-based health care: Lessons from Bangladesh to Boston. Boston: Management Sciences for Health, 2002: 87-111.

Bhuiya A, Ribaux CA. Rethinking community participation: prospects of health initiatives by indigenous self-help organizations in rural Bangladesh. Dhaka: International Centre for Diarrhoeal Disease Research Bangladesh, 1997. (Special publication no. 65). 31 p.

Bhuiya A, Sharmin T, Hanifi SMA. Nature of domestic violence against women in a rural area of Bangladesh: Implication for preventive interventions. J Health Popul Nutr 2003; 21(1): 48-54.

Bhuiya A, Yasmin F, Begum F, Rob U. Community participation in health, family planning and development programmes. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 1997. 53 p.

Bhuiya A. Health knowledge and behaviour in five unions of Chakaria. Dhaka: International Centre for Diarrhoeal Disease Research, Bangladesh, 1996. (Special publication no. 52). 38 p.

Bhuiya A., Mahmood SS., Rana AKMM, Ahmed SM, Chowdhury AMR. A multidimensional Approach to Measure Poverty in Rural Bangladesh. J Health Popul Nutr 2007;25(2):134-145.

Choudhury KK, Hanifi SMA, Mahmood SS, Bhuiya A. Sociodemographic Characteristics of Tobacco Consumers in a Rural Area of Bangladesh. J Health Popul Nutr 2007;25(4):456-64.

Choudhury KK, Hanifi SMA, Rasheed S, Bhuiya A. Gender inequality and severe malnutrition among children in a remote rural area of Bangladesh. J Health Popul Nutr 2000;18(3):123-30.

Fariba Alamgir, Papreen Nahar, Andrew E. Collins, Dr. Nibedita S. Ray-Bennett, Abbas Bhuiya 'Climate Change and Food Security: Health Risks and Vulnerabilities of the Poor in Bangladesh' 2010; The International Journal of Climate Change: Impacts and Responses, Volume 1, Issue 4, pp.37-54.

Hanifi SMA, Haq MZ, Aziz RR, and Bhuiya A. High concentration of childhood deaths in the low-lying areas of Chakaria HDSS, Bangladesh: Findings from a spatial analysis. Global Health Action Supplement 1, 2010, 70-76

Hanifi SMA, Mahmood SS, and Bhuiya A. Smoking has declined but not for all: Findings from a study in a rural area of Bangladesh. Asia-Pacific Journal of Public Health. 2010 May 24:1-10

Hanifi SMA, Bhuiya A. Family-planning services in a low-performing rural area of Bangladesh: insights from field observations. J Health Popl Nutr 2001;19(3):209-14.

Hossain SMM, Bhuiya A, Khan AR, Uhaa I. Community development and its impact on health: South Asian experience. BMJ 2004; 328:830-3.

Hossain SMM, Bhuiya A, Rasheed S. Correlates of perceived malarial episodes and treatment-seeking behavior in a malarial-endemic rural area in Bangladesh. Southeast Asian J Trop Med Public Health 2001;32(4):707-19.

Iqbal M, Rasheed S, Hanifi SMA, Bhuiya A. 2009. Reaching the poor with performance based payment for safe delivery services in rural Bangladesh. Medicus Mundi Bulletin 112:37-42

Jacob S. Siegel and David A. Swanson. The Methods and Materials of Demography, Second edition. Elsevier Academic Press, 2004. pp 196-340.

Nahar, P., F. Alamgir, et al. (2010). Contextualizing disaster in relation to human health in Bangladesh. Asian journal of water environment and Pollution 7(1): 55-62.

Rasheed S, Hanifi SMA, Iqbal M, NN, Bhuiya A. Policy of universal salt iodization in Bangladesh: do coastal people benefit? J Health Popul Nutr 2001;19(2):66-72.

#### APPENDIX A

# Midyear population by age and sex in the intervention and comparison areas, Chakaria, HDSS, 2009

Age	Inte	rvention a	area	Con	nparison	area	I	Both areas	5
(years)	Male	Female	Both	Male	Female	Both	Male	Female	Both
<1	287	269	556	224	236	460	511	505	1,016
1-4	1,288	1,222	2,510	1,017	918	1,935	2,305	2,140	4,445
5-9	1,684	1,569	3,253	1,288	1,238	2,526	2,972	2,807	5,779
10-14	1,746	1,706	3,452	1,500	1,400	2,900	3,246	3,106	6,352
15-19	1,658	1,698	3,356	1,386	1,525	2,911	3,044	3,223	6,267
20-24	1,268	1,206	2,474	1,118	999	2,117	2,386	2,205	4,591
25-29	894	736	1,630	747	598	1,345	1,641	1,334	2,975
30-34	607	706	1,313	492	554	1,046	1,099	1,260	2,359
35-39	602	599	1,201	463	560	1,023	1,065	1,159	2,224
40-44	528	533	1,061	397	437	834	925	970	1,895
45-49	487	456	943	446	431	877	933	887	1,820
50-54	379	398	777	323	377	700	702	775	1,477
55-59	328	290	618	348	301	649	676	591	1,267
60-64	275	205	480	205	144	349	480	349	829
65-69	181	153	334	183	154	337	364	307	671
70-74	135	101	236	108	82	190	243	183	426
75-79	123	97	220	76	73	149	199	170	369
80-84	52	44	96	37	31	68	89	75	164
85+	60	35	95	39	38	77	99	73	172
All	12,582	12,023	24,605	10,397	10,096	20,493	22,979	22,119	45,098

#### APPENDIX B

#### Percentage distribution of midyear population by age and sex in the intervention and comparison areas, Chakaria HDSS, 2009

Age	Inte	rvention a	rea	Co	mparison a	area	I	Both areas	
(years)	Male	Female	Both	Male	Female	Both	Male	Female	Both
<1	2.3	2.2	2.3	2.2	2.3	2.2	2.2	2.3	2.3
1-4	10.2	10.2	10.2	9.8	9.1	9.4	10.0	9.7	9.9
5-9	13.4	13.0	13.2	12.4	12.3	12.3	12.9	12.7	12.8
10-14	13.9	14.2	14.0	14.4	13.9	14.1	14.1	14.0	14.1
15-19	13.2	14.1	13.6	13.3	15.1	14.2	13.2	14.6	13.9
20-24	10.1	10.0	10.1	10.8	9.9	10.3	10.4	10.0	10.2
25-29	7.1	6.1	6.6	7.2	5.9	6.6	7.1	6.0	6.6
30-34	4.8	5.9	5.3	4.7	5.5	5.1	4.8	5.7	5.2
35-39	4.8	5.0	4.9	4.5	5.5	5.0	4.6	5.2	4.9
40-44	4.2	4.4	4.3	3.8	4.3	4.1	4.0	4.4	4.2
45-49	3.9	3.8	3.8	4.3	4.3	4.3	4.1	4.0	4.0
50-54	3.0	3.3	3.2	3.1	3.7	3.4	3.1	3.5	3.3
55-59	2.6	2.4	2.5	3.3	3.0	3.2	2.9	2.7	2.8
60-64	2.2	1.7	2.0	2.0	1.4	1.7	2.1	1.6	1.8
65-69	1.4	1.3	1.4	1.8	1.5	1.6	1.6	1.4	1.5
70-74	1.1	0.8	1.0	1.0	0.8	0.9	1.1	0.8	0.9
75-79	1.0	0.8	0.9	0.7	0.7	0.7	0.9	0.8	0.8
80-84	0.4	0.4	0.4	0.4	0.3	0.3	0.4	0.3	0.4
85+	0.5	0.3	0.4	0.4	0.4	0.4	0.4	0.3	0.4
All	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

### APPENDIX C

# Number of births by age of mother, Chakaria HDSS, 2009

Age (years)	Inte	rvention a	irea	Cor	nparison a	irea		Both areas	3
Age (years)	Male	Female	Both	Male	Female	Both	Male	Female	Both
10-14	0	3	3	0	0	0	0	3	3
15-19	81	66	147	72	62	134	153	128	281
20-24	109	109	218	70	78	148	179	187	366
25-29	46	54	100	48	40	88	94	94	188
30-34	48	27	75	34	21	55	82	48	130
35-39	12	18	30	8	12	20	20	30	50
40-44	3	6	9	0	3	3	3	9	12
45-49	1	0	1	0	0	0	1	0	1
50-54	0	1	1	0	0	0	0	1	1
All	300	284	584	232	216	448	532	500	1,032

#### APPENDIX D

# Number of deaths by age and sex, Chakaria HDSS, 2009

Age	Inte	rvention a	rea	Con	nparison a	ırea	]	Both areas	
(years)	Male	Female	Both	Male	Female	Both	Male	Female	Both
<1	20	12	32	12	16	28	32	28	60
1-4	6	7	13	2	6	8	8	13	21
5-9	1	2	3	1	1	2	2	3	5
10-14	2	1	3	1	0	1	3	1	4
15-19	2	3	5	1	3	4	3	6	9
20-24	4	2	6	1	3	4	5	5	10
25-29	0	1	1	1	0	1	1	1	2
30-34	4	0	4	0	0	0	4	0	4
35-39	0	2	2	0	1	1	0	3	3
40-44	1	3	4	1	1	2	2	4	6
45-49	8	4	12	3	2	5	11	6	17
50-54	5	2	7	2	3	5	7	5	12
55-59	5	2	7	6	5	11	12	7	19
60-64	5	4	10	2	2	4	7	6	13
65-69	7	7	14	6	5	11	13	12	25
70-74	2	3	5	3	7	10	5	10	15
75-79	6	5	11	6	5	11	12	10	22
80-84	6	2	8	3	1	4	9	3	12
85+	9	11	20	4	9	13	13	20	33
All	94	73	167	55	70	125	149	143	292

### APPENDIX E

# Causes of deaths by age and sex, Chakaria HDSS, 2009

Cause	All			Age (	years)		
Cause	age	<1	1-4	5-14	15-49	50-59	60+
Male							
Communicable diseases							
Diarrheal	3	1	0	0	2	0	0
Dysentery	1	1	0	0	0	0	0
Tuberculosis	1	0	0	0	1	0	0
Hepatitis	7	0	0	1	3	1	2
Respiratory infections	5	5	0	0	0	0	0
Maternal and neonatal conditions							
Neonatal	9	8	0	0	0	0	1
Other neonatal	9	9	0	0	0	0	0
Non–communicable diseases							
Malignant neoplasm	7	0	0	0	2	1	4
Neoplasm	11	1	0	1	3	0	6
Diabetes	5	0	0	0	0	2	3
Stroke	21	0	0	0	5	5	11
Other cardiovascular	4	1	0	0	0	2	1
Asthma/Bronchitis	17	0	0	0	1	2	14
Digestive disease	1	0	0	0	1	0	0
Senility	11	0	0	0	0	0	11
Epilepsy	2	0	1	0	1	0	0
Injuries							
Accident	6	0	1	1	3	1	0
Drowning	7	1	5	1	0	0	0
Homicide	1	0	0	0	1	0	0
Suicide	1	0	0	1	0	0	0
Unknown	20	5	1	0	3	5	6
All	149	32	8	5	26	19	59
Female							
Communicable diseases							
Diarrheal	4	0	1	0	3	0	0
Hepatitis	2	1	0	0	0	0	1
Respiratory infection	17	16	1	0	0	0	0
Malaria	1	0	0	0	1	0	0
Maternal and neonatal conditions							
Maternal death	1	0	0	0	1	0	0
Premature and lbw	1	1	0	0	0	0	0
Other neonatal	7	7	0	0	0	0	0

# Appendix E: (Contd...)

Cause	All			Age	(years)		
Cause	age	<1	1-4	5-14	15-49	50-59	60+
Nutritional	2	0	1	0	1	0	0
Non–communicable diseases							
Malignant neoplasm	6	0	0	0	1	4	1
Neoplasm	5	0	0	0	2	0	3
Diabetes	2	0	0	0	0	1	1
Hypertension disease	2	0	0	0	0	0	2
Stroke	20	0	0	0	5	3	12
Other cardiovascular	1	0	0	0	1	0	0
Asthma/Bronchitis	14	0	0	0	3	2	9
Other urinary	2	0	0	0	1	0	1
Epestaxis	1	1	0	0	0	0	0
Senility	26	0	0	0	0	0	26
Disease of uterus	1	0	0	0	0	1	0
Injuries							
Accident	2	0	2	0	0	0	0
Drowning	7	0	5	2	0	0	0
Suicide	2	0	0	0	2	0	0
Burn	1	0	1	0	0	O	0
Unknown	16	2	2	2	4	1	5
All	143	28	13	4	25	12	61

## APPENDIX F

## Number of migrants by age and sex, Chakaria HDSS, 2009

Age	Inte	rvention a	rea	Cor	nparison a	rea		Both areas	}
(years)	Male	Female	Both	Male	Female	Both	Male	Female	Both
In-migra	nts								
<1	13	10	23	7	8	15	20	18	38
1-4	27	37	64	12	30	42	39	67	106
5-9	36	36	72	32	25	57	68	61	129
10-14	63	44	107	28	39	67	91	83	174
15-19	46	185	231	34	142	176	80	327	407
20-24	34	68	102	20	55	75	54	123	177
25-29	33	27	60	20	20	40	53	47	100
30-34	16	15	31	6	4	10	22	19	41
35-39	14	3	17	9	3	12	23	6	29
40-44	9	1	10	3	5	8	12	6	18
45-49	3	1	4	5	2	7	8	3	11
50-54	4	7	11	3	5	8	4	7	11
55-59	3	10	13	1	4	5	6	15	21
60-64	4	8	12	0	7	7	5	12	17
65-69	4	6	10	1	8	9	4	13	17
70-74	2	4	6	2	6	8	3	12	15
75-79	4	3	7	3	1	4	6	9	15
80-84	2	1	3	3	2	5	5	2	7
85+	1	3	4	7	8	15	4	5	9
All	318	469	787	189	366	555	507	835	1,342
Out-mig									
<1	13	15	28	20	11	31	33	26	59
1-4	36	23	59	43	30	73	79	53	132
5-9	24	36	60	23	36	59	47	72	119
10-14	68	44	112	41	41	82	109	85	194
15-19	72	169	241	68	163	231	140	332	472
20-24	54	163	217	70	128	198	124	291	415
25-29	58	47	105	54	37	91	112	84	196
30-34	22	19	41	22	11	33	44	30	74
35-39	21	5	26	18	5	23	39	10	49
40-44	6	4	10	8	2	10	14	6	20
45-49	4	1	5	3	3	6	7	4	11
50-54	3	5	8	2	3	5	5	8	13
55-59	2	8	10	1	6	7	3	14	17
60-64	5	5	10	1	2	3	6	7	13
65-69	2	2	4	2	6	8	4	8	12
70-74	4	2	6	2	4	6	6	6	12
75-79	2	2	4	2	5	7	4	7	11
80-84	2	1	3	2	2	4	4	3	7
85+	3	2	5	0	1	970	3	3	1 022
All	401	553	954	382	497	879	783	1,049	1,832

#### APPENDIX G

#### Migration rate per 1,000 population by age and sex, Chakaria HDSS, 2009

Age	Inte	rvention a	rea	Cor	nparison a	irea		Both areas	
(years)	Male	Female	Both	Male	Female	Both	Male	Female	Both
In-migra	tion								
<1	45.3	37.2	41.4	31.3	33.9	32.6	39.1	35.6	37.4
1-4	21.0	30.3	25.5	11.8	32.7	21.7	16.9	31.3	23.9
5-9	21.4	22.9	22.1	24.8	20.2	22.6	22.9	21.7	22.3
10-14	36.1	25.8	31.0	18.7	27.8	23.1	28.0	26.7	27.4
15-19	27.8	108.8	68.8	24.5	93.2	60.5	26.3	101.4	65.0
20-24	26.7	56.4	41.2	17.9	54.9	35.4	22.6	55.7	38.5
25-29	37.0	36.8	36.9	26.8	33.4	29.7	32.3	35.3	33.6
30-34	26.3	21.2	23.6	12.2	7.2	9.6	20.0	15.1	17.4
35-39	23.3	5.0	14.2	19.5	5.4	11.8	21.6	5.2	13.1
40-44	17.0	1.9	9.4	7.5	11.4	9.6	12.9	6.2	9.5
45-49	6.2	2.2	4.2	11.2	4.6	8.0	8.6	3.4	6.0
50-54	10.6	17.5	14.2	9.3	13.3	11.4	5.7	9.0	7.4
55-59	9.1	34.5	21.0	2.9	13.3	7.7	8.9	25.4	16.6
60-64	14.5	38.8	24.9	0.0	48.3	20.0	10.4	34.2	20.4
65-69	22.1	39.5	30.0	5.5	51.9	26.7	11.0	42.5	25.4
70-74	14.9	39.2	25.4	18.5	73.2	42.1	12.4	65.2	35.2
75-79	32.5	30.9	31.8	39.5	13.7	26.8	30.2	52.9	40.7
80-84	38.5	22.7	31.3	81.1	62.5	72.5	56.2	26.3	42.4
85+	16.4	85.7	41.7	0.0	0.0	0.0	40.0	68.5	52.0
All	25.3	39.0	32.0	18.2	36.2	27.1	22.1	37.7	29.8
Out-mig	ration								
<1	45.3	55.8	50.4	89.3	46.6	67.4	64.6	51.5	58.1
1-4	28.0	18.8	23.5	42.3	12.0	16.0	34.3	24.8	29.7
5-9	14.2	22.9	18.4	33.4	24.3	28.9	15.8	25.7	20.6
10-14	38.9	25.8	32.5	15.3	25.7	20.3	33.6	27.4	30.5
15-19	43.5	99.4	71.8	29.6	26.9	28.2	46.0	103.0	75.3
20-24	42.5	135.2	87.6	60.8	162.8	109.0	51.9	131.9	90.3
25-29	64.9	64.0	64.5	93.7	214.0	147.2	68.3	63.1	65.9
30-34	36.2	26.8	31.2	109.5	66.8	86.9	40.0	23.8	31.3
35-39	34.9	8.4	21.7	47.6	19.7	32.3	36.7	8.7	22.1
40-44	11.3	7.5	9.4	45.1	11.4	27.5	15.1	6.2	10.5
45-49	8.2	2.2	5.3	17.9	4.6	11.4	7.5	4.5	6.0
50-54	7.9	12.5	10.3	9.3	8.0	8.6	7.1	10.3	8.8
55-59	6.1	27.6	16.2	5.7	10.0	7.7	4.4	23.7	13.4
60-64	18.1	24.3	20.7	4.9	41.4	20.0	12.5	19.9	15.6
65-69	11.0	13.2	12.0	5.5	13.0	8.9	11.0	26.1	17.9
70-74	29.9	19.6	25.4	18.5	73.2	42.1	24.8	32.6	28.2
75-79	16.3	20.6	18.2	26.3	54.8	40.3	20.1	41.2	29.8
80-84	38.5	22.7	31.3	54.1	156.3	101.4	44.9	39.5	42.4
85+	49.2	57.1	52.1	51.3	52.6	51.9	30.0	41.1	34.7
All	31.9	46.0	38.8	36.7	49.2	42.9	34.1	47.4	40.6

### APPENDIX H

#### Number of migrants by origin or destination, Chakaria HDSS, 2009

Origin/	All						Age (yea	ars)				
Destination	age	<5	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50+
In-migration												
Male												
Inside Bangladesh	446	61	64	83	83	44	36	18	13	6	5	33
Outside Bangladesh	61	0	2	1	0	11	18	6	10	5	3	5
Inside Chakaria	327	47	45	71	65	26	21	9	7	3	2	31
Outside Chakaria	85	11	16	10	14	12	8	5	5	2	1	1
Inside HDSS area	214	29	27	47	41	16	13	8	4	2	2	25
Outside HDSS area	153	25	27	27	30	18	10	4	3	2	1	6
Female												
Inside Bangladesh	831	82	56	84	311	138	50	18	8	4	5	75
Outside Bangladesh	4	2	0	0	1	0	1	0	0	0	0	0
Inside Chakaria	583	52	45	61	208	88	32	13	7	2	3	72
Outside Chakaria	208	26	8	13	93	42	15	5	1	1	2	2
Inside HDSS area	386	34	28	51	128	47	21	10	6	2	3	56
Outside HDSS area	328	31	22	19	139	66	23	6	2	1	2	17
Out-migration												
Male												
Inside Bangladesh	746	138	55	122	126	81	87	32	29	19	7	50
Outside Bangladesh	167	2	0	2	24	59	42	17	16	2	2	1
Inside Chakaria	366	58	30	80	73	41	28	11	10	3	3	29
Outside Chakaria	127	28	9	15	24	15	14	6	5	4	2	3
Inside HDSS area	249	40	20	48	59	31	18	5	5	1	2	20
Outside HDSS area	172	29	14	39	23	19	16	9	8	3	1	11
Female												
Inside Bangladesh	1,193	102	84	100	359	314	100	33	14	8	5	74
Outside Bangladesh	9	1	0	0	1	3	1	1	0	1	0	1
Inside Chakaria	641	51	46	50	207	168	44	13	5	5	3	49
Outside Chakaria	236	14	12	18	78	75	25	7	4	5	1	2
Inside HDSS area	420	32	30	32	134	108	28	5	4	3	2	42
Outside HDSS area	340	23	21	24	113	103	30	14	3	2	2	7

#### APPENDIX I

#### Number of in-migrants by reasons for migration, Chakaria HDSS, 2009

Reason for migration	All					A	Age (yea	ars)				
icason for inigration	age	<5	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50+
Male												
Family-related To join spouse Family friction/ breakdown Others	11 31 20	0 3 3	0 3 2	0 1 1	1 3 4	4 4 3	3 6 0	2 2 0	0 1 0	1 0 0	0 0 1	0 8 6
Work-related New job/job transfer To look for work/lost job others	98 156 0	0 1 0	2 11 0	7 54 0	19 43 0	13 18 0	23 6 0	9 8 0	11 4 0	6 3 0	5 2 0	3 6 0
Housing-related Wanted to own home/ new house	131	49	22	13	7	10	11	0	5	1	0	13
Education To acquire education	46	1	28	15	1	1	0	0	0	0	0	0
Reasons not reported	14	2	0	0	2	1	4	1	2	1	0	1
All	507	59	68	91	80	54	53	22	23	12	8	37
Female												
Family related To join spouse Family friction/breakdown Others	336 84 72	0 7 6	0 0 2	16 2 2	249 15 7	50 29 9	9 8 5	6 5 4	3 0 1	2 1 1	0 0 1	1 17 34
Work-related New job/job transfer To look for work/lost job	6 70	2 0	0 10	0 28	1 17	1 7	2 7	0	0	0	0	0
Others  Housing-related Wanted to own home/new house	213	67	25	19	31	26	15	3	2	0	2	23
Education To acquire education	51	3	24	16	6	0	1	1	0	0	0	0
Reasons not reported	3	0	0	0	1	1	0	0	0	1	0	0
All	835	85	61	83	327	123	47	19	6	6	3	75

## APPENDIX J

#### Number of out-migrants by reasons for migration, Chakaria HDSS, 2009

Reason for migration	All					:	Age (ye	ars)				
Reason for inigration	age	<5	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50+
Male												
Family-related												
To Join spouse	16	0	0	1	1	6	4	1	0	0	0	3
Family friction/	42	2	0	3	6	10	10	3	3	0	0	5
breakdown						_		_	_	_		_
Others	61	17	8	8	3	6	8	1	1	1	1	7
Work-related												
New job/job transfer	210	3	1	7	34	66	52	20	19	5	3	0
To look for work/ lost job	161	0	3	41	53	16	22	10	8	4	2	2
Others	0	0	0	0	0	0	0	0	0	0	0	0
	o o	O	Ü	· ·	O	O	Ü	Ü	Ü	Ü	Ü	O
Housing-related Wanted to own home/	217	89	24	25	22	9	14	8	7	4	0	15
new house	217	0)	21	23	22		17	O	,	T	O	13
Education												
To acquire education	53	1	11	22	17	2	0	0	0	0	0	0
Reasons not reported	21	0	0	2	3	8	2	1	1	0	1	3
All	783	112	47	109	140	124	112	44	39	14	7	35
Female												
Family-related												
To Join spouse	425	0	0	7	214	169	28	5	1	0	0	1
Family friction/ breakdown	64	2	2	1	15	19	6	0	1	0	1	17
Others	115	17	11	5	14	28	15	5	5	0	1	14
	110			3	- 1	_3	10	3	3	J	-	- 1
Work-related New job/job transfer	13	1	0	0	1	3	2	3	0	2	0	1
To look for work/lost	87	1	7	27	28	12	8	2	1	1	0	0
job												
Others	0	0	0	0	0	0	0	0	0	0	0	0
Housing-related												
Wanted to own home/ new house	293	59	35	29	49	55	23	14	2	3	2	22
Education												
To acquire education	42	0	17	16	7	0	1	1	0	0	0	0
Reasons not reported	11	0	0	0	4	5	1	0	0	0	0	1
All	1,050	80	72	85	332	291	84	30	10	6	4	56

#### APPENDIX K

#### Percentage of population by age and marital status, Chakaria HDSS, 2009

Age (years)	Married	Divorced	Abandoned	Widower/ Widow	Separated	Never married	Population
Male							
10-14	0.1	0.0	0.0	0.0	0.0	99.9	3,245
15-19	3.7	0.1	0.0	0.0	0.0	96.1	3,049
20-24	25.7	0.4	0.0	0.1	0.0	73.8	2,411
25-29	56.1	1.3	0.0	0.1	0.0	42.5	1,655
30-34	82.4	0.7	0.0	0.1	0.1	16.6	1,110
35-39	95.1	0.5	0.0	0.6	0.0	3.8	1,071
40-44	97.7	0.6	0.0	0.1	0.1	1.4	932
45-49	98.8	0.1	0.1	0.7	0.1	0.1	937
50-54	98.9	0.3	0.0	0.4	0.0	0.4	703
55-59	97.8	0.1	0.0	1.9	0.0	0.1	679
60-64	97.1	0.4	0.2	2.3	0.0	0.0	484
65-69	94.8	0.3	0.0	5.0	0.0	0.0	363
70-74	93.9	0.0	0.4	5.7	0.0	0.0	244
75-79	87.5	0.5	0.0	11.0	0.0	1.0	200
80-84	79.8	1.1	0.0	19.1	0.0	0.0	89
85+	73.0	0.0	0.0	27.0	0.0	0.0	100
All	47.2	0.4	0.0	0.8	0.0	51.5	17,272
Female							
10-14	0.9	0.0	0.0	0.0	0.0	99.1	3,106
15-19	31.5	0.2	0.0	0.1	0.0	68.1	3,235
20-24	69.8	1.6	0.4	0.6	0.0	27.5	2,225
25-29	87.9	1.4	1.2	1.7	0.0	7.8	1,338
30-34	92.6	1.2	0.9	2.8	0.2	2.1	1,265
35-39	89.9	1.7	2.2	6.0	0.1	0.2	1,156
40-44	87.3	1.2	1.4	9.3	0.2	0.4	971
45-49	80.6	1.0	1.8	16.3	0.0	0.1	888
50-54	71.1	0.6	1.9	25.8	0.1	0.3	778
55-59	60.1	1.2	1.0	36.7	0.2	0.3	592
60-64	48.7	0.0	0.6	50.7	0.0	0.0	351
65-69	39.5	0.3	1.6	57.8	0.3	0.3	306
70-74	27.7	1.6	0.0	70.7	0.0	0.0	184
75-79	15.9	0.6	0.6	82.4	0.0	0.6	170
80-84	9.2	0.0	0.0	90.8	0.0	0.0	76
85+	5.5	0.0	1.4	91.8	0.0	1.4	73
All	52.9	0.8	0.7	9.3	0.0	36.1	16,714

#### APPENDIX L

# Percentage of population by age and marital status, intervention area, Chakaria HDSS, 2009

Age (years)	Married	Divorced	Abandoned	Widower/ Widow	Separated	Never married	Population
Male							
10-14	0.1	0.0	0.0	0.0	0.0	99.8	1,746
15-19	3.6	0.1	0.0	0.0	0.0	96.3	1,658
20-24	24.4	0.3	0.0	0.1	0.0	75.2	1,283
25-29	54.6	1.6	0.0	0.2	0.0	43.6	901
30-34	81.5	1.0	0.0	0.2	0.2	17.2	615
35-39	95.4	0.7	0.0	0.3	0.0	3.6	605
40-44	97.6	0.6	0.0	0.2	0.0	1.7	532
45-49	99.0	0.2	0.0	0.8	0.0	0.0	490
50-54	98.7	0.3	0.0	0.5	0.0	0.5	379
55-59	97.9	0.3	0.0	1.5	0.0	0.3	330
60-64	97.1	0.0	0.4	2.5	0.0	0.0	276
65-69	94.5	0.0	0.0	5.5	0.0	0.0	181
70-74	96.3	0.0	0.7	3.0	0.0	0.0	135
75-79	89.5	0.8	0.0	8.1	0.0	1.6	124
80-84	76.9	1.9	0.0	21.2	0.0	0.0	52
85+	65.6	0.0	0.0	34.4	0.0	0.0	61
All	47.0	0.4	0.0	0.9	0.0	51.7	9,368
Female							
10-14	1.1	0.0	0.0	0.0	0.0	98.9	1,705
15-19	31.1	0.2	0.0	0.1	0.0	68.4	1,704
20-24	70.6	2.0	0.2	0.7	0.0	26.4	1,217
25-29	87.9	1.1	1.4	1.6	0.0	8.0	735
30-34	93.5	1.1	0.7	2.8	0.0	1.7	710
35-39	90.5	2.0	2.2	5.0	0.2	0.2	597
40-44	88.4	0.9	1.7	8.3	0.4	0.4	533
45-49	84.5	1.1	1.1	13.3	0.0	0.0	457
50-54	69.0	0.5	2.3	27.8	0.3	0.3	400
55-59	59.5	1.7	1.7	36.4	0.0	0.3	291
60-64	52.4	0.0	0.5	47.1	0.0	0.0	206
65-69	38.2	0.7	3.3	57.9	0.0	0.0	152
70-74	30.4	2.0	0.0	67.6	0.0	0.0	102
75-79	14.4	1.0	1.0	83.5	0.0	0.0	97
80-84	13.6	0.0	0.0	86.4	0.0	0.0	44
85+	2.9	0.0	0.0	94.3	0.0	2.9	35
All	53.2	0.9	0.7	8.9	0.0	36.2	8,985

#### APPENDIX M

# Percentage of population by age and marital status, comparison area, Chakaria HDSS, 2009

Age (years)	Married	Divorced	Abandoned	Widower/ Widow	Separated	Never married	Population
Male							
10-14	0.1	0.0	0.0	0.0	0.0	99.9	1,499
15-19	3.9	0.1	0.0	0.0	0.0	96.0	1,391
20-24	27.1	0.4	0.0	0.2	0.0	72.2	1,128
25-29	58.0	0.9	0.0	0.0	0.0	41.1	754
30-34	83.6	0.4	0.0	0.0	0.0	15.8	495
35-39	94.8	0.2	0.0	0.9	0.0	4.1	466
40-44	98.0	0.8	0.0	0.0	0.3	1.0	400
45-49	98.7	0.0	0.2	0.7	0.2	0.2	447
50-54	99.1	0.3	0.0	0.3	0.0	0.3	324
55-59	97.7	0.0	0.0	2.3	0.0	0.0	349
60-64	97.1	1.0	0.0	1.9	0.0	0.0	208
65-69	95.1	0.5	0.0	4.4	0.0	0.0	182
70-74	90.8	0.0	0.0	9.2	0.0	0.0	109
75-79	84.2	0.0	0.0	15.8	0.0	0.0	76
80-84	83.8	0.0	0.0	16.2	0.0	0.0	37
85+	84.6	0.0	0.0	15.4	0.0	0.0	39
All	47.5	0.3	0.0	0.8	0.0	51.4	7,904
Female							
10-14	0.7	0.0	0.0	0.0	0.0	99.3	1,401
15-19	31.9	0.3	0.1	0.0	0.0	67.7	1,531
20-24	68.9	1.2	0.5	0.4	0.0	28.9	1,008
25-29	87.9	1.8	1.0	1.8	0.0	7.5	603
30-34	91.5	1.3	1.3	2.7	0.4	2.5	555
35-39	89.3	1.4	2.1	7.0	0.0	0.2	559
40-44	86.1	1.6	1.1	10.5	0.0	0.5	438
45-49	76.6	0.9	2.6	19.5	0.0	0.2	431
50-54	73.3	0.8	1.6	23.8	0.0	0.3	378
55-59	60.8	0.7	0.3	36.9	0.3	0.3	301
60-64	43.4	0.0	0.7	55.9	0.0	0.0	145
65-69	40.9	0.0	0.0	57.8	0.6	0.6	154
70-74	24.4	1.2	0.0	74.4	0.0	0.0	82
75-79	17.8	0.0	0.0	80.8	0.0	1.4	73
80-84	3.1	0.0	0.0	96.9	0.0	0.0	32
85+	7.9	0.0	2.6	89.5	0.0	0.0	38
All	52.5	0.8	0.7	9.8	0.1	36.0	7,729

# APPENDIX N

# Chakaria HDSS project team, 2009

Name of Staff	Designation				
Dhaka					
Abbas Bhuiya	Project Director				
Mohammad Iqbal	Senior Operations Researcher				
SM Manzoor Ahmed Hanifi	Assistant Scientist				
Rumesa R Aziz	Research Investigator				
Tania Wahed	Senior Operations Researcher				
Shehrin Shaila Mahmood	Assistant Scientist				
Farhana Urni	Senior Statistical Officer				
Md. Kashem Iqbal	Senior Administrative Officer				
Ayesha Begum	Senior Data Management Assistant				
Chakaria					
Shahidul Hoque	Senior Field Research Officer				
Ariful Moula	Field Research Officer				
Mijanur Rahaman	Field Research Officer				
Ashish Paul	Senior Data Management Assistant				
Md. Sharif Al-Hasan	Field Research Supervisor				
Snehasish Dutta	Field Research Assistant				
Md. Rehmat Ali	Senior Field Assistant				
Afroza Yeasmin	Data Collector				
Armanul Maowa	Data Collector				
Aymun Nahar	Data Collector				
Fatema Johura Surma	Data Collector				
Fatema Zannat	Data Collector				
Helena Khanom Happy	Data Collector				
Hosaina Begum	Data Collector				
Ismat Jahan Khuki	Data Collector				
Kawsar Jannat	Data Collector				
Kawsar Jannat Mukta	Data Collector				
Kulsuma Aktar	Data Collector				
Mina Dhar	Data Collector				
Mobasseratul Zannat	Data Collector				
Monuara Begum	Data Collector				
Nazma Akter	Data Collector				
Nigar Sultana	Data Collector				
Noor Ayesha Begum Rawnak Zahan	Data Collector				
Riasmin Zannat	Data Collector Data Collector				
Rosan Ara	Data Collector  Data Collector				
Sabina Yesmin	Data Collector  Data Collector				
	Data Collector  Data Collector				
Setara Begum Shamima Khanam	Data Collector				
Tanjina Zannat Ara	Data Collector				
Tanjina Zannat Afa Tanjimul Zannat	Data Collector  Data Collector				
Zannatul Ferdous	Data Collector				
Zosna Begum	Data Collector				

